




12-2009

## Higher Order Difference Schemes for Heat Equation

Jianzhong Wang  
*Sam Houston State University*

Follow this and additional works at: <https://digitalcommons.pvamu.edu/aam>

 Part of the [Numerical Analysis and Computation Commons](#), and the [Partial Differential Equations Commons](#)

---

### Recommended Citation

Wang, Jianzhong (2009). Higher Order Difference Schemes for Heat Equation, *Applications and Applied Mathematics: An International Journal (AAM)*, Vol. 4, Iss. 2, Article 2.  
Available at: <https://digitalcommons.pvamu.edu/aam/vol4/iss2/2>

This Article is brought to you for free and open access by Digital Commons @PVAMU. It has been accepted for inclusion in *Applications and Applied Mathematics: An International Journal (AAM)* by an authorized editor of Digital Commons @PVAMU. For more information, please contact [hvkoshy@pvamu.edu](mailto:hvkoshy@pvamu.edu).



## Higher Order Difference Schemes for Heat Equation

**Jianzhong Wang**

Department of Mathematics and Statistics  
Sam Houston State University  
Huntsville, TX 77341 USA  
[mth\\_jxw@shsu.edu](mailto:mth_jxw@shsu.edu)

Received: July 10, 2009; Accepted: October 15, 2009

### Abstract

In this paper, we construct the explicit difference schemes for the heat equation with arbitrary high orders. We also show the validity of the new schemes by numerical simulations.

**Keywords:** Heat equation; explicit difference schemes; numerical solutions; high-order schemes

**MSC (2000) No.:** 65M06 65D25 35K05

### 1. Introduction

In this paper we study the construction of the explicit difference schemes for the 1-D heat equation

$$D_t u = c D_x^2 u, \quad (1)$$

where  $u = u(x, t)$  is temperature,  $c > 0$  is the heat conductivity,  $D_t = \frac{\partial}{\partial t}$ , and  $D_x^2 = \frac{\partial^2}{\partial x^2}$ . The operator form of (1) is

$$D_t = c D_x^2. \quad (2)$$

For a temperature function  $u(x, t)$ , let  $E_h$  ( $h \in R$ ) denote the spatial translation operator  $E_h u(\cdot, t) = u(\cdot + h, t)$ . We have  $E_{kh} = E_h^k, k \in Z$ . A Laurent polynomial of  $E_h$  :

$$A_h^{n,m} = \sum_{k=-m}^n a_k E_{kh}$$

is called a difference scheme of order  $s \in N$  if

$$A_h^{n,m}(x^k) = 0, \quad k = 0, 1, \dots, s-1.$$

A difference scheme of order 2 approximates  $D_x^2$ . Since the differential operator  $D_x^2$  is self-conjugate, we are only interested in the symmetric difference scheme of order 2:

$$A_h^n = \sum_{k=0}^n a_k (E_{kh} + E_{-kh}), \quad \text{with } A_h^n(1) = 0 \text{ and } A_h^n(x) = 0. \tag{3}$$

For a temperature function  $u$ , we define the temporal difference operator  $\Delta_t$  ( $t > 0$ ) by  $\Delta_t u(x, \cdot) = u(x, t + \cdot) - u(x, \cdot)$ . Thus, an explicit difference scheme for the heat equation (1) is

$$\frac{\Delta_t}{t} u = \frac{c}{h^2} A_h^n u. \tag{4}$$

Let  $\lambda = \frac{ct}{h^2} > 0$  be the constant multiple of the ratio of the time-step to the square of the space-step (TSR). Then,  $t = \frac{\lambda h^2}{c} (= O(h^2))$ . A difference scheme (4) is said to have order  $s \in N$  if

$$R(u) = \frac{\Delta_t}{t} u - \frac{c}{h^2} A_h^n u = O(h^s), \quad h \rightarrow 0.$$

Write

$$\delta_h^2 = E_h + E_{-h} - 2I, \tag{5}$$

where  $I$  is the identity operator. The simplest difference scheme for (1) is

$$\frac{\Delta_t}{t} u = \frac{c}{h^2} \delta_h^2 u, \tag{6}$$

which has order 2 and its stability condition is  $\lambda \leq \frac{1}{2}$  [Gerald Wheatley (1999), Richtmyer and Morton (1967)]. People are also interested in higher order difference schemes. In Qian et al. (2000), the authors proposed the following difference scheme of order 4:

$$\frac{\Delta_t}{t} u = \frac{c}{h^2} \delta_h^2 \left( I + \frac{6\lambda - 1}{12} \delta_h^2 \right) u, \tag{7}$$

and showed that when  $\lambda \leq \frac{2}{3}$  the scheme is stable. In this paper, we shall give a general formula for the construction of difference schemes for (1) with arbitrary orders and show the validity of

the formula by numerical simulations.

## 2. Construction of Difference Schemes

We start our construction from the exponential expansion of  $\Delta_t$  :

$$\Delta_t = \sum_{n=1}^{\infty} \frac{t^n D_t^n}{n!}.$$

Applying the heat equation (2) and recalling  $\lambda = \frac{ct}{h^2}$ , we have

$$\Delta_t = \sum_{n=1}^{\infty} \frac{\lambda^n h^{2n} D_x^{2n}}{n!}.$$

To illustrate our method, we first construct the difference schemes for (1) with order 2 and 4, respectively. The Taylor expansion of  $\delta_h^2$  in (5) is

$$\delta_h^2 = 2 \left( \sum_{n=1}^{\infty} \frac{h^{2n} D_x^{2n}}{(2n)!} \right), \quad (8)$$

which yields

$$\begin{aligned} \frac{\Delta_t}{t} - \left( \frac{c}{h^2} \right) \delta_h^2 &= c \sum_{n=1}^{\infty} \left( \frac{\lambda^{n-1}}{n!} - \frac{2}{(2n)!} \right) h^{2(n-1)} D_x^{2n} \\ &= c \sum_{n=1}^{\infty} \left( \frac{\lambda^n}{(n+1)!} - \frac{2}{(2n+2)!} \right) h^{2n} D_x^{2(n+1)}, \end{aligned}$$

i.e.,

$$\frac{\Delta_t}{t} - \left( \frac{c}{h^2} \right) \delta_h^2 = c \left( \frac{\lambda}{2} - \frac{1}{12} \right) h^2 D_x^4 + O(h^4). \quad (9)$$

The formula (9) shows that the simplest scheme (6) has order 2, and it achieves order 4 when  $\lambda = \frac{1}{6}$ .

To derive the difference scheme of order 4, we, replacing  $h$  in (8) by  $2h$ , derive the identity

$$\delta_{2h}^2 = \sum_{n=1}^{\infty} \frac{2^{2n+1} h^{2n} D_x^{2n}}{(2n)!}$$

and set the scheme to

$$\frac{\Delta_t}{t} = \frac{c}{h^2} (a\delta_h^2 + b\delta_{2h}^2), \tag{10}$$

where  $a$  and  $b$  are two real numbers to be determined. We have

$$\frac{\Delta_t}{t} - \frac{c}{h^2} (a\delta_h^2 + b\delta_{2h}^2) = c \sum_{k=1}^{\infty} \left( \frac{\lambda^{k-1}}{k!} - \frac{2}{(2k)!} (a + 4^k b) \right) h^{2(k-1)} D_x^{2k}.$$

Let

$$\begin{cases} a + 4b = 1 \\ a + 4^2 b = 6\lambda. \end{cases} \tag{11}$$

Then,

$$\begin{aligned} \frac{\Delta_t}{t} - \frac{c}{h^2} (a\delta_h^2 + b\delta_{2h}^2) &= c \sum_{k=3}^{\infty} \left( \frac{\lambda^{k-1}}{k!} - \frac{2}{(2k)!} (a + 4^k b) \right) h^{2(k-1)} D_x^{2k} \\ &= c \left( \frac{\lambda^2}{6} - \frac{1}{360} (a + 64b) \right) h^4 D_x^6 + O(h^6). \end{aligned}$$

The solution of (11) is

$$\begin{cases} a = \frac{4}{3} - 2\lambda \\ b = -\frac{1}{12} + \frac{1}{2}\lambda. \end{cases}$$

Therefore, setting  $a = \frac{4}{3} - 2\lambda$  and  $b = -\frac{1}{12} + \frac{1}{2}\lambda$ , we have

$$\frac{\Delta_t}{t} - \frac{c}{h^2} (a\delta_h^2 + b\delta_{2h}^2) = c \left( \frac{\lambda^2}{6} - \frac{1}{12}\lambda + \frac{1}{90} \right) h^4 D_x^6 + O(h^6),$$

i.e., the difference scheme

$$\frac{\Delta_t}{t} = \frac{c}{h^2} \left( \frac{4}{3} - 2\lambda \right) \delta_h^2 + \frac{c}{h^2} \left( -\frac{1}{12} + \frac{1}{2}\lambda \right) \delta_{2h}^2 \tag{12}$$

has order 4. By  $\delta_{2h}^2 = \delta_h^2 \delta_h^2 + 4\delta_h^2$ , we have

$$\begin{aligned} &\frac{c}{h^2} \left( \frac{4}{3} - 2\lambda \right) \delta_h^2 + \frac{c}{h^2} \left( -\frac{1}{12} + \frac{1}{2}\lambda \right) \delta_{2h}^2 \\ &= \frac{c}{h^2} \left[ \left( \frac{4}{3} - 2\lambda \right) + 4 \left( -\frac{1}{12} + \frac{1}{2}\lambda \right) \right] \delta_h^2 + \left( -\frac{1}{12} + \frac{1}{2}\lambda \right) (\delta_h^2)^2 \end{aligned}$$

$$= \frac{c}{h^2} \delta_h^2 \left( I + \frac{6\lambda - 1}{12} \delta_h^2 \right),$$

which shows that the scheme (12) is the same as the scheme (7) obtained in Qian, et al. (2000). To obtain the stability condition for the scheme (12), we denote  $E^t = I + \Delta_t$ , and rewrite the scheme (12) to the form of

$$E^t = I + \lambda \left( \frac{4}{3} - 2\lambda \right) \delta_h^2 + \lambda \left( -\frac{1}{12} + \frac{1}{2} \lambda \right) \delta_{2h}^2 = I + \lambda \delta_h^2 + \lambda \left( -\frac{1}{12} + \frac{1}{2} \lambda \right) (\delta_h^2)^2. \quad (13)$$

Let  $A(\omega)$  be representation of  $E^t$  in the Fourier domain. Then, by (13),

$$\begin{aligned} A(\omega) &= 1 + 2\lambda(\cos h\omega - 1) + 4\lambda \left( -\frac{1}{12} + \frac{1}{2} \lambda \right) (\cos h\omega - 1)^2 \\ &= 1 - 4\lambda \sin^2 \frac{h\omega}{2} + \lambda \left( 8\lambda - \frac{4}{3} \right) \sin^4 \frac{h\omega}{2} \end{aligned}$$

The stability condition of the scheme (12) is  $\max_{\omega \in R} |A(\omega)| \leq 1$ , which leads the stability condition  $\lambda \leq \frac{2}{3}$ .

We now develop the difference scheme for (1) with an arbitrary order. Assume that the difference scheme of order  $2m$  has the form

$$\frac{\Delta_t}{t} = \frac{c}{h^2} \sum_{j=1}^m a_j \delta_{jh}^2, \quad (14)$$

where the coefficient vector  $\mathbf{a} = [a_1, \dots, a_m]^T$  is to be determined. Recall that

$$\frac{\Delta_t}{t} - \frac{c}{h^2} \sum_{k=1}^m a_k \delta_{kh}^2 = c \sum_{k=1}^{\infty} \left( \frac{\lambda^{k-1}}{k!} - \frac{2}{(2k)!} \sum_{j=1}^m j^{2k} a_j \right) h^{2(k-1)} D_x^{2k}.$$

In order to obtain a scheme of order  $2m$ , the real numbers  $a_1, \dots, a_m$  have to satisfy

$$\sum_{j=1}^m j^{2k} a_j = \frac{(2k)!}{2k!} \lambda^{k-1}, \quad k = 1, \dots, m. \quad (15)$$

Write

$$V_m = \begin{bmatrix} 1 & 4 & 9 & \dots & m^2 \\ 1 & 4^2 & 9^2 & \dots & m^4 \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ 1 & 4^m & 9^m & \dots & m^{2m} \end{bmatrix}$$

and  $\mathbf{b} = [1, 6\lambda, \dots, \frac{(2m)!}{2^m} \lambda^{m-1}]^T$ . The matrix form of Equation (15) is

$$V_m \mathbf{a} = \mathbf{b}.$$

Since the Vandermonde matrix  $V_m$  is invertible, Equation (15) has the unique solution

$$\mathbf{a} = V_m^{-1} \mathbf{b}, \quad (16)$$

which yields

$$\frac{\Delta_t}{t} - \frac{c}{h^2} \sum_{k=1}^m a_k \delta_{kh}^2 = c \left( \frac{\lambda^m}{(m+1)!} - \frac{2}{(2m+2)!} \sum_{j=1}^m j^{2m+2} a_j \right) h^{2m} D_x^{2m+2} + O(h^{2m+2}),$$

i.e., the difference scheme (14) with  $a$  in (16) has order  $2m$ .

The scheme (14) can be rewritten to

$$E^t = I + \lambda \sum_{k=1}^m a_k \delta_{kh}^2. \quad (17)$$

Let  $A(\omega)$  be representation of  $E^t$  in the Fourier domain. Then,

$$A(\omega) = 1 - 2\lambda \sum_{k=1}^m a_k (1 - \cos kh\omega)$$

and the stability condition of the scheme (14) is  $\max_{\omega \in R} |A(\omega)| \leq 1$ . Therefore, a sufficient condition for the stability can be obtained by  $a_k \geq 0, k = 1, \dots, m$ , and  $1 - 2\lambda \sum_{k=1}^m a_k \geq 0$ .

As examples, we use (16) to derive the difference schemes of order 6 and 8, respectively.

### Example 1.

Let

$$\begin{cases} a_1 = \frac{3}{2} - \frac{13}{4} \lambda + \frac{5}{2} \lambda^2, \\ a_2 = -\frac{3}{20} + \lambda - \lambda^2, \\ a_3 = \frac{1}{90} - \frac{1}{12} \lambda + \frac{1}{6} \lambda^2, \end{cases}$$

which is the solution of the linear system

$$\begin{aligned}a_1 + 4a_2 + 9a_3 &= 1, \\a_1 + 4^2a_2 + 9^2a_3 &= 6\lambda, \\a_1 + 4^3a_2 + 9^3a_3 &= 60\lambda^2.\end{aligned}$$

Then, the difference scheme

$$\frac{\Delta_t}{t} = \frac{c}{h^2}(a_1\delta_h^2 + a_2\delta_{2h}^2 + a_3\delta_{3h}^2)$$

has order 6. We select  $\lambda$  in the following range

$$0.184\frac{1}{2} - \frac{1}{10}\sqrt{10} \leq \lambda \leq \frac{1}{2} + \frac{1}{10}\sqrt{100.816}$$

so that  $a_1, a_2, a_3$ , and  $[1 - 2\lambda(a_1 + a_2 + a_3)]$  are nonnegative, that ensures the stability of the scheme.

### Example 2.

Let

$$\begin{cases} a_1 = \frac{8}{5} - \frac{61}{15}\lambda + \frac{29}{6}\lambda^2 - \frac{7}{3}\lambda^3, \\ a_2 = -\frac{1}{5} + \frac{169}{120}\lambda - \frac{13}{6}\lambda^2 + \frac{7}{6}\lambda^3, \\ a_3 = \frac{8}{315} - \frac{1}{5}\lambda + \frac{1}{2}\lambda^2 - \frac{1}{3}\lambda^3, \\ a_4 = -\frac{1}{560} + \frac{7}{480}\lambda - \frac{1}{24}\lambda^2 + \frac{1}{24}\lambda^3, \end{cases}$$

which is the solution of the linear system

$$\begin{aligned}a_1 + 4a_2 + 9a_3 + 16a_4 &= 1, \\a_1 + 4^2a_2 + 9^2a_3 + 16^2a_4 &= 6\lambda, \\a_1 + 4^3a_2 + 9^3a_3 + 16^3a_4 &= 60\lambda^2, \\a_1 + 4^4a_2 + 9^4a_3 + 16^4a_4 &= 840\lambda^3.\end{aligned}$$

Then, the difference scheme

$$\frac{\Delta_t}{t} = \frac{c}{h^2}(a_1\delta_h^2 + a_2\delta_{2h}^2 + a_3\delta_{3h}^2 + a_4\delta_{4h}^2)$$



has order 8. To ensure the stability of the scheme, we select  $\lambda$  in the range [0.194,0.955].

### 3. Numerical simulations

To validate our theoretical results, we show a numerical example in this section. For comparison, we set the same initial condition as in Qian, et al. (2000) for the heat equation (1):

$$u(x, 0) = \sin(2k\pi x),$$

and seek for the unit-periodic (with respect to  $x$ ) solution of (1). The exact solution is

$$u(x, t) = e^{-4ck^2\pi^2 t} \sin(2k\pi x).$$

To apply our schemes to the equation, we let  $h > 0$  be the space-step and  $\tau > 0$  be the time-step, where  $h$  is chosen such that  $N = 1/h$  is an integer. The relation of  $\tau$  and  $h$  is given by  $\tau = \frac{\lambda h^2}{c}$ , where  $\lambda$  is chosen from the range of the stability. Let  $\hat{u}_m$  be the numerical solution obtained by the difference scheme of order  $2m$ . We measure the global error of the scheme at  $t = n\tau$  by

$$E_m(t) = \sqrt{\frac{1}{N} \sum_{i=0}^N (\hat{u}_m(ih, t) - u(ih, t))^2}$$

and show the pointwise error by the discrete function

$$Er_m(x, t) = \hat{u}_m(x, t) - u(x, t), \quad x = 0, h, 2h, \dots, Nh.$$

As pointed out in Qian, et al. (2000), the maximal global error  $E_m(t)$  is obtained at  $t_0 = \frac{1}{4ck^2\pi^2}$ , which is independent of schemes. Let  $n_0 = \frac{t_0}{\tau}$ , where  $\tau = \frac{\lambda h^2}{c}$ , which yields

$$n_0 = \text{round}\left(\frac{1}{4\lambda k^2 \pi^2 h^2}\right).$$

Then, the maximal error is obtained after  $n_0$  iterations of the schemes. In our numerical simulations, we set


$$k = 1, c = 0.001, \lambda = 0.5, h = 0.01.$$


Then, the global maximal error is obtained after 500 iterations. The following table presents the maximal global errors for all schemes of order 2, 4, 6, and 8.

Order of scheme	2	4	6	8
Maximal global error	1.7042e-004	4.4859e-008	4.7456e-012	1.4156e-16

**Remark:** The maximal global error of the scheme of order 8 already comes up to the machine epsilon  $2^{-64} = 2.2204e-16$ .

The table shows that the numerical results match the theoretical results very well. The following figures show the pointwise errors of difference schemes of order 2, 4, and 6 respectively.





### ***Acknowledgement***

The author would like to thank the editor and, especially, the reviewers for their valuable comments. The research is supported by NSF Grant of the United States, DMS-07-12925.

### **REFERENCES**

- Gerald, C. F. and Wheatley, P. Q. (1999). Applied Numerical Analysis, 6th ed., Addison Wesley.
- Qian, Y., Chen, H., Zhang, R., and Chen, S. (2000). A new fourth order finite difference scheme for the heat equation. *Communications in Nonlinear Science and Numerical Simulation* **5**, 151–157.
- Richtmyer, R. D. and Morton, K. W. (1967). Difference Methods for Initial-Value Problems, 2nd ed., Wiley, New York.