



6-2020

## On the Asymptotic Stability of a Nonlinear Fractional-order System with Multiple Variable Delays

Yener Altun  
*Van Yuzuncu Yil University*

Cemil Tunç  
*Van Yuzuncu Yil University*

Follow this and additional works at: <https://digitalcommons.pvamu.edu/aam>



Part of the [Ordinary Differential Equations and Applied Dynamics Commons](#)

### Recommended Citation

Altun, Yener and Tunç, Cemil (2020). On the Asymptotic Stability of a Nonlinear Fractional-order System with Multiple Variable Delays, *Applications and Applied Mathematics: An International Journal (AAM)*, Vol. 15, Iss. 1, Article 26.

Available at: <https://digitalcommons.pvamu.edu/aam/vol15/iss1/26>

This Article is brought to you for free and open access by Digital Commons @PVAMU. It has been accepted for inclusion in *Applications and Applied Mathematics: An International Journal (AAM)* by an authorized editor of Digital Commons @PVAMU. For more information, please contact [hvkoshy@pvamu.edu](mailto:hvkoshy@pvamu.edu).



## On the Asymptotic Stability of a Nonlinear Fractional-order System with Multiple Variable Delays

<sup>1</sup>Yener Altun and <sup>2</sup>Cemil Tunç

<sup>1</sup>Department of Business Administration  
Ercis Management Faculty  
Van Yuzuncu Yil University  
65080, Van-Turkey  
[yeneraltun@yyu.edu.tr](mailto:yeneraltun@yyu.edu.tr)

<sup>2</sup>Department of Mathematics  
Faculty of Sciences  
Van Yuzuncu Yil University  
65080, Campus, Van-Turkey  
[cemtunc@yahoo.com](mailto:cemtunc@yahoo.com)

Received: January 1, 2020; Accepted: March 24, 2020

### Abstract

In this paper, we consider a nonlinear differential system of fractional-order with multiple variable delays. We investigate asymptotic stability of zero solution of the considered system. We prove a new result, which includes sufficient conditions, on the subject by means of a suitable Lyapunov functional. An example with numerical simulation of its solutions is given to illustrate that the proposed method is flexible and efficient in terms of computation and to demonstrate the feasibility of established conditions by MATLAB-Simulink.

**Keywords:** Asymptotic stability, Lyapunov functional, fractional-order, variable delay

**MSC 2020 No.:** 34A08; 34A09; 34K20, 34K40

### 1. Introduction

The theory and applications of fractional calculus emerging with various applications in science and engineering fields have attracted great attention of researchers during the

last 20 years. At the same time, we note that fractional differential equations/systems with time-varying delays often occur in many scientific areas such as engineering techniques fields, physics, polymer rheology, neural networks, biophysics, blood flow phenomena, capacitor theory, electrical circuits, signal processing, biology, control theory and etc. (see Agarwal et al. (2010), Altun (2019), Chen et al. (2012), Chen et al. (2014), Deng and Deng (2014), Deng (2010), Diethelm (2010), Duarte-Mermoud et al. (2015), Hristova and Tunç (2019), Hu et al. (2015), Khan et al. (2018), Kilbas et al. (2006), Li et al. (2015), Li et al. (2010), Liu et al. (2017), Liu et al. (2016, 2017), Lu and Chen (2009), Matignon (1996), Podlubny (1999), Qian et al. (2010) and the references therein).

This paper, motivated by the results of Liu et al. (2016) and that can be found in Agarwal et al. (2010), Alidousti (2017), Altun (2019), Altun and Tunç (2019), Chen et al. (2012), Chen et al. (2014), Deng and Deng (2014), Deng (2010), Diethelm (2010), Duarte-Mermoud et al. (2015), Gözen and Tunç (2020), Grace et al. (2019), Graef et al. (2017), Hristova and Tunç (2019), Hu et al. (2015), Khan et al. (2018), Kilbas et al. (2006), Li et al. (2015), Li et al. (2010), Liu et al. (2017), Lu and Chen (2009), Matignon (1996), Podlubny (1999), Qian et al. (2010), Slyn'ko and Tunç (2019), Tan (2008), Tunç and Mohammed (2019), Tunç and Tunç (2016a, 2016b), Wang et al. (2012), Zhou et al. (2014), Zhang et al. (2018) and the sources therein.

## 2. Preliminaries

In this section, several basic definitions and lemmas related to fractional calculus are presented.

The fractional integral (Riemann–Liouville integral)  ${}_t D_t^{-q}$  with fractional-order  $q \in \mathbb{R}^+$  of a function  $x(t)$  is defined by

$${}_t D_t^{-q} x(t) = \frac{1}{\Gamma(q)} \int_{t_0}^t (t-\tau)^{q-1} x(\tau) d\tau,$$

(see Podlubny (1999)).

The fractional derivative (Riemann–Liouville derivative) of fractional-order  $q$  of a function  $x(t)$  is defined by

$${}_t D_t^q x(t) = \frac{1}{\Gamma(n-q)} \frac{d^n}{dt^n} \int_{t_0}^t (t-\tau)^{n-q-1} x(\tau) d\tau, \quad (n-1 \leq q < n),$$

where  $\Gamma(\cdot)$  is the Gamma function, (see Liu et al. (2016)).

If  $p > q > 0$ , then the equality

$${}_t D_t^p ({}_t D_t^{-q} f(t)) = {}_t D_t^{p-q} f(t)$$

holds for sufficiently good functions  $f(t)$ . In particular, this relation holds if  $f(t)$  is integrable (see Hu et al. (2015)).

Suppose  $x(t) \in R^n \rightarrow R^n$  is a vector of differentiable functions. Then, for any  $t \geq t_0$ , the following relationship holds:

$${}_{t_0}D_t^q(x^T(t)Mx(t)) \leq 2x^T(t)M {}_{t_0}D_t^q x(t), \quad 0 < q < 1,$$

where  $M \in R^{n \times n}$  is a constant, square, symmetric and positive semi-definite matrix (see Duarte-Mermoud et al. 2015)).

For any  $x, y \in R^n$ ,  $\alpha > 0$ ,  $\alpha \in R$ , the following inequality holds:

$$2x^T y \leq \alpha x^T x + \frac{1}{\alpha} y^T y,$$

(see Liu et al. (2016)).

Suppose  $U > 0$  and  $V \geq 0$  are real symmetric matrices and  $\varepsilon$  is a positive number. Then,

$$\varepsilon U > V \Leftrightarrow \lambda_{\max}(VU^{-1}) < \varepsilon \Leftrightarrow \lambda_{\max}(U^{-\frac{1}{2}}VU^{-\frac{1}{2}}) < \varepsilon,$$

(see Liu et al. (2016)).

### 3. Stability

Liu et al. (2016) considered the following fractional system with unbounded delay:

$${}_{t_0}D_t^q x(t) = Ax(t) + Bx(t - \tau(t)) + F_1(x(t)) + F_2(x(t - \tau(t))).$$

Liu et al. (2016), applying the Lyapunov second method, sufficient conditions on asymptotic stability of zero solution of this fractional nonlinear system with variable delay is obtained. The advantage of the employed method is that one may directly calculate integer-order derivative of the Lyapunov function used therein.

In this paper, we consider the following fractional-order nonlinear differential system with multiple variable delays:

$${}_{t_0}D_t^q x(t) = A_0 x(t) + G_0(x(t)) + \sum_{i=1}^m A_i x(t - h_i(t)) + \sum_{i=1}^m G_i(x(t - h_i(t))), \quad (1)$$

with the initial conditions

$${}_{t_0}D_t^{-(1-q)} x(t) = \mathcal{G}(t), \quad t \in [-\infty, 0],$$

where  ${}_{t_0}D_t^q$  is the Riemann–Liouville fractional derivative of order  $0 < q < 1$ ,  $t \in R^+$ ,  $R^+ = [0, \infty)$ ,  $x(t) \in R^n$ ,  $x(t)$  is the state vector and  $A_0, A_i \in R^{n \times n}$  are known constant

matrices. The nonlinear functions  $G_i \in R^n$  are continuous and satisfy  $G_i(0) = 0$ . The variable delays  $h_i(t) \geq 0$  are differentiable and they satisfy

$$h_i(t) \leq \delta_i < 1, \quad i = 1, 2, \dots, m, \quad (2)$$

where  $\delta_i \in R$ ,  $\delta_i > 0$ .

Suppose the functions  $G_j(x)$  are the higher order term in  $x$  such that

$$\lim_{\|x\| \rightarrow 0} \frac{\|G_j(x)\|}{\|x\|} = 0, \quad j = 0, 1, \dots, m. \quad (3)$$

It is clear that fractional system (1) includes and improves the fractional system discussed by Liu et al. (2016). The aim of this paper is to generalize and improve the work of Liu et al. (2016). These are the contributions of this paper to the subject and the relevant literature.

Throughout this paper, we use the following notations.  $R^n$  denotes the  $n$ -dimensional Euclidean space;  $R^{n \times n}$  is the set of all  $n \times n$  real matrices;  $\|\cdot\|$  denotes the Euclidean norm;  $L^T$  means the transpose of matrix  $L$ ;  $U$  is symmetric if  $U = U^T$ ;  $H$  is positive definite (or negative definite) if  $\langle Hx, x \rangle > 0$  (or  $\langle Hx, x \rangle < 0$ ) for all  $x \neq 0$ ;  $\|A\| = \sqrt{\lambda_{\max}(A^T A)}$  denotes the spectral norm of matrix  $A$ ;  $\lambda_{\min}(B)$  and  $\lambda_{\max}(B)$  denote the minimal and maximal eigenvalues of matrix  $B$ , respectively.

### Theorem 3.1.

The zero solution of fractional differential equation system (1) is asymptotically stable if there exist a positive-definite matrix  $P \in R^{n \times n}$  and a symmetric positive-definite matrix  $Q \in R^{n \times n}$  such that the following relations hold:

$$\begin{cases} P^T A_0 + A_0^T P + (m+1)Q = 0, \\ \sum_{i=1}^m \frac{1}{\sqrt{1-\delta_i}} \|PA_i\| - \lambda_{\min}(Q) < 0. \end{cases} \quad (4)$$

**Proof:**

We define following Lyapunov functional:

$$V(t) = {}_{t_0} D_t^{q-1} (x^T(t) P x(t)) + \sum_{i=1}^m \int_{t-h_i(t)}^t x^T(s) Q x(s) ds. \quad (5)$$

In view of Lemma 2.1, Lemma 2.2 and the inequality (2), from the time derivative  $V(t)$  along solutions of fractional system (1), we obtain the following:

$$\begin{aligned}
V_{\mathcal{L}}(t) &= {}_{t_0} D_t^q (x^T(t) P x(t)) + m x^T(t) Q x(t) - \sum_{i=1}^m (1 - \delta_i) x^T(t - h_i(t)) Q x(t - h_i(t)) \\
&\leq 2x^T(t) P {}_{t_0} D_t^q (x(t)) + m x^T(t) Q x(t) - \sum_{i=1}^m (1 - \delta_i) x^T(t - h_i(t)) Q x(t - h_i(t)) \\
&= 2x^T(t) P \{A_0 x(t) + G_0(x(t)) + \sum_{i=1}^m A_i x(t - h_i(t)) + \sum_{i=1}^m G_i(x(t - h_i(t)))\} \\
&\quad + m x^T(t) Q x(t) - \sum_{i=1}^m (1 - \delta_i) x^T(t - h_i(t)) Q x(t - h_i(t)) \\
&= x^T(t) (P^T A_0 + A_0^T P + m Q) x(t) + 2x^T(t) P \sum_{i=1}^m A_i x(t - h_i(t)) \\
&\quad + 2x^T(t) P G_0(x(t)) + 2x^T(t) P \sum_{i=1}^m G_i(x(t - h_i(t))), \\
&\quad - \sum_{i=1}^m (1 - \delta_i) x^T(t - h_i(t)) Q x(t - h_i(t)). \tag{6}
\end{aligned}$$

For the terms included in (6), except the first one, using Lemma 2.3, we get the following inequalities, respectively:

$$\begin{aligned}
2 \sum_{i=1}^m x^T(t) P A_i x(t - h_i(t)) &= 2 \sum_{i=1}^m x^T(t) P A_i Q^{-\frac{1}{2}} Q^{\frac{1}{2}} x(t - h_i(t)) \\
&\leq \sum_{i=1}^m \frac{1}{\alpha(1 - \delta_i)} x^T(t) P A_i Q^{-1} A_i^T P^T x(t) \\
&\quad + \sum_{i=1}^m \alpha(1 - \delta_i) x^T(t - h_i(t)) Q x(t - h_i(t)), \tag{7}
\end{aligned}$$

$$2x^T(t) P G_0(x(t)) \leq \frac{1}{\beta} x^T(t) P^2 x(t) + \beta G_0^T(x(t)) G_0(x(t)), \tag{8}$$

$$\begin{aligned}
2x^T(t) P \sum_{i=1}^m G_i(x(t - h_i(t))) &\leq \frac{1}{\mu} x^T(t) m P^2 x(t) \\
&\quad + \mu \sum_{i=1}^m G_i^T(x(t - h_i(t))) G_i(x(t - h_i(t))), \tag{9}
\end{aligned}$$

where  $\alpha$ ,  $\beta$  and  $\mu$  are some positive constants.

If we gather the inequalities (7)-(9) into (6) and use the equality in (4), then it follows that:

$$\begin{aligned}
V_{\mathcal{L}}(t) &\leq x^T(t) \left[ P \left( \sum_{i=1}^m \frac{1}{\alpha(1 - \delta_i)} A_i Q^{-1} A_i^T \right) P^T - Q + \left( \frac{1}{\beta} + \frac{m}{\mu} \right) P^2 \right] x(t) \\
&\quad + \sum_{i=1}^m (\alpha - 1)(1 - \delta_i) x^T(t - h_i(t)) Q x(t - h_i(t)) + \beta G_0^T(x(t)) G_0(x(t))
\end{aligned}$$

$$+ \mu \sum_{i=1}^m G_i^T(x(t-h_i(t)))G_i(x(t-h_i(t))). \quad (10)$$

By the definition of the spectral norm, we derive that

$$\begin{aligned} & \left[ \lambda_{\max} \left( Q^{-\frac{1}{2}} P \left( \sum_{i=1}^m \frac{1}{1-\delta_i} A_i Q^{-1} A_i^T \right) P^T Q^{-\frac{1}{2}} \right) \right]^{\frac{1}{2}} \\ &= \left[ \sum_{i=1}^m \frac{1}{1-\delta_i} \lambda_{\max} \left( Q^{-\frac{1}{2}} P A_i Q^{-1} A_i^T P^T Q^{-\frac{1}{2}} \right) \right]^{\frac{1}{2}} \\ &= \left( \sum_{i=1}^m \frac{1}{1-\delta_i} \left\| Q^{-\frac{1}{2}} P A_i Q^{-\frac{1}{2}} \right\|^2 \right)^{\frac{1}{2}} \\ &\leq \sum_{i=1}^m \frac{1}{\sqrt{1-\delta_i}} \| Q^{-\frac{1}{2}} \|^2 \| P A_i \| \\ &= \frac{1}{\lambda_{\min}(Q)} \sum_{i=1}^m \frac{1}{\sqrt{1-\delta_i}} \| P A_i \|. \end{aligned} \quad (11)$$

In view of the inequality given by (4), it follows that

$$\lambda_{\max} \left( Q^{-\frac{1}{2}} P \left( \sum_{i=1}^m \frac{1}{1-\delta_i} A_i Q^{-1} A_i^T \right) P^T Q^{-\frac{1}{2}} \right) < 1. \quad (12)$$

Hence, from (12), it is seen that there exists a positive constant  $\varepsilon$  such that

$$\lambda_{\max} \left( Q^{-\frac{1}{2}} P \left( \sum_{i=1}^m \frac{1}{1-\delta_i} A_i Q^{-1} A_i^T \right) P^T Q^{-\frac{1}{2}} \right) < \varepsilon < 1. \quad (13)$$

Since  $A_i Q^{-1} A_i^T \geq 0$ , owing to Lemma 2.4, it follows from (9) that

$$P \left( \sum_{i=1}^m \frac{1}{1-\delta_i} A_i Q^{-1} A_i^T \right) P^T < \varepsilon Q. \quad (14)$$

Let  $\alpha < 1$  such that  $0 < \frac{\varepsilon}{\alpha} < 1$ . Hence, in view of (14), we have

$$P \left( \sum_{i=1}^m \frac{1}{\alpha(1-\delta_i)} A_i Q^{-1} A_i^T \right) P^T - Q < \left( \frac{\varepsilon}{\alpha} - 1 \right) Q < 0. \quad (15)$$

Let us choose suitable positive constants  $\beta$  and  $\mu$  such that the following inequality holds:

$$K_0 = P \left( \sum_{i=1}^m \frac{1}{\alpha(1-\delta_i)} A_i Q^{-1} A_i^T \right) P^T - Q + \left( \frac{1}{\beta} + \frac{m}{\mu} \right) P^2 < 0. \quad (16)$$

Since  $\alpha < 1$ ,  $0 < \delta_i < 1$  and  $Q$  is positive, then it follows that

$$(\alpha - 1)(1 - \delta_i)Q < 0, \quad i = 1, 2, \dots, m.$$

Let

$$K_i = (\alpha - 1)(1 - \delta_i)Q. \quad (17)$$

Hence, from (10)-(17), we obtain the following inequality:

$$\begin{aligned} \mathcal{V}(t) \leq & x^T(t)K_0x(t) + \sum_{i=1}^m x^T(t - h_i(t))K_ix(t - h_i(t)) \\ & + \beta \|G_0(x(t))\|^2 + \mu \sum_{i=1}^m \|G_i(x(t - h_i(t)))\|^2. \end{aligned} \quad (18)$$

By noting the inequalities (16) and (17), we can choose a constant  $\sigma > 0$  such that

$$K_i + \sigma I < 0, \quad i = 0, 1, 2, \dots, m. \quad (19)$$

From (7), it follows that there exists a constant  $\mathcal{G}(\sigma, \beta, \mu) > 0$  such if  $\|x(t)\| < \mathcal{G}$ ,  $t \geq t_0$ , then the following two inequalities hold:

$$\|G_0(x(t))\|^2 \leq \frac{\sigma}{\beta} \|x(t)\|^2, \quad (20)$$

$$\sum_{i=1}^m \|G_i(x(t - h_i(t)))\|^2 \leq \frac{\sigma}{\mu} \sum_{i=1}^m \|x(t - h_i(t))\|^2. \quad (21)$$

Substituting (20) and (21) into (18), we obtain

$$\mathcal{V}(t) \leq x^T(t)(K_0 + \sigma I)x(t) + \sum_{i=1}^m x^T(t - h_i(t))(K_i + \sigma I)x(t - h_i(t)).$$

Taking into account the inequality (19), we can conclude that  $\mathcal{V}(t)$  is negative definite. Therefore, the zero solution of nonlinear fractional system (1) is asymptotically stable. This result completes the proof of the Theorem 3.1.

### Remark.

Let us consider nonlinear fractional differential system (1) for the scalar linear constant coefficients, when  $A_0 = a_0$ ,  $a_0 \in \mathfrak{R}$ ,  $G_0 = 0$ ,  $A_i = 0$ ,  $G_i = 0$  and  $q = 1$ . In this case, we have the following scalar linear differential equation of first order:

$$\frac{dx}{dt} = a_0x.$$

It is well known that all solutions of this equation are asymptotically stable if and only if  $a_0 < 0$ . This condition coincides with the condition (4) of Theorem 3.1. Here, we would not like to give the details of the discussion for the sake of the brevity.

We now give a numerical example to analyze the stability a fractional system with variable delay. Hence, we see the efficiency and applicability of the proposed method.

### Example 3.2.

In a special case of system (1), we consider the following fractional-order nonlinear system with variable delay:

$${}_{t_0}D_t^q x(t) = A_0 x(t) + A_1 x(t - h_1(t)) + G_0(x(t)) + G(x(t - h_1(t))), \quad (22)$$


where

$$q \in (0, 1), \quad x(t) = [x_1(t), x_2(t)]^T, \quad A_0 = \begin{bmatrix} -3 & 0 \\ 0 & -6 \end{bmatrix}, \quad A_1 = \begin{bmatrix} 0.1 & 0.2 \\ 0 & 1.1 \end{bmatrix},$$

$$h_1(t) = 0.25t, \quad G_0(x) = [x_1 \cos x_2, x_2 \sin x_1]^T,$$

$$G(x(t - h_1(t))) = [0.2x_1(t - h_1(t))e^{-x_1^2(t - h_1(t))}, 0.1x_2(t - h_1(t))e^{-x_2^2(t - h_1(t))}]^T.$$

Since  $\dot{h}_1(t) = 0.25$ , it is clear that  $\delta_1 = 0.25$ . Let  $Q = 3I_2$ . Then, it follows from condition (3.8) that  $P = \begin{bmatrix} 1 & 0 \\ 0 & 0.5 \end{bmatrix}$ . Next, a simple calculation yields that  $\|PA_1\| = 0.55$  and  $\sqrt{1 - \delta_1} \lambda_{\min}(Q) = 2.5980$ . Thus, the condition (4) holds. That is, the trivial solution of fractional-order nonlinear system (22) is asymptotic stable.



**Figure 1.** Numeric simulation for the asymptotic stability of solutions of equation (22) for  $h_1(t) = 0.25t$ .

## 4. Conclusions

In this paper, we derive certain new sufficient conditions on the asymptotic stability of zero solution of a nonlinear fractional system with multiple variable delays. We prove a new theorem on the subject by constructing a meaningful Lyapunov functional and using matrix inequalities. The proposed method avoids computing Riemann-Liouville fractional-order derivative of the given Lyapunov functional. When we compare our

criteria with the stability criteria can be found in the relevant literature, it is seen that our criteria are simple and suitable for applications. Further, a numerical example is given with numerical simulation (see Figure 1) to demonstrate the effectiveness of the stability criteria of the considered system.

## REFERENCES

- Agarwal, R. P., Benchohra, M. and Hamani, S. (2010). A survey on existence results for boundary value problems of nonlinear fractional differential equations and inclusions, *Acta Appl. Math.*, Vol. 109, No. 3, pp. 973–1033.
- Alidousti, J., Ghaziani, R. K. and Eshkaftaki, A. B. (2017). Stability analysis of nonlinear fractional differential order systems with Caputo and Riemann–Liouville derivatives, *Turk. J. Math.*, Vol. 41, No.5, pp. 1260–1278.
- Altun, Y. (2019). Further results on the asymptotic stability of Riemann–Liouville fractional neutral systems with variable delays, *Adv. Difference Equ.*, Vol. 2019, No. 437, pp. 1–13.
- Altun, Y. and Tunç, C. (2019). On exponential stability of solutions of nonlinear neutral differential systems with discrete and distributed variable lags, *Nonlinear Studies*, Vol. 26, No. 2, pp. 455–466.
- Chen F., Nieto J. J. and Zhou Y. (2012). Global attractivity for nonlinear fractional differential equations, *Nonlinear Anal. Real World Appl.*, Vol. 13, No.1, pp. 287–298.
- Chen, L., He, Y., Chai, Y. and Wu, R. (2014). New results on stability stabilization of a class of nonlinear fractional-order systems, *Nonlinear Dynam.*, Vol. 75, No.4, pp. 633–641.
- Deng, J. and Deng, Z. (2014). Existence of solutions of initial value problems for nonlinear fractional differential equations, *Appl. Math. Lett.*, Vol. 32, pp. 6–12.
- Deng, W. (2010). Smoothness and stability of the solutions for nonlinear fractional differential equations, *Nonlinear Anal.*, Vol. 72, No.3-4, pp. 1768–1777.
- Diethelm, K. (2010). *The analysis of fractional differential equations: An application-oriented exposition using differential operators of Caputo type*, Springer, Berlin, Germany.
- Duarte-Mermoud, M.A., Aguila-Camacho, N., Gallegos, J.A. and Castro-Linares, R. (2015). Using general quadratic Lyapunov functions to prove Lyapunov uniform stability for fractional order systems, *Commun. Nonlinear Sci. Numer. Simul.*, Vol. 22, No. 1-3, pp. 650–659.
- Gözen, M. and Tunç, C. (2020). A new result on exponential stability of a linear differential system of first order with variable delays, *Nonlinear Studies*, Vol. 27, No.1, 275–284.
- Grace, S. R., Graef, J. R. and Tunç, E. (2019). On the boundedness of nonoscillatory solutions of certain fractional differential equations with positive and negative terms. *Appl. Math. Lett.*, Vol. 97, pp.114–120.
- Graef, J. R., Grace, S. R. and Tunç, E. (2017). Asymptotic behavior of solutions of nonlinear fractional differential equations with Caputo-type Hadamard derivatives. *Fract. Calc. Appl. Anal.*, Vol. 20, no. 1, pp. 71–87.

- Hristova, S., and Tunç, C. (2019). Stability of nonlinear Volterra integro-differential equations with Caputo fractional derivative and bounded delays, *Electron. J. Differential Equations*, Vol. 2019, No. 30, pp. 1-11.
- Hu, J. B., Lu, G. P., Zhang, S. B. and Zhao, L. D. (2015). Lyapunov stability theorem about fractional system without and with delay, *Commun. Nonlinear Sci. Numer. Simul.*, Vol. 20, No.3, pp. 905–913.
- Khan, H., Tunç, C., Chen, W. and Khan, A. (2018). Existence theorems and Hyers-Ulam stability for a class of Hybrid fractional differential equations with p-Laplacian operator, *J. Appl. Anal. Comput.*, Vol. 8, No. 4, pp. 1211-1226.
- Kilbas, A. A., Srivastava, H. M. and Trujillo, J. J. (2006). *Theory and Application of Fractional Differential Equations*, Elsevier, New York, USA.
- Li, H., Zhou, S. and Li, H. (2015). Asymptotic stability analysis of fractional-order neutral systems with time delay, *Adv. Difference Equ.*, Vol. 2015, No. 325, pp.1–11.
- Li, Y., Chen, Y. and Podlubny, I. (2010). Stability of fractional-order nonlinear dynamic systems: Lyapunov direct method and generalized Mittag-Leffler stability, *Comput. Math. Appl.*, Vol. 59, No. 5, pp. 1810–1821.
- Liu, S., Wu, X. Zhang, Y. J. and Yang, R. (2017). Asymptotical stability of Riemann–Liouville fractional neutral systems, *Appl. Math. Lett.*, Vol. 69, pp. 168–173.
- Liu, S., Wu, X., Zhou, X. F. and Jiang, W. (2016). Asymptotical stability of Riemann–Liouville fractional nonlinear systems, *Nonlinear Dynam.*, Vol. 86, No.1, pp. 65–71.
- Liu, S., Zhou, X. F., Li, X. and Jiang, W. (2017). Asymptotical stability of Riemann–Liouville fractional singular systems with multiple time-varying delays, *Appl. Math. Lett.*, Vol. 65, pp. 32–39.
- Lu, J. G. and Chen, G. (2009). Robust stability and stabilization of fractional-order interval systems: An LMI approach, *IEEE Trans. Automat. Control*, Vol. 54, No. 6, pp. 1294–1299.
- Matignon, D. (1996). *Stability results on fractional differential equations with applications to control processing*, In: *Proceedings of IMACS-SMC, Lille, France*.
- Podlubny, I. (1999). *Fractional Differential Equations*, Academic Press., New York, USA.
- Qian, D., Li, C., Agarwal, R. P. and Wong, P. J. Y. (2010). Stability analysis of fractional differential system with Riemann–Liouville derivative, *Math. Comput. Modelling*, Vol. 52, No. 5-6, pp. 862–874.
- Slyn'ko, V., and Tunç, C. (2019). Stability of abstract linear switched impulsive differential equations, *Automatica J. IFAC*, Vol. 107, pp. 433–441.
- Tan, M. C. (2008). Asymptotic stability of nonlinear systems with unbounded delays, *J. Math. Anal. Appl.*, Vol. 337, pp. 1010–1021.
- Tunç, C. and Mohammed, S.A. (2019). On the asymptotic analysis of bounded solutions to nonlinear differential equations of second order, *Adv. Difference Equ.*, Vol. 2019, No. 461, pp. 1-19.
- Tunç, C. and Tunç, O. (2016). On the boundedness and integration of non-oscillatory solutions of certain linear differential equations of second order, *Journal of Advanced Research*, Vol. 7, No. 1, pp. 165-168.
- Tunç, E. and Tunç, O. (2016). On the oscillation of a class of damped fractional differential equations. *Miskolc Math. Notes*, Vol. 17 , no. 1, pp. 647–656.

- Wang, J., Lv, L. and Zhou Y. (2012). New concepts and results in stability of fractional differential equations, *Commun Nonlinear Sci. Numer. Simul.*, Vol. 17, No. 6, pp. 2530-2538.
- Zhou, X. F., Hu, L. G., Liu, S. and Jiang, W. (2014). Stability criterion for a class of nonlinear fractional differential systems, *Appl. Math. Lett.*, Vol. 28, pp. 25–29.
- Zhang, H., Ye, R., Cao, J., Ahmed, A., Li, X. and Ying, W. (2018). Lyapunov functional approach to stability analysis of Riemann-Liouville fractional neural networks with time-varying delays, *Asian J. Control*, Vol. 20 , no.5, 1938–1951.