



12-2010

Application of Homotopy Perturbation Method to Biological Population Model

Pradip Roul
Duisburg-Essen University

Follow this and additional works at: <https://digitalcommons.pvamu.edu/aam>



Part of the [Numerical Analysis and Computation Commons](#), and the [Partial Differential Equations Commons](#)

Recommended Citation

Roul, Pradip (2010). Application of Homotopy Perturbation Method to Biological Population Model, *Applications and Applied Mathematics: An International Journal (AAM)*, Vol. 5, Iss. 2, Article 2.
Available at: <https://digitalcommons.pvamu.edu/aam/vol5/iss2/2>

This Article is brought to you for free and open access by Digital Commons @PVAMU. It has been accepted for inclusion in *Applications and Applied Mathematics: An International Journal (AAM)* by an authorized editor of Digital Commons @PVAMU. For more information, please contact hvkoshy@pvamu.edu.



Available at
<http://pvamu.edu/aam>
Appl. Appl. Math.
ISSN: 1932-9466

Applications and Applied
Mathematics:
An International Journal
(AAM)

Vol. 05, Issue 2 (December 2010), pp. 272 – 281
(Previously, Vol. 5, Issue 10, pp. 1369 – 1378)

Application of Homotopy Perturbation Method to Biological Population Model

Pradip Roul

Institute for Numerical Mathematics
Duisburg-Essen University
D-47058, Duisburg, Germany
bapuroul@yahoo.com

Received: May 18, 2010; Accepted: August 2, 2010

Abstract

In this article, a well-known analytical approximation method, so-called the Homotopy perturbation method (HPM) is adopted for solving the nonlinear partial differential equations arising in the spatial diffusion of biological populations. The resulting solutions are compared with those of the existing solutions obtained by employing the Adomian's decomposition method. The comparison reveals that our approximate solutions are in very good agreement with the solutions by Adomian's method. Moreover, the results show that the proposed method is a more reliable, efficient and convenient one for solving the non-linear differential equations.

Keywords: Homotopy perturbation method (HPM); Adomian's decomposition method; biological population model; exact solution

MSC 2000 No.: 35K15, 35C05, 65D99, 65M99

1. Introduction

Many problems arising in scientific fields including mathematical biology, fluid dynamics, visco-elasticity and mathematical physics can be successfully modeled by the use of nonlinear partial differential equations. Several approaches such as the Tanh method [Evans (2005), Fan (2000)], the Homotopy-Perturbation method [He (1999a, 2000a, 2005), Janalizadeh (2008), Yildirim (2010a, 2010b), Koçak (2009), Ganji (2006)], the Adomian's decomposition method

[Adomian (1988,1994), Gorguis (2006), Momani (2005), Shakeri (2007)], the Variational Iteration Method [(He (1999b, 2000b, 2006), Mohyud-Din (2009), Abdou (2005), Xu (2009), Noor (2008), Shakeri (2007))], and some asymptotic methods [He (2006)] have been used to solve either linear or nonlinear differential equations. Among these methods, the variational iteration method and the homotopy perturbation method are the most efficient, convenient and effective methods for finding the analytical approximate solutions of nonlinear problems.

In this article, the nonlinear degenerate parabolic equations arise in the study of spatial diffusion of biological populations subject to some initial conditions. Consider the first-order time derivative, nonlinear biological model in the following form:

$$\frac{\partial u}{\partial t} = \frac{\partial^2 u^2}{\partial x^2} + \frac{\partial^2 u^2}{\partial y^2} + g(u), \quad (1)$$

with given initial condition $u(x, y, 0)$, where u and g denote the population density and population supply due to births and deaths, respectively.

In this study, g is considered as $g(u) = hu^k(1 - ru^d)$, where h, d, k, r are real numbers. It is worth pointing out that there are two examples of constitutive equations for $g(u)$: if $h = c$, $d = 1$, $r = 0$, this leads to Malthusian law [Gurtin (1977)], where c is a constant and $h = c_1, d = k = 1$, $r = c_2/c_1$, Verhulst law [Gurtin (1977)], c_1, c_2 are positive constants.

The purpose of this paper is to extend the homotopy perturbation method for computing the approximate analytical solutions of the nonlinear biological population model and then see how these solutions compare with the available exact solutions implemented by Shakeri et al. [Shakeri (2007)] adopting an Adomian decomposition method. The homotopy perturbation method was originally proposed by He for nonlinear differential problems [He (1999a, 2000a)]. It's main feature is the condition of homotopy by introducing an embedment parameter p , which takes the value from 0 to 1. If $p = 0$, the system of equations (homotopy equations) generally reduces to a very simplified form, which yields a rather simple solution. On the other hand, when $p = 1$, it turns out to be the original problem and provides the required solution. The approximate solutions obtained using the HPM converges rapidly to the exact solution without any restrictive assumptions, linearization or transformations. In contrast to the traditional perturbation method [Holemen (1995), Nayfeh (2000)], the HPM method does not need a small parameter in the system. The detailed description of HPM is described in Section 2.

This article is organized as follows. Section 2 is devoted to a short description of the analysis of homotopy perturbation method. In Section 3, we present the analytical approximate solutions obtained by implementing the HPM to the nonlinear biological population model followed by the comparison of results between the approximate solutions and the solutions obtained by adopting an Adomian decomposition method. Finally in Section 4, we summarize and discuss the results.

2. Analysis of Homotopy Perturbation Method

To illustrate the basis ideas of the homotopy-perturbation method, we consider the following nonlinear differential equation:

$$Lu(\tau) + Nu(\tau) + g(\tau) = 0, \quad (2)$$

with boundary conditions

$$B\left(u, \frac{\partial u}{\partial n}\right) = 0, \quad \tau \in \Gamma, \quad (3)$$

where L is a linear operator, N denotes a non-linear operator, and $g(\tau)$ is a known analytical function, $u(\tau)$ is an unknown function, Γ is the boundary of the domain Ω and B is a boundary operator.

By means of Homotopy perturbation method [He (1999a, 2000a)], we can construct a Homotopy $v(\tau, p)$ for Equation (2) as follows:

$$v(\tau, p) : \Omega \times [0, 1] \rightarrow R,$$

which satisfies

$$H(v(\tau), p) = (1-p)(Lv(\tau) - Lu_0(\tau)) + p(Lv(\tau) + Nv(\tau) + g(\tau)), \quad (4)$$

or

$$H(v(\tau), p) = Lv(\tau) - Lu_0(\tau) + p(Lu_0(\tau) + Nv(\tau) + g(\tau)), \quad (5)$$

where $p \in [0, 1]$ is an embedding parameter, $v(\tau)$ is an unknown function and u_0 is an initial approximation that must satisfy the boundary condition (3).

If $p = 0$, then Equation (5) becomes

$$H(v(\tau), 0) = Lv(\tau) - Lu_0(\tau) = 0 \quad (6)$$

and when $p = 1$, Equation (5) takes the original form of the Equation (2), i.e.,

$$H(v(\tau), 1) = Lv(\tau) + Nv(\tau) + g(\tau) = 0. \quad (7)$$

To solve the problem (2), it is necessary to use the perturbation technique as discussed in Hólemen (1995) and Nayfeh (2000). So the combination of the perturbation method and the

homotopy method is known as the homotopy-perturbation method. By applying the perturbation technique the solution of Equation (2) can be expressed as a power series in p

$$v = v_0 + pv_1 + p^2v_2 + p^3v_3 + \dots \quad (8)$$

Letting $p = 1$ in Equation (8), the approximate solution of (2) can be obtained easily as follows

$$u = \lim_{p \rightarrow 1} (v_0 + pv_1 + p^2v_2 + p^3v_3 + \dots) = v_0 + v_1 + v_2 + v_3 + \dots \quad (9)$$

The detailed convergence analysis of the HPM has been discussed in He (1999a, 2000a). The rate of convergence of power series (8), stated in He (1999a), depends upon the nonlinear operator of Equation (2) which satisfies the following two conditions:

- I. The second derivative of nonlinear operator $Nv(\tau)$ must be small enough, otherwise the parameter may be large, that is, p approaches to 1.
- II. The norm of $L^{-1} \frac{\partial N}{\partial v}$ must be smaller than 1.

3. Application of Homotopy Perturbation Method

In this section, the homotopy perturbation method described in the previous section for solving three different types of problems arising in biological population models is applied. Then comparison is made with the available analytical results obtained by Shakeri et al. (2007) using the Adomian's decomposition method to assess the accuracy and the effectiveness of the homotopy perturbation method.

Example 1.

Let us consider the following biological population model:

$$u_t(x, y, t) = \frac{\partial^2 u^2(x, y, t)}{\partial x^2} + \frac{\partial^2 u^2(x, y, t)}{\partial y^2} - u(x, y, t) \left(1 + \frac{8}{9} u(x, y, t) \right), \quad (10)$$

$$\text{subject to the initial condition } u(x, y, 0) = \exp\left(\frac{1}{3}(x + y)\right). \quad (11)$$

According to the homotopy perturbation method, we can construct the homotopy for Equation (10) which satisfies:

$$(1-p) \left(\frac{\partial u(x,y,t)}{\partial t} - \frac{\partial u_0(x,y,t)}{\partial t} \right) + p \left(\frac{\partial u(x,y,t)}{\partial t} - \frac{\partial^2 u^2(x,y,t)}{\partial x^2} - \frac{\partial^2 u^2(x,y,t)}{\partial y^2} + u(x,y,t) \left(1 + \frac{8}{9} u(x,y,t) \right) \right). \quad (12)$$

Substituting (8) into (12) and equating the terms with identical powers of p , we obtain the following set of linear partial differential equations

$$p^0: \frac{\partial u_0(x,y,t)}{\partial t} = 0, \quad u_0(x,y,t) = \exp\left(\frac{1}{3}(x+y)\right). \quad (13)$$

$$p^1: \frac{\partial u_1(x,y,t)}{\partial t} - \frac{\partial^2 u_0^2(x,y,t)}{\partial x^2} - \frac{\partial^2 u_0^2(x,y,t)}{\partial y^2} + u_0(x,y,t) + \frac{8}{9} u_0^2(x,y,t) = 0, \quad u_1(x,y,0) = 0, \quad (14)$$

$$p^2: \frac{\partial u_2(x,y,t)}{\partial t} - 2 \frac{\partial^2 (u_0(x,y,t)u_1(x,y,t))}{\partial x^2} - 2 \frac{\partial^2 (u_0(x,y,t)u_1(x,y,t))}{\partial y^2} + u_1(x,y,t) + \frac{16}{9} u_0(x,y,t)u_1(x,y,t) = 0, \quad (15)$$

⋮

Solving the above equations, we obtain the following approximations

$$u_0(x,y,t) = \exp\left(\frac{1}{3}(x+y)\right), \quad (16)$$

$$u_1(x,y,t) = -t \exp\left(\frac{1}{3}(x+y)\right), \quad (17)$$

$$u_2(x,y,t) = \frac{t^2}{2} \exp\left(\frac{1}{3}(x+y)\right), \quad (18)$$

⋮

and so on, in the same manner the rest of the components can be obtained using the Maple package.

According to the HPM, we can obtain the solution in a series form as follows

$$u(x,y,t) = \exp\left(\frac{1}{3}(x+y)\right) - t \exp\left(\frac{1}{3}(x+y)\right) + \frac{t^2}{2!} \exp\left(\frac{1}{3}(x+y)\right) - \dots$$

$$\begin{aligned}
 &= \exp\left(\frac{1}{3}(x+y)\right)\left(1-t+\frac{t^2}{2!}-\dots\right) \\
 &= \exp\left(\frac{1}{3}(x+y)\right)\sum_{n=0}^{\infty}\frac{(-1)^n t^n}{n!},
 \end{aligned} \tag{19}$$

which has the exact solution

$$u(x, y, t) = \exp\left(\frac{1}{3}(x+y)\right)\exp(-t) = \exp\left(\frac{1}{3}(x+y)-t\right). \tag{20}$$

From the above solution process, it can be seen clearly that, the approximate solution converges very fast to its exact solution. The solution in Equation (20) which obtained by HPM is absolutely same as that of the solution investigated by Shkeri et al. [Shakeri (2007)] using the Adomian decomposition method. Furthermore, the main advantage in using the HPM for solving the considered model is that the exact solutions obtained successfully without requiring a small parameter in the equation and without calculating the complicated Adomian’s polynomials.

Example 2.

Let us consider the following biological population model:

$$u_t(x, y, t) = \frac{\partial^2 u^2(x, y, t)}{\partial x^2} + \frac{\partial^2 u^2(x, y, t)}{\partial y^2} + hu(x, y, t), \tag{21}$$

with the initial condition $u(x, y, 0) = \sqrt{xy}$.

Similarly, by using the homotopy perturbation method, a homotopy of (21) can be obtained as follows

$$(1-p)\left(\frac{\partial u(x, y, t)}{\partial t} - \frac{\partial u_0(x, y, t)}{\partial t}\right) + p\left(\frac{\partial u(x, y, t)}{\partial t} - \frac{\partial^2 u^2(x, y, t)}{\partial x^2} - \frac{\partial^2 u^2(x, y, t)}{\partial y^2} - hu(x, y, t)\right). \tag{22}$$

Substituting (8) into (22) and equating the terms with identical powers of p , we obtain the following set of linear partial differential equations

$$p^0: \frac{\partial u_0(x, y, t)}{\partial t} = 0, \quad u_0(x, y, t) = \sqrt{xy}, \tag{23}$$

$$p^1: \frac{\partial u_1(x, y, t)}{\partial t} - \frac{\partial^2 u_0^2(x, y, t)}{\partial x^2} - \frac{\partial^2 u_0^2(x, y, t)}{\partial y^2} - hu_0(x, y, t) = 0, \quad u_1(x, y, 0) = 0, \tag{24}$$

$$p^2 : \frac{\partial u_2(x, y, t)}{\partial t} - 2 \frac{\partial^2 (u_0(x, y, t)u_1(x, y, t))}{\partial x^2} - 2 \frac{\partial^2 (u_0(x, y, t)u_1(x, y, t))}{\partial y^2} - hu_1(x, y, t) = 0,$$

$$u_2(x, y, 0) = 0, \quad (25)$$

⋮

Using the initial approximation $u_0(x, y, t) = \sqrt{xy}$, and solving the above equations, we obtain the approximations as follows

$$u_0(x, y, t) = \sqrt{xy}, \quad (26)$$

$$u_1(x, y, t) = h\sqrt{xy}t, \quad (27)$$

$$u_2(x, y, t) = h^2\sqrt{xy}\frac{t^2}{2!}, \quad (28)$$

⋮

$$u_n(x, y, t) = h^n\sqrt{xy}\frac{t^n}{n!}. \quad (29)$$

Inserting the values of $u_0, u_1, u_2, \dots, u_n$ in Equation (9), yields the exact solution of (21) as follows

$$u(x, y, t) = \lim_{n \rightarrow \infty} \left(\sqrt{xy} + h\sqrt{xy}t + h^2\sqrt{xy}\frac{t^2}{2!} + \dots + h^n\sqrt{xy}\frac{t^n}{n!} \right)$$

$$= \lim_{n \rightarrow \infty} \sqrt{xy} \sum_{n=0}^{\infty} \frac{(ht)^n}{n!} = \sqrt{xy}e^{ht}, \quad (30)$$

which is the same exact solution obtained by Shakeri et al. (2007) using the Adomian decomposition method, if we use the parameter $h = 1/5$.

Example 3.

Consider the following biological population model:

$$u_t(x, y, t) = \frac{\partial^2 u^2(x, y, t)}{\partial x^2} + \frac{\partial^2 u^2(x, y, t)}{\partial y^2} + u(x, y, t), \quad (31)$$

subject to the initial condition $u(x, y, 0) = \sqrt{\sin x \sinh y}$.

For solving (31) by the homotopy-perturbation technique we consider the following homotopy

$$(1-p) \left(\frac{\partial u(x,y,t)}{\partial t} - \frac{\partial u_0(x,y,t)}{\partial t} \right) + p \left(\frac{\partial u(x,y,t)}{\partial t} - \frac{\partial^2 u^2(x,y,t)}{\partial x^2} - \frac{\partial^2 u^2(x,y,t)}{\partial y^2} u(x,y,t) \right). \quad (32)$$

Substituting the value of u from Equation (8) into (32) and equating the terms of the same powers of p , it yields that

$$p^0: \frac{\partial u_0(x,y,t)}{\partial t} = 0, \quad u_0(x,y,t) = \sqrt{\sin x \sinh y}. \quad (33)$$

$$p^1: \frac{\partial u_1(x,y,t)}{\partial t} - \frac{\partial^2 u_0^2(x,y,t)}{\partial x^2} - \frac{\partial^2 u_0^2(x,y,t)}{\partial y^2} - u_0(x,y,t) = 0, \quad u_1(x,y,0) = 0, \quad (34)$$

$$p^2: \frac{\partial u_2(x,y,t)}{\partial t} - 2 \frac{\partial^2 (u_0(x,y,t)u_1(x,y,t))}{\partial x^2} - 2 \frac{\partial^2 (u_0(x,y,t)u_1(x,y,t))}{\partial y^2} - u_1(x,y,t) = 0, \quad (35)$$

$$u_2(x,y,0) = 0,$$

⋮

Using the initial approximation $u_0(x,y,t) = \sqrt{\sin x \sinh y}$, and solving the above equations, we obtain the approximations as follows

$$u_0(x,y,t) = \sqrt{\sin x \sinh y}, \quad (36)$$

$$u_1(x,y,t) = \sqrt{\sin x \sinh yt}, \quad (37)$$

$$u_2(x,y,t) = \sqrt{\sin x \sinh y} \frac{t^2}{2!}, \quad (38)$$

⋮

$$u_n(x,y,t) = \sqrt{\sin x \sinh y} \frac{t^n}{n!}. \quad (39)$$

Therefore, the exact solution of (31) can be expressed as

$$u(x,y,t) = \lim_{n \rightarrow \infty} \left(\sqrt{\sin x \sinh y} + \sqrt{\sin x \sinh y} t + \sqrt{\sin x \sinh y} \frac{t^2}{2!} + \dots + \sqrt{\sin x \sinh y} \frac{t^n}{n!} \right)$$

$$= \lim_{n \rightarrow \infty} \sqrt{\sin x \sinh y} \left(1 + t + \frac{t^2}{2!} + \dots + \frac{t^n}{n!} \right) = \sqrt{\sin x \sinh y} e^t. \quad (40)$$

4. Conclusions

The main goal of this work was to employ homotopy perturbation method for finding the approximate analytical solution of biological population models. Three examples were presented in this study to illustrate the reliability and applicability of the method. The analytical solution in each of the examples obtained in terms of an infinite series with easily computable components which converges very rapidly to the exact solution without using any restrictive assumption, perturbation or discretization of the variables. Furthermore, the approximate solutions obtained using HPM are in excellent agreement with those obtained by the decomposition method of Adomian. However, due to its ease in calculations, the HPM is a more reliable and powerful mathematical tool that can be applied to other non-linear partial differential equations.

Acknowledgements

The author wishes to thank the anonymous referees for their valuable suggestions to improve the quality of the paper.

REFERENCES

- Adomian, G. (1988). A review of the decomposition method in applied mathematics, *J. Math. Anal. Appl.* 135, pp. 501-544.
- Adomian, G. (1994). *Solving Frontier problems of physics: The decomposition method*, Kluwer, Academic, Dordrecht.
- Abdou, M.A., Soliman, A.A. (2005). Variational iteration method for solving Burger's and coupled Burger's equations. *J. Comput. Appl. Math.* 181, pp. 245-251.
- Evans, D. J., Raslan, K.R. (2005). The Than-Function method for solving some important nonlinear partial differential equation. *Internat. J. Computational Mathematic.* 82, pp. 897-905.
- Fan, E. (2000). Than-Function method and its applications to nonlinear equations. *Phys. Lett. A* 277, pp. 212-218.
- Ganji, D. D., Sadighi, A. (2006). Application of He's homotopy-perturbation method to nonlinear coupled systems of reaction-diffusion equations. *International Journal Nonlinear Science Numerical solution.* 7 (4), 411-418.
- Gorguis, A. (2006). A comparison between Cole-Hopf transformation and the decomposition method for solving Burgers's equation. *Appl. Math. Comput.* 173 (1), pp. 126-136.
- Gurtin, M.E. and MacCamy, R.C. (1977). On the diffusion of biological populations, *Math. Biosci.* 33, pp. 35-49.
- He, J. H. (1999a). Homotopy-perturbation technique, *Comput. Methods Appl. Mech. Engrg.* 178, pp. 257-262.
- He, J. H. (1999b). Variational iteration method: a kind of nonlinear analytical technique: Some examples, *International Journal of Nonlinear Mechanics*, 34, pp. 699-708.

- He, J. H. (2000a). A coupling method of a homotopy technique and a perturbation technique for nonlinear problems, *Int. Nonlinear Mech.* 35, pp. 37-43.
- He, J. H. (2000b). Variational iteration method for autonomous ordinary differential systems, *Applied Mathematics and Computation*, 114, pp. 115-123.
- He, J. H. (2005). Application of homotopy perturbation method to nonlinear wave equations. *Chaos Solitons and Fractals*, 26, pp. 695-700.
- He, J. H. (2006). Some asymptotic methods for strongly nonlinear equations. *International J. Modern Physics. B* 20, pp. 1141-1199.
- He, J.H. & Wu, X. H. (2006). Construction of solitary solution and compaction-like solution by variational iteration method. *Chaos Solitons and Fractals*, 29 (1), pp. 108-113.
- Holmes, M. H. (1995). *Introduction to perturbation methods*. Springer, Berlin.
- Janalizadeh, A., Barari, A., Ganji, D. D. (2008). Application of the homotopy-perturbation method for solving non-linear wave equations. *Journal of Physics: Conference Series* 96, 12080.
- Koçak, H., Yıldırım, A. (2009). Numerical solution of 3D Green's function for the dynamic system of anisotropic elasticity, *Physics Letters A*, 373 (35), pp. 3145-3150.
- Mohyud-Din, S. T., Noor, M. A., Noor, K. I. (2009). Variational iteration method for solving Telegraph equations. *Applications and Applied Mathematics*, 4, pp. 114-121.
- Momani, S. (2005). Analytical approximate solution for fractional heat-like and wave-like equations with variable coefficients using the decomposition method. *Appl. Math. Comput.* 165, pp. 459-472.
- Nayfeh, A.H. (2000). *Perturbation methods*, Wiley, New York.
- Noor, M. A., Mohyud-Din, S. T. (2008). Variational iteration method for solving higher order non-linear boundary value problems using He's polynomial. *Int. J. Nonlin. Sci. Num. Simulation.* 9 (2), pp. 141-157.
- Shakeri, F. and Dehghan, M. (2007). Numerical solution of a biological population model using He's variational iteration method, *Computers & Mathematics with applications*, 54, pp. 1197.
- Xu, L. (2009). The Variational iteration method for fourth-order boundary value problems. *Chaos Solitons and Fractals*, 39 (3), pp. 1386-1394.
- Yıldırım, A., Momani, S. (2010a). Series solutions of a fractional oscillator by means of the homotopy perturbation method, *International Journal of Computer Mathematics*, 87 (5) pp. 1072-1082.
- Yıldırım, A. and Gülkanat, Y. (2010b). Analytical approach to fractional Zakharov-Kuznetsov equations by He's homotopy perturbation method, *Communications in Theoretical Physics*, 53 (6), pp. 1005-1010.