



12-2010

On the Eigenvalue and Inertia Problems for Descriptor Systems

Asadollah Aasaraai
University of Guilan

Kameleh N. Pirbazari
University of Guilan

Follow this and additional works at: <https://digitalcommons.pvamu.edu/aam>



Part of the [Analysis Commons](#), and the [Ordinary Differential Equations and Applied Dynamics Commons](#)

Recommended Citation

Aasaraai, Asadollah and Pirbazari, Kameleh N. (2010). On the Eigenvalue and Inertia Problems for Descriptor Systems, *Applications and Applied Mathematics: An International Journal (AAM)*, Vol. 5, Iss. 2, Article 27.

Available at: <https://digitalcommons.pvamu.edu/aam/vol5/iss2/27>

This Article is brought to you for free and open access by Digital Commons @PVAMU. It has been accepted for inclusion in *Applications and Applied Mathematics: An International Journal (AAM)* by an authorized editor of Digital Commons @PVAMU. For more information, please contact hvkoshy@pvamu.edu.



On the Eigenvalue and Inertia Problems for Descriptor Systems

Asadollah Aasaraai and Kameleh Nasiri Pirbazari

Department of Mathematics
 University of Guilan
 P.O. Box 41335-1914
 Rasht, Iran

AASARAAI@Gmail.com ; k-nasiri@guilan.ac.ir

Received: February 14, 2010; Accepted: October 12, 2010

Abstract

The present study is intended to demonstrate that for a descriptor system with matrix pencil (C, G) , there exists a matrix F such that matrix FG and matrix pencil (C, G) have the same positive and negative eigenvalues. It is also shown that matrix F can be calculated as a contour integral. On the other hand, different representations for matrix F are introduced.

Keywords: Descriptor system; matrix pencil; Weierstrass canonical normal form; generalized eigenvalues; inertia

MSC 2000 No.: 45CXX; 34HXX

1. Introduction

Matrix pencils appear in the process of analyzing descriptor systems

$$\begin{cases} Cx(t) = Gx(t) + Bu(t) \\ y = Lx(t) + Du(t) \end{cases} \quad (1)$$

where $C, G \in R^{n \times n}$, $x \in R^n$, $u \in R^m$, $B \in R^{n \times m}$, $L \in R^{p \times n}$, $D \in R^{p \times m}$. It is generally known that the stability properties of system (1) can be characterized in terms of the eigenvalues of pencil $\lambda C - G$ and that the controllability and observability properties of system (1) depend on

the rank of matrix pencils $\lambda C - G$ and $\lambda C^T - G^T$, respectively. See Cobb (1984), Mehrmann and Stykel (2005) and Yip and Sincovec (1981). Note that the eigenvalues of $\lambda C - G$ may be ill-conditioned in the sense that they may change largely even for small perturbations in C and G . On the other hand, there are different methods for various types of eigenvalues problems, see Mehrmann and Voss (2005) and Van den Eshof (2002). For a historical background of eigenproblems, one can refer to Ciarlet (2001).

First, some basic concepts that are later needed for the purposes of the present paper are to be introduced. Matrix $\lambda C - G$ is called a matrix pencil where $C, G \in R^{n \times n}$ and $\lambda \in C$. Matrix pencil is also shown by matrix pair (C, G) . Pencil (C, G) is called *regular* if there exists a $\lambda \in C$ such that $\det(\lambda C - G) \neq 0$; otherwise, it is called *singular*. Descriptor system (1) is called regular if pencil (C, G) is regular. The analyses in this paper are based on regular descriptor systems. Eigenvalues of matrix pencil (C, G) are defined to be the roots of characteristic polynomial $\det(\lambda C - G) = 0$. $sp(C, G)$ is the set of eigenvalues of pencil (C, G) . Value $\lambda_0 \in C$ is called a finite eigenvalue of $\lambda C - G$, if $\det(\lambda_0 C - G) = 0$. If matrix C is singular, then $\lambda C - G$ is said to have an eigenvalue at infinity. Two matrix pencils of (C_1, G_1) and (C_2, G_2) are called *equivalent* if there exist nonsingular matrices T and S such that $\lambda C_1 - G_1 = \lambda T C_2 S - T G_2 S$. Let $\lambda C - G$ be a regular pencil with $C, G \in R^{n \times n}$. There exist two nonsingular matrices T and $S \in C^{n \times n}$ such that

$$TCS = \text{diag}(I_{n_1}, N), \quad TGS = \text{diag}(J, I_{n_2}), \quad (2)$$

where matrices J and N are in Jordan canonical form and matrix N is a nilpotent matrix; that is, there exists an integer ν such that $N^\nu = 0$ but $N^{\nu-1} \neq 0$.

The representation (TCS, TGS) of pencil (C, G) is called Weierstrass canonical normal form. See Dai (1989). The inertia of a regular pencil $\lambda C - G$ is defined by a quadruple of integers

$$In(C, G) = \{\pi_-(C, G), \pi_+(C, G), \pi_0(C, G), \pi_\infty(C, G)\},$$

where $\pi_-(C, G)$, $\pi_+(C, G)$ and $\pi_0(C, G)$ are the numbers of eigenvalues with negative, positive and zero real parts, respectively and where $\pi_\infty(C, G)$ is the number of infinite eigenvalues of $\lambda C - G$.

2. Reduction Generalized Eigenvalue and Inertia Problem to Standard Form

Theorem1. There exists a matrix F such that

$$\pi_+(C, G) = \pi_+(FG) \quad \pi_-(C, G) = \pi_-(FG).$$

Proof:

Assume that matrix pencil (C, G) is in canonical form (TCS, TGS) ,

$$TCS = \text{diag}(I_{n_f}, N), \quad TGS = \text{diag}(J, I_{n_m}),$$

where matrices J and N are in Jordan canonical form and matrix N is a nilpotent matrix; then we define the matrix as

$$F = \text{Sdiag}(I_{n_f}, 0)T.$$

Therefore,

$$\begin{aligned} FG &= \text{Sdiag}(I_{n_f}, 0)TG = \text{Sdiag}(I_{n_f}, 0)TGSS^{-1} \\ &= \text{Sdiag}(I_{n_f}, 0)\text{diag}(J, I_{n_m})S^{-1} = \text{Sdiag}(J, 0)S^{-1} \end{aligned}$$

Thus,

$$\text{sp}(FG) = \text{sp}(J) \cup \{0\}.$$

Also, it is generally known that the finite eigenvalues of $\lambda C - G$ are the eigenvalues of matrix J , see Malyshev (1990) and Lewis(1985), so

$$\pi_+(\mathcal{C}, \mathcal{G}) = \pi_+(J) = \pi_+(FG) \quad \pi_-(\mathcal{C}, \mathcal{G}) = \pi_-(J) = \pi_-(FG).$$

In fact, it is hence proved that matrix pencil $(\mathcal{C}, \mathcal{G})$ and matrix FG have the same positive and negative eigenvalues.

In the next step, we want to show the relation of F with coefficients of Laurent series of $(\lambda C - G)^{-1}$ at infinity. It is generally known that Laurent series of $(\lambda C - G)^{-1}$ at infinity are in the following form

$$(\lambda C - G)^{-1} = \sum_{n=-\infty}^{+\infty} h_n \lambda^{-n-1},$$

where

$$h_n = \text{Sdiag}(J^n, 0)T \quad n = 0, 1, 2, \dots$$

and

$$h_n = \text{Sdiag}(0, -N^{-n-1})T \quad n = -1, -2, \dots$$

See Lewis (1985).

Theorem 2. Let c be a closed simple curve such that the finite eigenvalues of $\lambda C - G$ lie inside c , then

$$h_n = \frac{1}{2\pi i} \oint_c \lambda^n (\lambda C - G)^{-1} d\lambda \quad n \geq 0$$

Proof:

We have

$$(\lambda C - G)^{-1} = \left[\lambda T^{-1} \text{diag} (L_{n_f}, N) S^{-1} - T^{-1} \text{diag} (J, L_{n_m}) S^{-1} \right]^{-1} =$$

$$S \text{diag} (\lambda L_{n_f} - J, \lambda N - L_{n_m})^{-1} T = S \text{diag} [(\lambda L_{n_f} - J)^{-1}, (\lambda N - L_{n_m})^{-1}] T.$$

So,

$$\frac{1}{2\pi i} \oint_c \lambda^n (\lambda C - G)^{-1} d\lambda =$$

$$\frac{1}{2\pi i} S \text{diag} \left[\oint_c \lambda^n (\lambda L_{n_f} - J)^{-1} d\lambda, \oint_c \lambda^n (\lambda N - L_{n_m})^{-1} d\lambda \right] T$$

But, c includes all finite eigenvalues of $\lambda C - G$. Hence,

$$\frac{1}{2\pi i} \oint_c \lambda^n (\lambda L_{n_f} - J)^{-1} d\lambda = J^n$$

and since N is nilpotent,

$$\frac{1}{2\pi i} \oint_c \lambda^n (\lambda N - L_{n_m})^{-1} d\lambda = 0$$

Thus,

$$\frac{1}{2\pi i} \oint_c \lambda^n (\lambda C - G)^{-1} d\lambda = S \text{diag} (J^n, 0) T = h_n \quad n \geq 0$$

Theorem 3. Let c be a closed simple curve such that the finite eigenvalues of $\lambda C - G$ lie inside c . Furthermore, c includes origin. Then,

$$h_n = \frac{-1}{2\pi i} \oint_c \lambda^n (\lambda C - G)^{-1} d\lambda \quad n < 0$$

Proof:

Since \mathfrak{c} includes origin,

$$\frac{-1}{2\pi i} \oint_{\mathfrak{c}} \lambda^n (\lambda L_{n_f} - D)^{-1} d\lambda = 0$$

and

$$\frac{-1}{2\pi i} \oint_{\mathfrak{c}} \lambda^n (\lambda N - L_{n_g})^{-1} d\lambda = -N^{-n-1}$$

Thus,

$$\frac{-1}{2\pi i} \oint_{\mathfrak{c}} \lambda^n (\lambda \mathcal{C} - \mathcal{G})^{-1} d\lambda = \text{Sdiag}(0, -N^{-n-1})T \quad n \leq 0$$

So far, a generalized eigenvalue and inertia problem have been reduced to a standard eigenvalue and inertia problem by using Weierstrass canonical normal form. Also the coefficients of Laurent series of $(\lambda \mathcal{C} - \mathcal{G})^{-1}$ at infinity have been described as contour integrals. In sequel it will be shown that matrix F can be calculated as a contour integral. Then, different representations for matrix F will be obtained.

Corollary 1. F is the coefficient of λ^{-1} in the Laurent series of $(\lambda \mathcal{C} - \mathcal{G})^{-1}$ at infinity.

Proof:

The coefficient of λ^{-1} in the Laurent series of $(\lambda \mathcal{C} - \mathcal{G})^{-1}$ at infinity is equal to

$$h_0 = \frac{1}{2\pi i} \oint_{\mathfrak{c}} (\lambda \mathcal{C} - \mathcal{G})^{-1} d\lambda = \text{Sdiag}(L_{n_f}, 0)T = F.$$

3. Some Representations for Matrix F

In the previous section, matrix F was defined in terms of two matrices T, S . This section attempts to find different representations for F . One of them is a representation that just depends on matrix S ; the other describes F in terms of matrix T ; and the third representation describes the relation between F and the constant term of Laurent series of $(\lambda \mathcal{C} - \mathcal{G})^{-1}$ at infinity.

Lemma1. There exists a representation for F that just depends on matrix S .

Proof:

We define matrix T_r as follows:

$$T_r = S \text{dtag}(L_{n_f}, 0) S^{-1},$$

where S is the matrix defined in canonical normal form. We would then have

$$\begin{aligned} CT_r &= T^{-1} T C S S^{-1} T_r = T^{-1} \text{dtag}(L_{n_f}, N) S^{-1} S \text{dtag}(L_{n_f}, 0) S^{-1} = \\ &T^{-1} \text{dtag}(L_{n_f}, 0) S^{-1}. \end{aligned} \quad (3)$$

Also,

$$I - T_r - S S^{-1} - S \text{dtag}(L_{n_f}, 0) S^{-1} - S \text{dtag}(0, I_{n_m}) S^{-1}$$

and

$$\begin{aligned} G(I - T_r) &= T^{-1} T G S S^{-1} (I - T_r) \\ &= T^{-1} \text{dtag}(J, I_{n_m}) S^{-1} S \text{dtag}(0, I_{n_m}) S^{-1} \\ &= T^{-1} \text{dtag}(0, I_{n_m}) S^{-1}. \end{aligned} \quad (4)$$

By adding relations (3) and (4)

$$CT_r + G(I - T_r) = T^{-1} \text{dtag}(L_{n_f}, 0) S^{-1} + T^{-1} \text{dtag}(0, I_{n_m}) S^{-1} = T^{-1} \text{dtag}(L_{n_f}, I_{n_m}) S^{-1},$$

So,

$$[CT_r + G(I - T_r)]^{-1} = S \text{dtag}(L_{n_f}, I_{n_m}) T.$$

Thus,

$$T_r [CT_r + G(I - T_r)]^{-1} = S \text{dtag}(L_{n_f}, 0) S^{-1} S \text{dtag}(L_{n_f}, I_{n_m}) T = S \text{dtag}(L_{n_f}, 0) T.$$

But, the last term is equal to F and the proof is then complete.

Lemma2. There exists a representation for F that just depends on matrix T .

Proof:

The proof is similar to Lemma 1. It is sufficient to define

$$T_i = T^{-1} \text{diag}(L_{n_f}, \mathbf{0}) T$$

It, then, follows that

$$F = [T_i C + (I - T_i) G]^{-1} T_i$$

Theorem 4. There exists a matrix E such that

$$I - T_p = -EG, \quad I - T_i = -GE.$$

Proof:

Let E be the constant term of Laurent series of $(\lambda C - G)^{-1}$ at infinity.

$$E = h_{-1} = \frac{-1}{2\pi i} \oint_{\mathcal{C}} \lambda^{-1} (\lambda C - G)^{-1} d\lambda$$

According to computations in Theorem 2,

$$\lambda^{-1} (\lambda C - G)^{-1} = S \text{diag}[\lambda^{-1} (\lambda L_{n_f} - J)^{-1}, \lambda^{-1} (\lambda N - L_{n_w})^{-1}] T$$

So,

$$E = \frac{-1}{2\pi i} S \text{diag} \left[\oint_{\mathcal{C}} \lambda^{-1} (\lambda L_{n_f} - J)^{-1} d\lambda, \oint_{\mathcal{C}} \lambda^{-1} (\lambda N - L_{n_w})^{-1} d\lambda \right] T$$

Since \mathcal{C} includes the origin

$$\frac{-1}{2\pi i} \oint_{\mathcal{C}} \lambda^{-1} (\lambda L_{n_f} - J)^{-1} d\lambda = \mathbf{0}$$

and

$$\frac{-1}{2\pi i} \oint_{\mathcal{C}} \lambda^{-1} (\lambda N - L_{n_w})^{-1} d\lambda = -L_{n_w}$$

Therefore,

$$E = -S \text{diag}(\mathbf{0}, L_{n_w}) T$$

and

$$-EG = S \text{diag}(0, I_{n_m}) T T^{-1} \text{diag}(J, I_{n_m}) S^{-1} = S \text{diag}(0, I_{n_m}) S^{-1}.$$

But,

$$I - T_r = I - S \text{diag}(I_{n_r}, 0) S^{-1} = S \text{diag}(0, I_{n_m}) S^{-1}.$$

Hence,

$$I - T_r = -EG.$$

Similar computations show that

$$I - T_l = -GE.$$

Corollary 2. The following equalities hold

$$F = (I + EG)[C(I + EG) - GEG]^{-1} = [(I + GE)C - GEG]^{-1}(I + GE).$$

Proof:

It follows immediately from Lemma 1 and Lemma 2 and Theorem 4.

4. Conclusion

In this paper, a method for the reduction of a generalized eigenvalue and inertia problem to a standard eigenvalue and inertia problem is presented. The method is based on the use of Weierstrass canonical normal form and matrix F . In fact, it is demonstrated that matrix FG and matrix pencil (C, G) have the same positive and negative eigenvalues. Also according to the analyses presented, it is stated that matrix F can be described in terms of coefficients of Laurent series of $(\lambda C - G)^{-1}$ at infinity. As part of this study, other representations for F have also been offered. It is to be noted that the regularity of matrix pencil (C, G) was the only assumption used in this paper and the results are valid even if C and G are singular matrices.

REFERENCES

Ciarlet, P.G and Lions, J.L. (2001). Computational Methods for Large Eigenvalues Probl Hand book of Numerical Analysis. vol. VIII. North-Holland (Elsevier) 2001, pp. 3-179.

- Cobb, D.J. (1984). Controllability, observability and duality in singular systems. IEEE Transactions on Automatic Control.
- Dai, L. (1989). Singular Control Systems, volume 118 of Lecture Notes in Control and Information Sciences. Springer-Verlag, Berlin.
- Lewis, F.L. (1985). Fundamental, reachability and observability matrices for discrete des systems. IEEE Trans. Automat Control
- Malyshev, A. (1990). Computing invariant subspaces of a regular linear pencil of matrices Siberian Math. J. 30(4): 559-567
- Mehrmann, V and Stykel, T. (2005). Balanced truncation model reduction for large scale systems in descriptor form. Volume 45 of Lecture Notes in Computational Science and Engineering. Springer-Verlag, Berlin.
- Mehrmann, V. and Voss, H. (2005). Nonlinear eigenvalue problems: a challenge for modern eigenvalue methods, Mitt. Gesell. Ang. Math. Mech. 27, 121-151.
- Van den Eshof, J. (2002). The convergence of Jacobi-Davidson iteration for Hermitian eigenvalue problems, Num. Lin. Alg. Appl. 9, 163-179
- Yip, E.L. and Sincovec, R.F. (1981). Solvability, controllability and observability of continuous descriptor systems. IEEE Trans. Automat. Control