



6-2011

Positive Solutions for n th Order Differential Equations Under Some Conditions

S. N. Odda
Qassim University

Follow this and additional works at: <https://digitalcommons.pvamu.edu/aam>



Part of the [Ordinary Differential Equations and Applied Dynamics Commons](#)

Recommended Citation

Odda, S. N. (2011). Positive Solutions for n th Order Differential Equations Under Some Conditions, *Applications and Applied Mathematics: An International Journal (AAM)*, Vol. 6, Iss. 1, Article 19. Available at: <https://digitalcommons.pvamu.edu/aam/vol6/iss1/19>

This Article is brought to you for free and open access by Digital Commons @PVAMU. It has been accepted for inclusion in *Applications and Applied Mathematics: An International Journal (AAM)* by an authorized editor of Digital Commons @PVAMU. For more information, please contact hvkoshy@pvamu.edu.



Available at
<http://pvamu.edu/aam>
Appl. Appl. Math.
ISSN: 1932-9466

Applications and Applied
Mathematics:
An International Journal
(AAM)

Vol. 6, Issue 1 (June 2011) pp. 232 – 239
(Previously, Vol. 6, Issue 11, pp. 1973 – 1980)

Positive Solutions for n th Order Differential Equations Under Some Conditions

S.N. Odda

Department of Mathematics
Faculty of Computer Science
Qassim University
Burieda, Saudi Arabia
nabhan100@yahoo.com

Received: January 13, 2011; Accepted: May 26, 2011

Abstract

This paper presents an existence and nonexistence of positive solutions for the nonlinear boundary value problems. We prove that the n th order nonlinear differential equation has at least one positive solution by using appropriate fixed point theorems.

Keywords: Fixed-point theorem, Banach space, nonlinear n th order differential equations, Positive solutions

AMS 2010 No.: 34 B15, 34 B18

1. Introduction

We are concerned, in this paper, with the existence of positive solutions for the following boundary value problem for n th order differential equations

$$u^{(n)}(t) = f(t, u(t)), \quad 0 < t < 1, \quad (1.1)$$

$$u(0) = u'(0) = u''(0) = \dots = u^{(n-2)}(0) = u^{(n-1)}(1) = 0, \quad (1.2)$$

$$\alpha u'(1) + \beta u''(1) = 0, \text{ where } \beta, \alpha \geq 0, \alpha + \beta > 0 \quad (1.3)$$

Problems of the above type occur frequently in science, engineering, mathematical physics, economics and biology [Zhang (2006)].

The nonlinear n th order differential equations studied in this paper are an existence and nonexistence of positive solutions by using object of mathematical investigations [El-shahed (2009), El-Shahed and Hassan (2010), Guo and Lakshmikantham (1988), Sun and Wen (2006), Agarwal and O'Regan (1999) and Agarwal et al. (1999)]. However, there are few papers investigating the existence of positive solutions of n th impulsive differential equations by using the fixed point theorem of cone expansion and compression. The objective of the present paper is to fill this gap and the results presented are new and original. Also, several results obtained in Agarwal et al. (1999) are generalized.

2. Notation, Definition and Auxiliary Results

Theorem 2.1 [Agarwal et al. (2001), Agarwal and O'Regan (1999)]:

Assume that U is a relatively open subset of convex set K in Banach space E . Let

$N : \bar{U} \rightarrow K$ be a compact map with $0 \in U$. Then, either

- (i) N has a fixed point in \bar{U} ; or
- (ii) There is $u \in U$ and $\lambda \in (0,1)$ such that $u = \lambda N u$.

At first, we find the solution $u(t)$, for the problem

$$u^{(n)}(t) = y(t), \quad 0 < t < 1, \quad (2.1)$$

$$u(0) = u'(0) = u''(0) = \dots = u^{(n-2)}(0) = u^{(n-1)}(1) = 0, \quad (2.2)$$

$$\alpha u'(1) + \beta u''(1) = 0, \text{ where } \beta, \alpha \geq 0, \alpha + \beta > 0. \quad (2.3)$$

Applying Laplace transforms to equation (2.1), we have:

$$s^n \bar{u}(s) - s^{n-1} u(0) - s^{n-2} u'(0) - s^{n-3} u''(0) - \dots - s u^{(n-2)}(0) - u^{(n-1)}(0) = \bar{y}(s), \quad (2.4)$$

where $\bar{u}(s)$ and $\bar{y}(s)$ is the Laplace transform of $u(t)$ and $y(t)$ respectively. So,

$$\bar{u}(s) = \frac{s^{n-1}u(0)}{s^n} + \frac{s^{n-2}u'(0)}{s^n} + \frac{s^{n-3}u''(0)}{s^n} + \dots + \frac{su^{(n-2)}(0)}{s^n} + \frac{u^{(n-1)}(0)}{s^n} + \frac{\bar{y}(s)}{s^n}.$$

Therefore, by the inverse Laplace transform and using the boundary condition (2.2) and (2.3), we obtain the final form of $u(t)$ as:

$$u(t) = \frac{\alpha + (n - 2)\beta}{[\alpha + \beta](n - 2)!} \int_0^1 \frac{t^2}{2!} y(s) ds - \frac{\alpha}{[\alpha + \beta](n - 2)!} \int_0^1 \frac{t^2(1 - s)^{n-2}}{2!} y(s) ds - \frac{\beta}{[\alpha + \beta](n - 3)!} \int_0^1 \frac{t^2(1 - s)^{n-3}}{2!} y(s) ds - \int_0^1 \frac{t^{n-1}}{(n - 1)!} y(s) ds + \int_0^t \frac{(t - s)^{n-1}}{(n - 1)!} y(s) ds. \tag{2.5}$$

Definition 2.1.

An operator is called completely continuous if it is continuous and maps bounded sets into precompacts.

Definition 2.2.

Let E be a real Banach space. A nonempty closed convex set $K \subset E$ is called cone of E if it satisfies the following conditions:

- (i) $x \in K, \sigma \geq 0$ implies $\sigma x \in K$; and
- (ii) $x \in K, -x \in K$ implies $x = 0$.

3. Main Result

Consider the family of problems:

$$u^{(n)}(t) = f(t, u(t)), \quad 0 < t < 1, \tag{3.1}$$

$$u(0) = u'(0) = u''(0) = \dots = u^{(n-2)}(0) = u^{(n-1)}(1) = 0, \tag{3.2}$$

$$\alpha u'(1) + \beta u''(1) = 0, \text{ where } \beta, \alpha \geq 0, \alpha + \beta > 0. \tag{3.3}$$

Hence, (3.1), (3.2) and (3.3) are equivalent to the integral equation

$$\begin{aligned}
 u(t) = & \frac{\alpha + (n-2)\beta}{[\alpha + \beta](n-2)!} \int_0^1 \frac{t^2}{2!} f(s, u(s)) ds - \frac{\alpha}{[\alpha + \beta](n-2)!} \int_0^1 \frac{t^2(1-s)^{n-2}}{2!} f(s, u(s)) ds \\
 & - \frac{\beta}{[\alpha + \beta](n-3)!} \int_0^1 \frac{t^2(1-s)^{n-3}}{2!} f(s, u(s)) ds - \int_0^1 \frac{t^{n-1}}{(n-1)!} f(s, u(s)) ds \\
 & + \int_0^t \frac{(t-s)^{n-1}}{(n-1)!} f(s, u(s)) ds.
 \end{aligned} \tag{3.4}$$

Defining $T : X \rightarrow X$ as:

$$\begin{aligned}
 Tu(t) = & \frac{\alpha + (n-2)\beta}{[\alpha + \beta](n-2)!} \int_0^1 \frac{t^2}{2!} f(s, u(s)) ds - \frac{\alpha}{[\alpha + \beta](n-2)!} \int_0^1 \frac{t^2(1-s)^{n-2}}{2!} f(s, u(s)) ds \\
 & - \frac{\beta}{[\alpha + \beta](n-3)!} \int_0^1 \frac{t^2(1-s)^{n-3}}{2!} f(s, u(s)) ds - \int_0^1 \frac{t^{n-1}}{(n-1)!} f(s, u(s)) ds + \int_0^t \frac{(t-s)^{n-1}}{(n-1)!} f(s, u(s)) ds,
 \end{aligned} \tag{3.5}$$

where $X=C[0, I]$ is the Banach space endowed with the sup norm.

We have the following result for operator T .

Lemma 3.1:

Assume that $f : [0,1] \times R \rightarrow R$ is continuous function, then T is completely continuous operator.

Proof:

It is easy to see that T is continuous. For $u \in M = \{u \in X : \|u\| \leq l, l > 0\}$, we obtain,

$$\begin{aligned}
 Tu(t) = & \left| \frac{\alpha + (n-2)\beta}{[\alpha + \beta](n-2)!} \int_0^1 \frac{t^2}{2!} f(s, u(s)) ds - \frac{\alpha}{[\alpha + \beta](n-2)!} \int_0^1 \frac{t^2(1-s)^{n-2}}{2!} f(s, u(s)) ds \right. \\
 & \left. - \frac{\beta}{[\alpha + \beta](n-3)!} \int_0^1 \frac{t^2(1-s)^{n-3}}{2!} f(s, u(s)) ds - \int_0^1 \frac{t^{n-1}}{(n-1)!} f(s, u(s)) ds + \int_0^t \frac{(t-s)^{n-1}}{(n-1)!} f(s, u(s)) ds \right| \\
 \leq & \frac{\alpha + (n-2)\beta}{[\alpha + \beta](n-2)!} \int_0^1 \frac{t^2}{2!} |f(s, u(s))| ds + \frac{\alpha}{[\alpha + \beta](n-2)!} \int_0^1 \frac{t^2(1-s)^{n-2}}{2!} |f(s, u(s))| ds \\
 & + \frac{\beta}{[\alpha + \beta](n-3)!} \int_0^1 \frac{t^2(1-s)^{n-3}}{2!} |f(s, u(s))| ds + \int_0^1 \frac{t^{n-1}}{(n-1)!} |f(s, u(s))| ds + \int_0^t \frac{(t-s)^{n-1}}{(n-1)!} |f(s, u(s))| ds \\
 \leq & \frac{\alpha + (n-2)\beta}{[\alpha + \beta](n-2)!} \frac{L}{2!} + \frac{\alpha}{[\alpha + \beta](n-2)!} \frac{L}{2!} + \frac{\beta}{[\alpha + \beta](n-3)!} \frac{L}{2!} + \frac{L}{(n-1)!} + \frac{L}{n!},
 \end{aligned}$$

where $L = \max_{0 \leq t \leq 1, \|u\| \leq 1} |f(t, u(t))| + 1$, so $T(M)$ is bounded. Next we shall show the equi-continuity of $\overline{T(M)}$, $\forall u \in M, \forall \varepsilon > 0, t_1 < t_2 \in [0, 1]$.

Let

$$\eta < \left\{ 2! \frac{\varepsilon[\alpha + \beta](n-2)!}{5L[\alpha + (n-2)\beta]}, 2! \frac{\varepsilon[\alpha + \beta](n-1)!}{5L\alpha}, 2! \frac{\varepsilon[\alpha + \beta](n-2)!}{5L\beta} \right\},$$

$$\sigma < \left\{ \frac{\varepsilon(n-1)!}{5L} \right\},$$

$$\gamma < \left\{ \varepsilon \frac{n!}{5L} \right\}, \quad t_2^2 - t_1^2 < \eta, \quad t_2^{n-1} - t_1^{n-1} < \sigma, \quad t_2^n + t_1^n < \gamma.$$

Then, we have

$$\begin{aligned} |Tu(t_2) - Tu(t_1)| &= \left| \frac{\alpha + (n-2)\beta}{[\alpha + \beta](n-2)!} \int_0^{t_2^2 - t_1^2} \frac{(t_2^2 - t_1^2 - s)^{n-2}}{2!} f(s, u(s)) ds - \frac{\alpha}{[\alpha + \beta](n-2)!} \int_0^{t_2^2 - t_1^2} \frac{(t_2^2 - t_1^2)(1-s)^{n-2}}{2!} f(s, u(s)) ds \right. \\ &\quad - \frac{\beta}{[\alpha + \beta](n-3)!} \int_0^{t_2^{n-1} - t_1^{n-1}} \frac{(t_2^{n-1} - t_1^{n-1} - s)^{n-3}}{2!} f(s, u(s)) ds - \int_0^{t_2^{n-1} - t_1^{n-1}} \frac{t_2^{n-1} - t_1^{n-1}}{(n-1)!} f(s, u(s)) ds \\ &\quad \left. + \int_0^{t_2} \frac{(t_2 - s)^{n-1}}{(n-1)!} f(s, u(s)) ds - \int_0^{t_1} \frac{(t_1 - s)^{n-1}}{(n-1)!} f(s, u(s)) ds \right| \\ &\leq \frac{\alpha + (n-2)\beta}{[\alpha + \beta](n-2)!} \frac{L(t_2^2 - t_1^2)}{2!} + \frac{\alpha}{[\alpha + \beta](n-1)!} \frac{L(t_2^2 - t_1^2)}{2!} \\ &\quad + \frac{\beta}{[\alpha + \beta](n-2)!} \frac{L(t_2^{n-1} - t_1^{n-1})}{2!} + \frac{L(t_2^{n-1} - t_1^{n-1})}{(n-1)!} + \frac{Lt_1^n}{n!} + \frac{Lt_1^n}{n!} \\ &\leq \frac{\varepsilon}{5} + \frac{\varepsilon}{5} + \frac{\varepsilon}{5} + \frac{\varepsilon}{5} + \frac{\varepsilon}{5}. \end{aligned}$$

Thus, $\overline{T(M)}$ is equicontinuous. The Arzela-Ascoli theorem implies that the operator T is completely continuous.

Theorem 3.1:

Assume that $f : [0, 1] \times R \rightarrow R$ is continuous function, and there exist constants

$$0 < c_1 < \left(\frac{2!(\alpha + \beta)(n-1)!}{n\alpha + (n-1)^2\beta}, \frac{n!}{n+1} \right), \quad c_2 > 0,$$

such that $|f(t, u(t))| \leq c_1|u| + c_2$, for all $t \in [0,1]$. Then, the boundary value problem (3.1)-(3.3) has a solution.

Proof:

Following [Yang (2005), Zhang (2006) and Odda (2010)], we will apply the nonlinear alternative theorem to prove that T has one fixed point. Let $\Omega = \{u \in X : \|u\| < R\}$, be open subset of X , where

$$R > \left(4 \left\{ \frac{n\alpha + (n-1)^2\beta}{2!(n-1)!(\alpha + \beta)} |u| c_1, \frac{(n+1)c_1}{n!} |u|, \frac{n\alpha + (n-1)^2\beta}{2!(n-1)!(\alpha + \beta)} |u| c_2, \frac{(n+1)c_2}{n!} |u| \right\} \right).$$

We suppose that there is a point $u \in \partial\Omega$ and $\lambda \in (0,1)$ such that $u = Tu$. For $u \in \partial\Omega$, we have:

$$\begin{aligned} Tu(t) &= \left| \frac{\alpha + (n-2)\beta}{[\alpha + \beta](n-2)!} \int_0^1 \frac{t^2}{2!} f(s, u(s)) ds - \frac{\alpha}{[\alpha + \beta](n-2)!} \int_0^1 \frac{t^2(1-s)^{n-2}}{2!} f(s, u(s)) ds \right. \\ &\quad \left. - \frac{\beta}{[\alpha + \beta](n-3)!} \int_0^1 \frac{t^2(1-s)^{n-3}}{2!} f(s, u(s)) ds - \int_0^1 \frac{t^{n-1}}{(n-1)!} f(s, u(s)) ds + \int_0^t \frac{(t-s)^{n-1}}{(n-1)!} f(s, u(s)) ds \right| \\ &\leq \frac{\alpha + (n-2)\beta}{[\alpha + \beta](n-2)!} \int_0^1 \frac{t^2}{2!} |f(s, u(s))| ds + \frac{\alpha}{[\alpha + \beta](n-2)!} \int_0^1 \frac{t^2(1-s)^{n-2}}{2!} |f(s, u(s))| ds \\ &\quad + \frac{\beta}{[\alpha + \beta](n-3)!} \int_0^1 \frac{t^2(1-s)^{n-3}}{2!} |f(s, u(s))| ds + \int_0^1 \frac{t^{n-1}}{(n-1)!} |f(s, u(s))| ds + \int_0^t \frac{(t-s)^{n-1}}{(n-1)!} |f(s, u(s))| ds \\ &\leq \frac{\alpha + (n-2)\beta}{[\alpha + \beta](n-2)!} \int_0^1 \frac{t^2}{2!} (c_1|u(s)| + c_2) ds + \frac{\alpha}{[\alpha + \beta](n-2)!} \int_0^1 \frac{t^2(1-s)^{n-2}}{2!} (c_1|u(s)| + c_2) ds \\ &\quad + \frac{\beta}{[\alpha + \beta](n-3)!} \int_0^1 \frac{t^2(1-s)^{n-3}}{2!} (c_1|u(s)| + c_2) ds + \int_0^1 \frac{t^{n-1}}{(n-1)!} (c_1|u(s)| + c_2) ds + \int_0^t \frac{(t-s)^{n-1}}{(n-1)!} (c_1|u(s)| + c_2) ds \\ &\leq \frac{\alpha + (n-2)\beta}{2![\alpha + \beta](n-2)!} (c_1|u(s)| + c_2) + \frac{\alpha}{2![\alpha + \beta](n-1)!} (c_1|u(s)| + c_2) \\ &\quad + \frac{\beta}{2![\alpha + \beta](n-2)!} (c_1|u(s)| + c_2) + \frac{1}{(n-1)!} (c_1|u(s)| + c_2) + \frac{1}{(n)!} (c_1|u(s)| + c_2) \\ &\leq \frac{\alpha + (n-2)\beta}{2![\alpha + \beta](n-2)!} (c_1|u(s)|) + \frac{\alpha}{2![\alpha + \beta](n-1)!} (c_1|u(s)|) \\ &\quad + \frac{\beta}{2![\alpha + \beta](n-2)!} (c_1|u(s)|) + \frac{1}{(n-1)!} (c_1|u(s)|) + \frac{1}{(n)!} (c_1|u(s)|) \\ &+ \frac{\alpha + (n-2)\beta}{2![\alpha + \beta](n-2)!} (c_2) + \frac{\alpha}{2![\alpha + \beta](n-1)!} (c_2) + \frac{\beta}{2![\alpha + \beta](n-2)!} (c_2) + \frac{1}{(n-1)!} (c_2) + \frac{1}{(n)!} (c_2) \\ &< \frac{R}{4} + \frac{R}{4} + \frac{R}{4} + \frac{R}{4} = R, \end{aligned}$$

which implies that $\|T\| \neq R = \|u\|$, that is a contradiction. Then, the nonlinear alternative theorem implies that T has a fixed point $u \in \overline{\Omega}$, that is, problem (3.1)-(3.3), has a solution $u \in \overline{\Omega}$.

Finally, we give an example to illustrate the results obtained in this paper.

Example.

From the equation (3.1) –(3.3) we solve the boundary value problem

$$u^{(5)}(t) = \frac{u+1}{u^2+7}. \quad (3.6)$$

Theorem 3.1 with $\alpha = 1$ and $\beta = 1$, we find $0 < c_1 < \min(\frac{2!(2)4!}{5+(4)^2}, \frac{5!}{6})$. So, we conclude that the problem (3.6) has a solution.

Acknowledgements

The author is grateful to the reviewers for their useful comments.

REFERENCES

- Agarwal, R. P., Meehan, M. and O'Regan, D. (2001). Fixed Point and Applications, Cambridge University Press, Cambridge.
- Agarwal, R.P. and O'Regan, D. (1999). Global existence for nonlinear operator inclusion, Computers Math-Applic.38 (11/12), 131-139.
- Agarwal, R.P. and O'Regan, D. and Wong, P.J.Y. (1999). Positive solutions of differential, difference and integral equations, Kluwer Academic, Dordrecht.
- El-Shahed, M. (2009). Positive solutions for nonlinear singular third order boundary value problem, Communications in Nonlinear Science and Numerical Simulation, Vol.14, 424-429.
- El-Shahed, M. and Hassan, H.A. (2010). Positive solutions of q-Difference equation, American Mathematical Society, Vol. 138, No.5, 1733-1738.
- Guo, D. and Lakshmikantham, V. (1988). Nonlinear Problems in Abstract Cones, Academic Press, San Diego.
- Odda, S. N. (2010). Positive Solutions for Boundary Value Problems of Higher Order differential equations, International Mathematical Forum, Vol. 5, No. 43, 2131 – 2136.
- Odda, S. N. (2010). Existence solution for 5th order differential equation under some conditions, Applied Mathematics, Vol. 1, 279-282.

- Sun, H. and Wen, W. (2006). On the Number of positive solutions for a nonlinear third order boundary value problem, *International Journal of Difference Equations*, 1, 165- 176.
- Yang, B. (2005). Positive solutions for a fourth order boundary value problem, *Electronic Journal of Qualitative Theory of Differential Equations*, 3, 1-17.
- Zhang, Shuqin (2006). Existence of solution for a boundary value problem of fractional order, *Acta Mathematica Scientica*, 26B, 220-228.