



6-2016

ON EXTENSION OF MITTAG-LEFFLER FUNCTION

Ekta Mittal

The IIS University

Rupakshi M. Pandey

Amity University Uttar Pradesh

Sunil Joshi

Manipal University

Follow this and additional works at: <https://digitalcommons.pvamu.edu/aam>



Part of the [Analysis Commons](#), and the [Special Functions Commons](#)

Recommended Citation

Mittal, Ekta; Pandey, Rupakshi M.; and Joshi, Sunil (2016). ON EXTENSION OF MITTAG-LEFFLER FUNCTION, *Applications and Applied Mathematics: An International Journal (AAM)*, Vol. 11, Iss. 1, Article 20.

Available at: <https://digitalcommons.pvamu.edu/aam/vol11/iss1/20>

This Article is brought to you for free and open access by Digital Commons @PVAMU. It has been accepted for inclusion in *Applications and Applied Mathematics: An International Journal (AAM)* by an authorized editor of Digital Commons @PVAMU. For more information, please contact hvkoshy@pvamu.edu.



Available at
<http://pvamu.edu/aam>
Appl. Appl. Math.
ISSN: 1932-9466

**Applications and
Applied Mathematics:**
An International Journal
(AAM)

Vol. 11, Issue 1 (June 2016), pp. 307 - 316

ON EXTENSION OF MITTAG-LEFFLER FUNCTION

Ekta Mittal¹, Rupakshi Mishra Pandey² and Sunil Joshi³

¹Department of Mathematics

The IIS University

Jaipur, Rajasthan, India

ekta.jaipur@gmail.com

²Department of Mathematics

Amity University Uttar Pradesh

Noida, India

rup_ashi@yahoo.com

³Department of Mathematics and Statistics

Manipal University

Jaipur, Rajasthan, India.

sunil.joshi@jaipur.manipal.edu

Received: May 5, 2015; Accepted: March 21, 2016

Abstract

In this paper, we study the extended Mittag-Leffler function by using generalized beta function and obtain various differential properties, integral representations. Further, we discuss Mellin transform of these functions in terms of generalized Wright hyper geometric function and evaluate Laplace transform, and Whittaker transform in terms of extended beta function. Finally, several interesting special cases of extended Mittag-Leffler functions have also be given.

Keywords: Beta Functions; Mellin Transform; Laplace Transform; Whittaker Transform; Extended Riemann Liouville Fractional derivative operator

MSC 2010 No. : 33E12, 44A10

1. Introduction

The Mittag-Leffler function occurs naturally in the solution of fractional order and integral equation. The importance of such functions in physics and engineering is steadily increasing. Some application of the Mittag-Leffler is carried out in the Study of Kinetic Equation, Study of Lorenz System, Random Walk, Levy Flights and Complex System and also in applied problems such as fluid flow, electric network, probability and statistical distribution theory. Haubold et al. (2011) studied various properties of Mittag-Leffler function.

In 1903, the Swedish mathematician Gosta Mittag-Leffler introduced the function.

$$E_{\alpha}(z) = \sum_{n=0}^{\infty} \frac{z^n}{\Gamma(\alpha n + 1)}, \quad (1)$$

where z is a complex variable, Γ is a Gamma function, and $\alpha \geq 0$. It is a direct generalization of the exponential function for $\alpha = 1$, and for $0 < \alpha < 1$, it interpolates between exponential and hypergeometric function $1/(1-z)$. The generalization of $E_{\alpha}(z)$ was further studied by Wiman (1905) as:

$$E_{\alpha,\beta}(z) = \sum_{n=0}^{\infty} \frac{z^n}{\Gamma(\alpha n + \beta)}, \quad (\alpha, \beta \in \mathbb{C}, \operatorname{Re}(\alpha) > 0, \operatorname{Re}(\beta) > 0), \quad (2)$$

Prabhakar (1971) introduced the function $E_{\alpha,\beta}^{\gamma}(z)$ in as:

$$E_{\alpha,\beta}^{\gamma}(z) = \sum_{n=0}^{\infty} \frac{(Y)_n}{\Gamma(\alpha n + \beta)} \frac{z^n}{n!}, \quad (\alpha, \beta, \gamma \in \mathbb{C}, \operatorname{Re}(\alpha) > 0, \operatorname{Re}(\beta) > 0, \operatorname{Re}(\gamma) > 0), \quad (3)$$

where $(Y)_n$ is the Pochhammer symbol by Rainville (1960) such that

$$(Y)_n = \frac{\Gamma(Y+n)}{\Gamma(Y)}, \quad (Y)_0 = 1,$$

$$(Y)_n = Y(Y+1)(Y+2)\dots(Y+n-1), \quad \text{for } n \geq 1.$$

Shukla and Prajapati (2007) introduced the extended Mittag-Leffler function $E_{\alpha,\beta}^{\gamma,q}(z)$ which is defined as follows:

$$E_{\alpha,\beta}^{\gamma,q}(z) = \sum_{n=0}^{\infty} \frac{(Y)_{qn}}{\Gamma(\alpha n + \beta)} \frac{z^n}{n!},$$

$$\{\alpha, \beta, \gamma \in \mathbb{C}, \operatorname{Re}(\alpha) > 0, \operatorname{Re}(\beta) > 0, \operatorname{Re}(\gamma) > 0, q \in (0,1) \cup \mathbb{N}\}, \quad (4)$$

$E_{\alpha,\beta}^{\nu,q}(z)$ Converge absolutely $\forall |z| < 1$, if $q = \text{Re}(\alpha) + 1$, $\forall z$ and for Beta function defined as:

$$B(\alpha, \beta) = \int_0^1 u^{\alpha-1} (1-u)^{\beta-1} du = \frac{\Gamma(\alpha) \Gamma(\beta)}{\Gamma(\alpha + \beta)}, \quad \text{Re}(\alpha) > 0, \text{Re}(\beta) > 0. \quad (5)$$

Laplace transform of the function $f(z)$ is defined as:

$$L(f(z); s) = \int_0^{\infty} e^{-sz} f(z) dz. \quad (6)$$

Mellin-transform and its inverse Transform of the function $f(z)$ is defined as:

$$M[f(z); s] = f^*(s) = \int_0^{\infty} z^{s-1} f(z) dz, \quad \text{Re}(s) > 0, \quad (7)$$

$$f(z) = M^{-1}[f^*(s); z] = \frac{1}{2\pi i} \int_L f^*(s) z^{-s} ds. \quad (8)$$

Whittaker transforms (Whittaker and Watson (1996)) is defined as:

$$\int_0^{\infty} e^{-t/2} t^{\nu-1} W_{\lambda,\mu}(t) dt = \frac{\Gamma\left(\frac{1}{2} + \mu + \nu\right) \Gamma\left(\frac{1}{2} - \mu + \nu\right)}{\Gamma(1 - \lambda + \nu)}, \quad (9)$$

where $\text{Re}(\mu \pm \nu) > -\frac{1}{2}$ and $w_{\lambda,\mu}(t)$ is the Whittaker confluent hyper geometric function.

Wright generalized hypergeometric function is defined as:

$$\begin{aligned} {}_p\Psi_q \left[(a_1, A_1), \dots, (a_p, A_p); (b_1, B_1), \dots, (b_q, B_q); z \right] \\ = \sum_{n=0}^{\infty} \frac{\prod_{i=1}^p \Gamma(a_i + A_i n)}{\prod_{j=1}^q \Gamma(b_j + B_j n)} \cdot \frac{z^n}{n!}. \end{aligned} \quad (10)$$

The classical Riemann-Liouville fractional derivative of order μ is usually defined by

$$D_z^{\mu} [f(z)] = \frac{1}{\Gamma(-\mu)} \int_0^z f(t) (z-t)^{-\mu-1} dt, \quad (\text{Re}(\mu) < 0),$$

where the integration path is a line from 0 to z in the complex t -plane. For the case $(m-1) < \text{Re}(\mu) < m$ ($m = 1, 2, 3, \dots$), it is defined by

$$\begin{aligned} D_z^\mu [f(z)] &= \frac{d^m}{dz^m} D_z^{\mu-m} \{f(z)\} \\ &= \frac{d^m}{dz^m} \left[\frac{1}{\Gamma(-\mu+m)} \int_0^z f(t)(z-t)^{-\mu+m-1} dt \right]. \end{aligned}$$

The extended Riemann-Liouville fractional derivative operator was defined by Özarslan and Özergin (2010) as follows:

$$D_z^{\mu,p} [f(z)] = \frac{1}{\Gamma(-\mu)} \int_0^z f(t)(z-t)^{-\mu-1} \exp\left(\frac{-pz^2}{t(z-t)}\right) dt, \quad (11)$$

$\text{Re}(\mu) < 0, \text{Re}(p) > 0$ and for $(m-1) < \text{Re}(\mu) < m, (m = 1, 2, 3, \dots)$.

$$\begin{aligned} D_z^{\mu,p} [f(z)] &= \frac{d^m}{dz^m} D_z^{\mu-m} [f(z)] \\ &= \frac{d^m}{dz^m} \left[\frac{1}{\Gamma(-\mu+m)} \int_0^z f(t)(z-t)^{-\mu+m-1} \exp\left(\frac{-pz^2}{t(z-t)}\right) dt \right], \quad (12) \end{aligned}$$

where the path of integration is a line from 0 to z in the complex t -plane. For the case $p = 0$, we obtain the classical Reimann-Liouville fraction derivative operator.

2. Main Result

The extended Mittag-Leffler function can be given as:

$$\begin{aligned} E_{\alpha,p}^{(\gamma,c);q}(z;p) &= \sum_{n=0}^{\infty} \frac{B_p(\gamma+nq, c-\gamma)}{B(\gamma, c-\gamma)} \frac{(c)_{nq}}{\Gamma(\alpha n + \beta)} \cdot \frac{z^n}{n!}, \quad (13) \\ &\quad p \geq 0, \text{Re}(c) > 0, \text{Re}(\gamma) > 0, q < \text{Re}(\alpha) + 1, \end{aligned}$$

where

$$\begin{aligned} B_p(x,y) &= \int_0^1 u^{x-1} (1-u)^{y-1} \exp\left[\frac{-p}{u(1-u)}\right] du, \quad (14) \\ &\quad \text{Re}(p) > 0, \text{Re}(x) > 0, \text{Re}(y) > 0. \end{aligned}$$

The above Mittag-Leffler function can be derived by using the following relations which are given by Chaudhary et al. (2004), Chaudhary and Zubair (2001):

$$\frac{(\gamma)_{nq}}{(c)_{nq}} = \frac{B(\gamma + nq, c - \gamma)}{B(\gamma, c - \gamma)},$$

$$E_{\alpha, \beta}^{\gamma, q}(z) = \sum_{n=0}^{\infty} \frac{(\gamma)_{qn}}{\Gamma(\alpha n + \beta)} \cdot \frac{(c)_{nq}}{(c)_{nq}} \frac{z^n}{n!}. \quad (15)$$

Now, we state some theorems on this function

3. Fractional Derivative of Extended Mittag-Leffler function

Theorem 1.

Let $p \geq 0$, $\operatorname{Re}(\mu) > 0$, $\operatorname{Re}(\lambda) > 0$, $\operatorname{Re}(\alpha) > 0$, $\operatorname{Re}(\beta) > 0$, where $q < \operatorname{Re}(\alpha) + 1$, then

$$D_z^{\lambda - \mu, p} \left[z^{\lambda - 1} E_{\alpha, \beta}^{(\mu); q}(z^q; p) \right]$$

$$= \frac{z^{\mu - 1}}{\Gamma(\mu - \lambda)} B(\lambda, \mu - \lambda) E_{\alpha, \beta}^{(\lambda, \mu); q}(z^q; p), \quad (16)$$

Proof:

Taking L.H.S. and using the extended Riemann-Liouville fractional derivative

$$D_z^{\lambda - \mu, p} \left[z^{\lambda - 1} E_{\alpha, \beta}^{(\mu); q}(z^q; p) \right]$$

$$= \frac{1}{\Gamma(\mu - \lambda)} \int_0^z t^{\lambda - 1} E_{\alpha, \beta}^{(\mu); q}(t^q; p) (z - t)^{-\lambda + \mu - 1} \exp\left[\frac{-p z^2}{t(z - t)}\right] dt,$$

on putting $t = zu$ and using Equation (4), we obtain

$$\frac{z^{\mu - 1}}{\Gamma(\mu - \lambda)} \int_0^1 u^{\lambda - 1} (1 - u)^{-\lambda + \mu - 1} \exp\left[\frac{-p}{u(1 - u)}\right] \sum_{k=0}^{\infty} \frac{(\mu)_{kq}}{\Gamma(\alpha k + \beta)} \frac{z^{qk} u^{qk}}{k!} du.$$

After changing the order of summation and integration, we get the desired result.

Theorem 2.

$$\frac{d^n}{dz^n} \left\{ E_{\alpha, \beta}^{(\gamma, c); q}(z; p) \right\} = (\gamma)_{qn} E_{\alpha, \beta + nq}^{\gamma + nq, c + nq}(z; p). \quad (17)$$

Proof:

Taking first derivative with respect to z of Equation (14), we get

$$\begin{aligned} \frac{d}{dz} \left[E_{\alpha, \beta}^{(\gamma, c); q} (z; \rho) \right] &= \sum_{n=0}^{\infty} \frac{B_{\rho}(\gamma + (n+1)q, c - \gamma)}{B(\gamma, c - \gamma)} \frac{(c)_{(n+1)q}}{\Gamma(\alpha(n+1) + \beta)} \frac{z^n}{n!} \\ &= \sum_{n=0}^{\infty} \frac{B_{\rho}(\gamma + q + nq, c - \gamma)}{B(\gamma, c - \gamma)} \frac{(c)_q (c+q)_{nq}}{\Gamma(\alpha(n+1) + \beta)} \frac{z^n}{n!} \\ &= (\gamma)_q E_{\alpha, \beta + \alpha}^{(\gamma+q; c+q); q} (z; \rho). \end{aligned}$$

Further, applying the process n times, we get the result.

Corollary 1.

For the extended Mittag-Leffler function, the following differentiation formula holds:

$$\frac{d^n}{dz^n} \left[z^{\beta-1} E_{\alpha, \beta}^{(\gamma, c); q} (\lambda z^{\alpha}; \rho) \right] = z^{\beta-n-1} E_{\alpha, \beta-n}^{(\gamma, c); q} (\lambda z^{\alpha}; \rho),$$

in Equation (17). We put $z = \lambda z^{\alpha}$ and multiply by $z^{\beta-1}$. Then taking z derivative n times, we get required result.

Similarly, another interesting derivative formula for the extended Mittag-Leffler function is

$$E_{\alpha, \beta}^{(\gamma, c); q} (z; \rho) = \beta E_{\alpha, \beta+1}^{(\gamma, c); q} (z; \rho) + \alpha z \frac{d}{dz} E_{\alpha, \beta+1}^{(\gamma, c); q} (z; \rho), \quad (18)$$

4. Integral Representation of Extended Mittag-Leffler function**Theorem 3.**

For Extended MittagLeffler function, we have

$$E_{\alpha, \beta}^{(\gamma, c); q} (Z; \rho) = \frac{1}{B(\gamma, c - \gamma)} \int_0^1 t^{\gamma-1} (1-t)^{c-\gamma-1} e^{-\frac{\rho}{t(1-t)}} E_{\alpha, \beta}^{c; q} (t^{\alpha} z) dt, \quad (19)$$

where $p \geq 0, \operatorname{Re}(c) > 0, \operatorname{Re}(\gamma) > 0, \operatorname{Re}(\alpha) > 0, \operatorname{Re}(\beta) > 0, q < \operatorname{Re}(\alpha) + 1$.

Proof:

Using Equation (14) in Equation (13), we get

$$E_{\alpha,\beta}^{(\gamma,c);q} = \sum_{k=0}^{\infty} \int_0^1 t^{\gamma+kq-1} (1-t)^{c-\gamma-1} e^{-p/t(1-t)} \frac{(c)_{kq}}{B(\gamma,c-\gamma)} \cdot \frac{1}{\Gamma(\alpha k + \beta)} \cdot \frac{z^k}{k!} dt.$$

On interchanging order of summation and integration in the above equation, we get

$$\int_0^1 t^{\gamma-1} (1-t)^{c-\gamma-1} e^{-\frac{p}{t(1-t)}} \sum_{k=0}^{\infty} \frac{(c)_{kq}}{B(\gamma,c-\gamma)} \cdot \frac{(t^q z)^k}{\Gamma(\alpha k + \beta) k!} dt, \quad (20)$$

and using Equation (4) in Equation (20), we get the desired result.

Corollary 2.

$$\begin{aligned} & E_{\alpha,\beta}^{(\gamma,c);q}(z; \rho) \\ &= \frac{1}{B(\gamma,c-\gamma)} \int_0^{\infty} \frac{u^{\gamma-1}}{(u+1)^c} \exp\left(\frac{-p(1+u)^2}{u}\right) E_{\alpha,\beta}^{c;q}\left[\left(\frac{u}{1+u}\right)^q z; \rho\right] du. \end{aligned} \quad (21)$$

The above result can be prove by putting $t = \frac{u}{1+u}$ in Theorem 3.

Corollary 3.

$$\begin{aligned} & E_{\alpha,\beta}^{(\gamma,c);q}(z; \rho) \\ &= \frac{1}{B(\gamma,c-\gamma)} \left[2 \int_0^{\pi/2} \sin^{2\gamma-1}\theta \cos^{2(c-\gamma)-1}\theta \exp\left(\frac{-p}{\sin^2\theta \cos^2\theta}\right) \cdot E_{\alpha,\beta}^{c;q}(\sin^{2q}\theta z) d\theta \right]. \end{aligned} \quad (22)$$

Taking $t = \sin^2\theta$ in theorem 3, we get above trigonometric form of Extended Beta function.

5. Integral transform of extended Mittag-Leffler function**Theorem 4.**

The Mellin transform of the extended Mittag-Leffler function is given by

$$\begin{aligned} & M\left[E_{\alpha,\beta}^{(\gamma,c);q}(z; \rho); s\right] \\ &= \frac{\Gamma s \Gamma(c+s-\gamma)}{\Gamma \gamma \Gamma(c-\gamma)} {}_2\psi_2\left[\begin{matrix} (c,q), (\gamma+s,q) \\ (\beta,\alpha), (c+2s,q) \end{matrix}; z\right]. \end{aligned} \quad (23)$$

Proof:

We start with

$$M \left[E_{\alpha, \beta}^{(\gamma, c); q} (z; \rho); s \right] = \int_0^{\infty} \rho^{s-1} E_{\alpha, \beta}^{(\gamma, c); q} (z; \rho) d\rho, \quad (24)$$

and using Equation (19) in Equation (24), we get

$$\begin{aligned} M \left[E_{\alpha, \beta}^{(\gamma, c); q} (z; \rho); s \right] \\ = \frac{1}{B(\gamma, c-\gamma)} \int_0^{\infty} \rho^{s-1} \left[\int_0^1 t^{\gamma-1} (1-t)^{c-\gamma-1} e^{-\frac{\rho}{t(1-t)}} \right] E_{\alpha, \beta}^{c; q} (t^q z) dt d\rho. \end{aligned}$$

Now, changing the order of integration and putting $u = \frac{\rho}{t(1-t)}$, we get

$$\begin{aligned} M \left[E_{\alpha, \beta}^{(\gamma, c); q} (z; \rho); s \right] \\ = \frac{\Gamma s}{B(\gamma, c-\gamma)} \int_0^1 t^{\gamma+s-1} (1-t)^{c+s-\gamma-1} \cdot \sum_{k=0}^{\infty} \frac{(c)_{kq}}{\Gamma(\alpha k + \beta)} \cdot \frac{(t^q z)^k}{k!} dt. \end{aligned}$$

Finally, using Beta function, [Equation (5)] in the above case, we get the desired result.

Corollary 4.

Taking $s=1$ in Equation (23), we get

$$\int_0^{\infty} E_{\alpha, \beta}^{(\gamma, c); q} (z; \rho) d\rho = \frac{\Gamma(c+1-\gamma)}{\Gamma\gamma \Gamma(c-\gamma)} {}_2\Psi_2 \left[\begin{matrix} (c, q), (\gamma+1, q) \\ (\beta, \alpha), (c+2, q) \end{matrix}; z \right]. \quad (25)$$

Corollary 5.

Taking Inverse Mellin transform

$$\begin{aligned} E_{\alpha, \beta}^{(\gamma, c); q} (z; \rho) d\rho \\ = \frac{1}{2\pi i} \frac{1}{\Gamma\gamma \Gamma c-\gamma} \int_{\gamma-i\infty}^{\gamma+i\infty} \Gamma s \Gamma(c+s-\gamma) {}_2\Psi_2 \left[\begin{matrix} (c, q), (\gamma+s, q) \\ (\beta, \alpha), (c+2s, q) \end{matrix}; z \right] \cdot \rho^{-s} ds, \quad (26) \end{aligned}$$

Similarly, Laplace and Whittaker transform [4] of Extended Mittag-Leffler function are as follows:

$$\begin{aligned} L \left[z^{b-1} E_{\alpha, \beta}^{(\gamma, c); q} (z^q; p); s \right] &= \int_0^{\infty} z^{b-1} e^{-sz} E_{\alpha, \beta}^{(\gamma, c); q} (z^q; p) dz \\ &= \frac{\Gamma b}{s^b} \sum_{k=0}^{\infty} \frac{B_p(\gamma + kq, c - \gamma)}{B(\gamma, c - \gamma)} \frac{(c)_{kq}}{k!} \frac{(b)_k}{s^k}, \end{aligned} \quad (27)$$

and

$$\begin{aligned} &\int_0^{\infty} e^{-ft/2} t^{\varepsilon-1} W_{\lambda, \psi}(ft) E_{\alpha, \beta}^{(\gamma, c); q} (w t^{\eta}) dt \\ &= f^{-\varepsilon} \sum_{k=0}^{\infty} \frac{B_p(\gamma + kq, c - \gamma)}{B(\gamma, c - \gamma)} \frac{(c)_{kq}}{\Gamma(\alpha k + \beta)} \frac{1}{k!} \left(\frac{\omega}{f^{\eta}} \right)^k \frac{\Gamma\left(\varepsilon + \eta k - \psi + \frac{1}{2}\right) \Gamma\left(\varepsilon + \eta k + \psi + \frac{1}{2}\right)}{\Gamma(\varepsilon + \eta k - \lambda + 1)}. \end{aligned} \quad (28)$$

Using the definition of extended Mittag-Leffler function [Equation (14)] in middle term of Equation (27), and changing the order of integration and summation, and using Laplace Transform

$$\int_0^{\infty} e^{-st} t^{n-1} dt = \frac{\Gamma n}{s^n},$$

we get the required result. Taking L.H.S. of Equation (28) and put $ft = v$ and using definition of Extended Mittag-Leffler function changing order of summation and integration and using Equation (9) we get the desired result.

6. Special Cases

- (i) If we put $q=1$, in above Theorems 1, 3, and 4, it reduces to the result given recently by Özarlan and Yilmaz (2014).
- (ii) If we put $c=q=1$, and $p=0$, in the above Theorems 1, 3, and 4, it reduces to Prabhakar function (1971).

7. Conclusion

In this investigation, we established and evaluated some fractional derivative formulas involving extended Mittag Leffler function by using extended beta function and also evaluated integral representation of this extended Mittag-leffler function. Other than this, we obtained some integral transforms in terms of generalized Wright function and extended beta function. The approach presented in this investigation is general but can be extended to establish other properties of special functions.

Acknowledgement

We thank both referees and the Editor-in-Chief Professor Aliakbar Montazer Haghighi for their valuable suggestions and constructive comments towards the improvement of this present investigation.

REFERENCES

- Chaudhry, M. A., Qadir, A., Srivastava, H. M., & Paris, R. B. (2004). Extended hypergeometric and confluent hypergeometric functions. *Applied Mathematics and Computation*, Vol. 159, No.2, pp. 589-602.
- Chaudhry, M. A. & Zubair, S. M. (2001). *On a class of incomplete gamma functions with applications*. Chapman and Hall (CRC Press Company), Boca Raton, London, New York and Washington, D.C.
- Haubold, H. J., Mathai, A. M., & Saxena, R. K. (2011). Mittag-Leffler functions and their applications. *Journal of Applied Mathematics*, Article ID 298628, doi: 10.1155/2011/298628
- Khan, M. A., & Ahmed, S. (2013). On some properties of the generalized Mittag-Leffler function, *Springer Plus*, Vol. 2, No.1, pp.1-9.
- Mittag-Leffler, G. M. (1903). Une généralisation de l'intégrale de Laplace-Abel. *CR Acad. Sci. Paris (Ser. II)*, Vol. 137, pp. 537-539.
- Özarslan, M. A., & Yılmaz, B. (2014). The extended Mittag-Leffler function and its properties. *Journal of Inequalities and Applications*, Vol. 2014, No.1, pp.85.
- Özarslan, M. A., & Özergin, E. (2010). Some generating relations for extended hypergeometric functions via generalized fractional derivative operator. *Mathematical and Computer Modelling*, Vol. 52, No.9, pp.1825-1833.
- Rainville, E. D. (1960). *Special functions*, Vol. 442, Macmillan, New York.
- Samko, S.G., Kilbas, A.A., Marichev, O.I. (1993). *Fractional Integrals and Derivatives: Theory and Applications*, Gordon & Breach, New York
- Shukla, A. K. & Prajapati, J. C. (2007). On a generalization of Mittag-Leffler function and its properties. *Journal of Mathematical Analysis and Applications*, Vol. 336, No.2, pp.797-811.
- Whittaker, E. T., & Watson, G. N. (1996). *A course of modern analysis*, Cambridge University press, Cambridge.
- Wiman, A. (1905). Über den Fundamentalsatz in der Theorie der Funktionen $E_\alpha(x)$, *Acta Mathematica*, Vol. 29, No.1, pp.191-201.