



12-2016

Some Results on f-Simultaneous Chebyshev Approximation

Sh. Al-Sharif
Yarmouk University

Kh. Qaraman
Yarmouk University

Follow this and additional works at: <https://digitalcommons.pvamu.edu/aam>



Part of the [Numerical Analysis and Computation Commons](#)

Recommended Citation

Al-Sharif, Sh. and Qaraman, Kh. (2016). Some Results on f-Simultaneous Chebyshev Approximation, *Applications and Applied Mathematics: An International Journal (AAM)*, Vol. 11, Iss. 2, Article 31. Available at: <https://digitalcommons.pvamu.edu/aam/vol11/iss2/31>

This Article is brought to you for free and open access by Digital Commons @PVAMU. It has been accepted for inclusion in *Applications and Applied Mathematics: An International Journal (AAM)* by an authorized editor of Digital Commons @PVAMU. For more information, please contact hvkoshy@pvamu.edu.



Some Results on f -Simultaneous Chebyshev Approximation

¹Sh. Al-Sharif & ²Kh. Qaraman

Mathematics Department
Yarmouk University
Irbid, Jordan

¹sharifa@yu.edu.jo; ²math811@yahoo.com

Received: February 26, 2015; Accepted: October 30, 2015

Abstract

Let X be Hausdorff topological vector space and f be a real valued continuous function on X . In this paper we introduce and study the concept of f -simultaneous approximation of a nonempty subset K of X as a generalization to the problem of simultaneous approximation. Further we present some results regarding f -simultaneous approximation in the quotient space.

Keywords: f -best simultaneous approximation; f -simultaneous Chebyshev approximation; f -simultaneously proximal sets; f -quasi-simultaneously Chebyshev

MSC 2010 No.: 41A65, 41A50

1. Introduction

Let K be a subset of a Hausdorff topological vector space X and f be a real valued continuous function on X . For $x \in X$, set $F_K(x) = \inf_{k \in K} f(x - k)$. A point $k_0 \in K$ is called f -best approximation to x in K if $F_K(x) = f(x - k_0)$. The set $P_K^f(x) = \{k_0 \in K : F_K(x) = f(x - k_0)\}$ denotes the set of all f -best approximations to x in K . Note that this set may be empty. The set K is said to be f -proximal (f -Chebyshev) if for each $x \in X$, $P_K^f(x)$ is non-empty (singleton). The notion of f -best approximation in a vector space X was given by Breckner and Brosowski, and in a Hausdorff topological vector space X by Narang. For a

Hausdorff locally convex topological vector space and a continuous sublinear functional f on X , Breckner, Brosowski and Govindarajulu proved certain results on best approximation relative to the functional f . By using the existence of elements of f -best approximation some results on fixed point were proved by Pai and Veermani.

As a generalization to the problem of simultaneous approximation see Saidi and Singer, we introduce the concept of best f -simultaneous approximation as follows.

Definition 1.

Let f be a real valued continuous function on a Hausdorff topological vector space X . A subset A of X is called f -bounded if there exists $M > 0$ such that $|f(x)| \leq M$ every $x \in A$.

Note that f -bounded set need not be bounded in the classical sense, for example if $f(x) = e^{-x}$, the set $[0, \infty)$ is an f -bounded subset of real numbers.

Definition 2.

Let X be a Hausdorff topological real vector space, f be a real valued continuous function on X and K be a non-empty subset of X . A point $k_0 \in K$ is called f -best simultaneous approximation in K if there exists an f -bounded subset A of X such that

$$F_K(A) = \inf_{k \in K} \sup_{a \in A} |f(a - k)| = \sup_{a \in A} |f(a - k_0)|.$$

The set of all f -best simultaneous approximations to an f -bounded subset A of X in K is denoted by

$$P_K^f(A) = \left\{ k \in K : F_K(A) = \sup_{a \in A} |f(a - k)| \right\}.$$

The set K is called f -simultaneously proximal (f – simultaneously Chebyshev) if for each f -bounded set A in X , $P_K^f(A) \neq \phi$ (singleton).

We note that if $f(x) = \|x\|$ ($f(x) = \|x\| + \epsilon$), then the concept of f -best approximation is precisely best approximation, (best ϵ -approximation) (see Khalil, Rezapour, Singer, and others).

A set K is said to be inf-compact at a point $x \in X$ (Pai and Veermani), if each minimizing sequence in K (i.e. $f(x - k_n) \rightarrow F_K(x)$) has a convergent subsequence in K . The set K is called inf-compact if it is inf-compact at each $x \in X$. A subset K of X is called f -compact (Moghaddam) if for every sequence $\{k_n\}$ in K , there exist a subsequence $\{k_{n_i}\}$ of $\{k_n\}$ and $k_0 \in K$ such that $f(k_{n_i} - k_0) \rightarrow 0$. It is easy to see that if K is f -compact or inf-compact, then K is f -simultaneously proximal.

In this paper we introduce and study the concept of f -simultaneous approximation of a subspace K of a Hausdorff topological real vector space X , existence and uniqueness. Certain results regarding f -simultaneous approximation in quotient spaces is obtained by generalizing some of the results in Moghaddam.

Throughout this paper X is a Hausdorff topological real vector space and f is a real valued continuous function on X .

2. f -Simultaneous Approximation

In this section we give some characterization of f -proximal sets in X . We begin by the following definitions:

Definition 3.

A function $f : X \rightarrow \mathbb{R}$ is called

- (1) absolutely subadditive if $|f(x + y)| \leq |f(x)| + |f(y)|$ for all $x, y \in X$.
- (2) absolutely homogeneous if $f(\alpha x) = |\alpha| f(x)$, for all $x \in X$ and all $\alpha \in \mathbb{R}$.

Definition 4.

A subset K of X is called f -closed if for all sequences $\{k_m\}$ of K and for all $x \in X$ such that $f(x - k_m) \rightarrow 0$ we have $x \in K$.

Theorem 5.

Let K be a subset of X . Then,

- (1) $F_{K+y}(A + y) = F_K(A)$, for all f -bounded sets $A \subset X$, $y \in X$.
- (2) $P_{K+y}^f(A + y) = P_K^f(A) + y$, for all f -bounded sets $A \subset X$, $y \in X$.
- (3) K is f -simultaneously proximal (f -simultaneously Chebyshev) if and only if $K + y$ is f -simultaneously proximal (f -simultaneously Chebyshev) for every $y \in X$.

More over if f is absolutely homogeneous function, then

- (4) $F_{\lambda K}(\lambda A) = |\lambda| F_K(A)$, for all f -bounded sets $A \subset X$ and $\lambda \in \mathbb{R}$.
- (5) $P_{\lambda K}^f(\lambda A) = \lambda P_K^f(A)$, for all f -bounded sets $A \subset X$ and $\lambda \in \mathbb{R}$.
- (6) K is f -simultaneously proximal (f -simultaneously Chebyshev) if and only if λK is f -simultaneously proximal (f -simultaneously Chebyshev), $\lambda \in \mathbb{R}$.

Proof:

- (1) Let $A \subset X$, f -bounded set. Then

$$F_{K+y}(A + y) = \inf_{w \in K} \sup_{a \in A} |f((a + y) - (w + y))| = F_K(A).$$

- (2) The equation

$$\sup_{a \in A} |f(a - k_0)| = \inf_{k \in K} \sup_{a \in A} |f((a + y) - (k + y))| = \inf_{k \in K} \sup_{a \in A} |f(a - k)|,$$

implies that $k_0 + y \in P_{K+y}^f(A + y)$ if and only if $k_0 \in P_K^f(A)$. Thus,

$$P_{K+y}^f(A + y) = P_K^f(A) + y.$$

(3) Follows immediately from part two.

(4) Let $A \subset X$, be f -bounded set, $\lambda \in \mathbb{R}$. Then,

$$F_{\lambda K}(\lambda A) = \inf_{k \in K} \sup_{a \in A} |f(\lambda a - \lambda k)| = |\lambda| \inf_{k \in K} \sup_{a \in A} |f(a - k)| = |\lambda| F_K(A).$$

(5) If $\lambda = 0$, we are done. If $\lambda \neq 0$ and $k_0 \in P_{\lambda K}^f(\lambda A)$, then $k_0 \in \lambda K$ and

$$\sup_{a \in A} |f(\lambda a - k_0)| = \inf_{k \in K} \sup_{a \in A} |f(\lambda a - \lambda k)|.$$

This implies that

$$\sup_{a \in A} \left| f\left(a - \frac{1}{\lambda} k_0\right) \right| = F_K(A),$$

which implies that $\frac{1}{\lambda} k_0 \in P_K^f(A)$.

(6) Follows immediately from part 5. \square

Theorem 6.

Let f be an absolutely homogeneous real valued function on X and M be a subspace of X . Then

(1) $F_M(\lambda A) = |\lambda| F_M(A)$, for all for all f -bounded sets $A \subset X$ and $\lambda \in \mathbb{R} - \{0\}$.

(2) $P_M^f(\lambda A) = \lambda P_M^f(A)$, for all for all f -bounded sets $A \subset X$ and $\lambda \in \mathbb{R} - \{0\}$.

Proof:

(1) Let $A \subset X$ be an f -bounded set and $\lambda \neq 0 \in \mathbb{R}$. Then,

$$F_M(\lambda A) = \inf_{m \in M} \sup_{a \in A} |f(\lambda a - m)| = |\lambda| \inf_{m' \in M} \sup_{a \in A} |f(a - m')| = |\lambda| F_M(A).$$

(2) Let $m_0 \in P_M^f(\lambda A)$. Then,

$$\begin{aligned} \sup_{a \in A} |\lambda| \left| f\left(a - \frac{1}{\lambda} m_0\right) \right| &= \sup_{a \in A} |f(\lambda a - m_0)| \\ &= \inf_{m \in M} \sup_{a \in A} |f(\lambda a - m)| \\ &= \inf_{m' \in M} \sup_{a \in A} |\lambda| \left| f(a - m') \right|. \end{aligned}$$

Therefore,

$$\sup_{a \in A} \left| f\left(a - \frac{1}{\lambda} m_0\right) \right| = \inf_{m' \in M} \sup_{a \in A} |f(a - m')| = F_M(A),$$

for all $\lambda \in \mathbb{R} - \{0\}$, which implies that $\frac{1}{\lambda} m_0 \in P_M^f(A)$, and so $m_0 \in \lambda P_M^f(A)$. \square

For a subset K of X , let us define \widehat{K}_F be such that

$$\widehat{K}_F = \left\{ A \subset X : F_K(A) = \sup_{a \in A} f(a) \right\}.$$

Using this we prove the following theorem characterizing f -simultaneously proximal sets.

Theorem 7.

Let K be a subspace of X . Then K is f -simultaneously proximal in X if and only if every f -bounded subset A of X can be written as $B + k$ for some $k \in K$ and $B \in \widehat{K}_F$.

Proof:

Suppose the condition hold. Let $A \subset X$ be an f -bounded subset of X . By assumption there exists $k_0 \in K$ and $B \in \widehat{K}_F$ such that $A = B + k_0$. Hence $A - k_0 \in \widehat{K}_F$. Therefore,

$$\begin{aligned} \sup_{a \in A} |f(a - k_0)| &= F_K(A - k_0) \\ &= \inf_{k \in K} \sup_{a \in A} |f(a - k_0 - k)| \\ &= \inf_{k' \in K} \sup_{a \in A} |f(a - k')| = F_K(A). \end{aligned}$$

Hence K is f -simultaneously proximal.

Conversely, suppose K is f -simultaneously proximal and $A \subset X$ be an f -bounded subset of X . Then there exists $k_0 \in K$ such that

$$\sup_{a \in A} |f(a - k_0)| = \inf_{k \in K} \sup_{a \in A} |f(a - k)| = \inf_{k' \in K} \sup_{a \in A} |f(a - (k' + k_0))|.$$

where $k = k' + k_0$. Hence,

$$\sup_{a \in A} |f(a - k_0)| = F_K(A - k_0).$$

Consequently $A - k_0 \in \widehat{K}_F$. So there exists $B \in \widehat{K}_F$ such that $A - k_0 = B$ or $A = B + k_0$. \square

Theorem 8.

Let f be a real valued continuous function on X such that $x = 0$ iff $f(x) = 0$. If K is f -simultaneously proximal, then K is f -closed.

Proof:

Let $\{k_m\}$ be a sequence of K and $x \in X$, such that $f(x - k_m) \rightarrow 0$. Taking $A = \{x\}$, we have

$$F_K(A) = \inf_{k \in K} \sup_{a \in A} |f(a - k)| \leq |f(x - k_m)| \rightarrow 0.$$

Since K is f -simultaneously proximal, there exists $k_0 \in K$ such that

$$F_K(A) = |f(x - k_0)| = 0.$$

Hence, $f(x - k_0) = 0$. Using assumption it follows that $x - k_0 = 0$. Hence $x = k_0 \in K$ and K is f -closed. \square

3. f -Simultaneous Approximation in Quotient Space

Let M a closed subspace of X . Then a function $\tilde{f} : (X/M) \rightarrow \mathbb{R}$ can be defined as follows:

$$\tilde{f}(x + M) = \inf_{y \in M} |f(x + y)|.$$

Proposition 9.

Let M a closed subspace of X . If A is f -bounded in X , then A/M is \tilde{f} -bounded in X/M .

Proof:

Let A be an f -bounded subset in X . Since M is a subspace, for $x + M \in A/M$

$$\left| \tilde{f}(x + M) \right| = \inf_{y \in M} |f(x + y)| \leq |f(x)|.$$

Consequently since A is an f -bounded subset of X , it follows that A/M is \tilde{f} -bounded in X/M . \square

Theorem 10.

Let M a closed subspace of X . If B is \tilde{f} -bounded in X/M , then there exists an f -bounded subset A of X such that $B = A/M$.

Proof:

Let B be a non empty \tilde{f} -bounded in X/M . Let $C = \bigcup_{b \in B} b$. Claim $B = \{\bar{x} = x + M : x \in C\}$.

Indeed if $b \in B$, then $b = x_b + M$ for some $x_b \in X$. But M is a subspace. Thus $x_b = x_b + 0 \in x_b + M \subset C$. Hence $b = x_b + M \in \{\bar{x} = x + M : x \in C\}$ and $B \subseteq \{\bar{x} = x + M : x \in C\}$. Similarly if $x \in C$, then $x \in b_x + M$ for some $b_x + M \in B$. This implies that $x = b_x + m_x$ for some $m_x \in M$. Hence $x + M = b_x + m_x + M = b_x + M \in B$. Therefore $\{\bar{x} = x + F : x \in C\} \subseteq B$.

Now clearly C is not bounded unless M is trivial. Note that B is \tilde{f} -bounded. So there exists $K > 0$ such that $|\tilde{f}(b)| \leq K$ for all $b \in B$. Consider the set $A = \{x \in C : |f(x)| \leq K + 1\} \subseteq C$. Now we claim that for all $x \in C$,

$$\bar{x} \cap A = (x + M) \cap A \neq \phi.$$

Given $x \in C$. Since

$$|\tilde{f}(x + M)| = \inf_{m \in M} |f(x + m)| \leq K,$$

there exists $m_x \in M$ such that $|f(x + m_x)| < K + 1$. But $x + m_x \in x + M \subseteq C$. Hence $x + m_x \in (x + M) \cap A \neq \phi$. Claim $B = A/M$. Since $A \subseteq C$, we have $A/M \subseteq \{\bar{x} = x + M : x \in C\} = B$. To show the other inclusion, let $b \in B = \{\bar{x} = x + M : x \in C\}$. Then $b = x_b + M$ for some $x_b \in C$. But $(x_b + M) \cap A \neq \phi$. Thus there exists $a \in A$ such that $a = x_b + m_a \in x_b + M$. Therefore $b = x_b + M = (x_b + m_a) + M = a + M \in A/M$. Hence $B \subseteq A/M$. Consequently $A/M = B$. \square

Theorem 11.

Let K be a subspace of X and M be a closed f -proximal subspace of K . If k_0 is a point of f -best simultaneous approximation to $A \subset X$ in K , then $k_0 + M$ is an \tilde{f} -best simultaneous approximation to A/M in K/M .

Proof:

Suppose $k_0 + M$ is not \tilde{f} -best simultaneous approximation to A/M in K/M . Then for at least $k \in K$, say $k_1 \in K$, we have

$$\sup_{a \in A} \tilde{f}(a - k_1 + M) < \sup_{a \in A} \tilde{f}(a - k_0 + M).$$

Since

$$\sup_{a \in A} \tilde{f}(a - k_0 + M) = \sup_{a \in A} \inf_{m \in M} |f(a - k_0 + m)| \leq \sup_{a \in A} |f(a - k_0)|,$$

we have

$$\sup_{a \in A} \tilde{f}(a - k_1 + M) = \sup_{a \in A} \inf_{m \in M} |f(a - k_1 + m)| < \sup_{a \in A} |f(a - k_0)|.$$

But M is f -proximal, so for some $m_0 \in M$ we have

$$\sup_{a \in A} |f(a - k_1 + m_0)| = \sup_{a \in A} \inf_{m \in M} |f(a - k_1 + m)| < \sup_{a \in A} |f(a - k_0)|.$$

Since $M \subset K$, it follows that $k_1 - m_0 \in K$. Therefore k_0 not f -best simultaneous approximation to A in K , which is a contradiction. \square

Corollary 12.

Let K be a subspace of X and M is a closed f -proximal subspace of K . If K is f -simultaneously proximal in X , then K/M is \tilde{f} -simultaneously proximal in X/M .

Proof:

Let B be an \tilde{f} -bounded subset of X/M . Then by Theorem 10, there exists f -bounded subset $A \subset X$ such that $B = A/M$. If K is f -simultaneously proximal in X , then there exists at least $k_0 \in K$ such that k_0 is f -best simultaneous approximation to A in K . By Theorem 11, $k_0 + M$ is an \tilde{f} -best simultaneous approximation to A/M in K/M , so K/M is \tilde{f} -simultaneously proximal in X/M . \square

Theorem 13.

Let K be a subspace of X and M is a closed f -proximal subspace of K . If K/M is \tilde{f} -simultaneously proximal in X/M , then K is f -simultaneously proximal in X .

Proof:

Let A be an f -bounded subset of X . By Proposition 9, A/M is \tilde{f} -bounded in X/M . Since K/M is \tilde{f} -simultaneously proximal in X/M , then there exists $k_0 + M \in K/M$ such that $k_0 + M$ is \tilde{f} -best simultaneous approximation to A/M from K/M , so

$$\begin{aligned} \sup_{a \in A} \tilde{f}(a - k_0 + M) &= \inf_{k \in K} \sup_{a \in A} \tilde{f}(a - k + M) \\ &= \inf_{k \in K} \sup_{a \in A} \inf_{m \in M} |f(a - k + m)| \\ &\leq \inf_{k \in K} \sup_{a \in A} |f(a - k + m)| \\ &= \inf_{k \in K} \sup_{a \in A} |f(a - k')|, \end{aligned} \tag{1}$$

where, $k' = k - m \in K$. Since M is f -proximal, there exists $m_0 \in M$ such that

$$\sup_{a \in A} |f(a - k_0 - m_0)| = \sup_{a \in A} \inf_{m \in M} |f(a - k_0 + m)| = \sup_{a \in A} \tilde{f}(a - k_0 + M). \quad (2)$$

Consequently, combining (1) and (2) since $M \subset K$, it follows that

$$\begin{aligned} \sup_{a \in A} |f(a - k_0 - m_0)| &\leq \inf_{k' \in K} \sup_{a \in A} |f(a - k')| \\ &\leq \sup_{a \in A} |f(a - k_0 - m_0)| \end{aligned}$$

Hence,

$$\sup_{a \in A} |f(a - k_0 + m_0)| = \inf_{k \in K} \sup_{a \in A} |f(a - k')|$$

So $k_0 + m_0$ is an f -best simultaneous approximation to A from K and K is f -simultaneously proximal in X . \square

Theorem 14.

Let W and M be two subspaces of X . If M is a closed f -proximal subspace of X , then the following assertions are equivalent:

- (1) W/M is \tilde{f} -simultaneously proximal in X/M ,
- (2) $W + M$ is f -simultaneously proximal in X .

Proof:

(1) \Rightarrow (2). Since $(W + M)/M = W/M$ and M are f -simultaneously proximal, using Theorem 13, it follows that $W + M$ is f -simultaneously proximal in X .

(2) \Rightarrow (1). Since $W + M$ is f -simultaneously proximal and $M \subseteq W + M$, by Corollary 12, $(W + M)/M = W/M$ is simultaneously f -proximal. \square

Theorem 15.

Let K, M be two subspaces of X such that, $M \subset K$. If M is closed f -simultaneously proximal in X and K is f -simultaneously Chebyshev in X , then K/M is \tilde{f} -simultaneously Chebyshev in X/M .

Proof:

Suppose not. Then there exists A , f -bounded subset of X such that $A/M \in X/M$ is \tilde{f} -bounded and $k_1 + M, k_2 + M \in P_{K/M}^{\tilde{f}}(A/M)$ such that $k_1 + M \neq k_2 + M$. Thus $k_1 - k_2 \notin M$. Since M is an f -simultaneously proximal in X , then

$$P_M^f(A - k_1) \neq \phi, \text{ and } P_M^f(A - k_2) \neq \phi.$$

Let $m_1 \in P_M^f(A - k_1)$, and $m_2 \in P_M^f(A - k_2)$. By Theorem 13, $k_1 + m_1$ and $k_2 + m_2$ are f -best simultaneous approximations to A from K . Since K is f -simultaneously Chebyshev in X , then $k_1 + m_1 = k_2 + m_2$ and hence $k_1 - k_2 = m_1 - m_2 \in M$, which is a contradiction. \square

Definition 16.

A subset K of X is called f -quasi-simultaneously Chebyshev if $P_K^f(A)$ is non empty and f -compact set in X , for all f -bounded subsets of X .

Theorem 17.

Let M be a closed f -simultaneously proximal subspace of X and K is f -quasi-simultaneously Chebyshev of X such that $M \subset K$. Then K/M is \tilde{f} -quasi-simultaneously Chebyshev in X/M .

Proof:

Since K is f -simultaneously proximal in X , By Corollary 12, K/M is \tilde{f} -simultaneously proximal in X/M . Let B be an \tilde{f} -bounded subset of X/M . Then, by Theorem 10, $B = A/M$ for an f -bounded subset A of X . If $(k_n + M)$ a sequence in $P_{K/M}^{\tilde{f}}(A/M)$, by the proof of Theorem 13, for every n , there exists $m_n \in M$ such that $k_n + m_n = k'_n \in P_K^f(A)$. But since M is a subspace, we have

$$k'_n + M = k_n + m_n + M = k_n + M.$$

Since K is f -quasi-simultaneously Chebyshev in X , the sequence $\{k_n\}$ has a subsequence $\{k_{n_i}\}$ such that $f(k_{n_i} - k_0) \rightarrow 0$ for some $k_0 \in P_K^f(A)$. But

$$\tilde{f}(k_{n_i} - k_0 + M) \leq |f(k_{n_i} - k_0)| \rightarrow 0.$$

Therefore,

$$\tilde{f}(k_{n_i} - k_0 + M) \rightarrow 0$$

and

$$\tilde{f}((k_{n_i} + M) - (k_0 + M)) \rightarrow 0.$$

Hence, $P_{K/M}^{\tilde{f}}(A/M)$ is \tilde{f} -compact and K/M is \tilde{f} -quasi-simultaneously Chebyshev. This complete the proof. \square

Definition 18.

A topological vector space X is said to have the f -property if every f -bounded sequence in X has an f -convergent subsequence, where f is a real valued continuous function on X .

Note that the space $X = l^2$ has the f -property for every projection $f : X \rightarrow \mathbb{R}$, and if $f(x) = \|x\|$, then every finite dimensional Banach space has the f -property.

Proposition 19.

Let f be an absolutely homogeneous subadditive continuous real valued function on a topological vector space X and K be an f -closed subspace of X . Then for any f -bounded subset A of X , $P_K^f(A)$ is f -closed.

Proof:

Let K be an f -closed subspace of X and A be an f -bounded subset of X . If $\{k_m\}$ is a sequence in $P_K^f(A)$ and $x \in X$ such that $f(k_m - x) \rightarrow 0$, then $x \in K$ since K is f -closed. Further

$$\begin{aligned} \inf_{k \in K} \sup_{a \in A} |f(a - k)| &= \sup_{a \in A} |f(a - k_m)| \\ &= \sup_{a \in A} |f((a - x) - (k_m - x))| \\ &\geq \sup_{a \in A} ||f(a - x)| - |f(k_m - x)||. \end{aligned}$$

Taking the limit as $m \rightarrow \infty$, we get

$$\inf_{k \in K} \sup_{a \in A} |f(a - k)| \geq \sup_{a \in A} |f(a - x)|.$$

Consequently,

$$\inf_{k \in K} \sup_{a \in A} |f(a - k)| = \sup_{a \in A} |f(a - x)|.$$

Hence $x \in P_K^f(A)$ and $P_K^f(A)$ is f -closed. \square

Theorem 20.

Let f be a real valued sub-additive continuous function on a topological vector space X that has the f -property and M be a closed subspace of X . If W is a subspace of X such that $W + M$ is f -closed, then the following assertions are equivalent:

- (1) W/M is \tilde{f} -simultaneously quasi-Chebyshev in X/M .
- (2) $W + M$ is f -simultaneously quasi-Chebyshev in X .

Proof:

(1) \Rightarrow (2) Since M is f -simultaneously proximal by Theorem 14, $W + M$ is f -simultaneously proximal in X . Let A be an arbitrary f -bounded set in X . Then $P_{W+M}^f(A) \neq \phi$. Now to show that $P_{W+M}^f(A)$ is f -compact, we need to show that every sequence in $P_{W+M}^f(A)$ has an f -convergent subsequence. Let $\{g_n\}_{n=1}^{\infty}$ be an arbitrary sequence in $P_{W+M}^f(A)$. Then by Theorem 11, for each $n > 1$, $g_n + M \in P_{(W+M)/M}^{\tilde{f}}(A/M)$. Since $P_{(W+M)/M}^{\tilde{f}}(A/M)$ is \tilde{f} -compact, one

can choose $g_0 \in W + M$ with $g_0 + M \in P_{(W+M)/M}^{\tilde{f}}(A/M)$ and $\{g_{n_k} + M\}_{k=1}^{\infty}$ is \tilde{f} -convergent to $g_0 + M$ for some subsequence $\{g_{n_k} + M\}_{k=1}^{\infty}$ of $\{g_n + M\}_{n=1}^{\infty}$. That means,

$$\tilde{f}(g_0 - g_{n_k} + M) = \inf_{m \in M} |f(g_0 - g_{n_k} - m)| \rightarrow 0.$$

Now, since M is f -proximal in X , there exists $m_{n_k} \in M$ such that $m_{n_k} \in P_M^f(g_0 - g_{n_k})$, for every $k \geq 1$, and hence

$$|f(g_0 - g_{n_k} - m_{n_k})| = \inf_{m \in M} |f(g_0 - g_{n_k} - m)|.$$

Therefore,

$$\lim_{k \rightarrow \infty} f(g_0 - g_{n_k} - m_{n_k}) = 0.$$

On the other hand, $\{g_{n_k}\}_{k=1}^{\infty}$ is an f -bounded sequence because $g_n \in P_{W+M}^f(A)$. In fact $|f(g_n)| \leq 2 \sup_{a \in A} |f(a)|$. Since M has the f -property, with out loss of generality, we may assume that for some $m_0 \in M$, $f(m_{n_k} - m_0) \rightarrow 0$. Let $g' = g_0 - m_0$. Then, $g' \in W + M$ and

$$\begin{aligned} f(g' - g_{n_k}) &= f(g_0 - m_0 - g_{n_k}) \\ &\leq f(g_0 - g_{n_k} - m_{n_k}) + f(m_{n_k} - m_0), \end{aligned}$$

$\forall k \geq 1$. Thus $\lim_{k \rightarrow \infty} f(g' - g_{n_k}) = 0$. Since $\{g_{n_k}\}_{k=1}^{\infty} \in P_{W+M}^f(A)$, for every $k \geq 1$, and $P_{W+M}^f(A)$ is f -closed, since $W + M$ is f -closed by Proposition 19, we conclude that $g' \in P_{W+M}^f(A)$. Hence $P_{W+M}^f(A)$ is f -compact.

(2) \Rightarrow (1) Since M and $W + M$ are f -simultaneously proximal and $M \subseteq W + M$, then $(W + M)/M = W/M$ is \tilde{f} -simultaneously proximal in X/M .

Now, let A be an arbitrary f -bounded set in X . Then $P_{W/M}^{\tilde{f}}(A/M)$ is non-empty. So from the hypothesis we have $W + M$ is f -simultaneously quasi-Chebyshev in X , and hence $P_{W+M}^f(A)$ is f -compact in X . Using Theorem 11, we conclude that

$$P_{(W+M)/M}^{\tilde{f}}(A/M) = \pi \left(P_{W+M}^f(A) \right),$$

where $\pi : X \rightarrow X/M$, $\pi(x) = x + M$, is continuous. Consequently $P_{W/M}^{\tilde{f}}(A/M)$ is \tilde{f} -compact. Therefore, W/M is f -simultaneously quasi-Chebyshev in X . \square

Note that Theorem 20 is still true if the restriction $W + M$ is f -closed is replaced by the condition that the function $f(x) = 0$ if and only if $x = 0$ and use Theorem 8 to prove that $W + M$ is f -closed.

4. Conclusions

In this paper we introduce and study the concept of f -simultaneous approximation of a nonempty subset K of Hausdorff topological vector space X , existence and uniqueness as a generalization to the problem of simultaneous approximation in the sense that if the function f is taken to be the usual norm, the problem is turned out to be precisely the problem of best approximation in the usual sense. Further we obtain some results regarding f -simultaneous approximation in the quotient space.

REFERENCES

- Breckner, W. W. and Brosowski, B. (1971). Ein kriterium zur charakterisierung von sonnen, *Mathematika*, Vol. 13, pp. 181-188.
- Govindarajulu, P. and Pai, D. V. (1980). On properties of sets related to f -projections, *J. Math. Anal. Appl.*, Vol. 73, pp. 457-465.
- Khalil, R. and Abushamla, W. (2003). f -proximality in function spaces, *Dirasat*, Vol. 30, pp. 93-96.
- Moghaddam, M. A. (2010). On f -best approximation in quotient topological vector spaces, *Int. Math. Forum.*, Vol. 12, pp. 587-595.
- Narang, T. D. (1985). On f -best approximation in topological spaces, *Arch. Math.*, Vol. 4, pp. 229-234.
- Narang, T. D. (1986). Approximation relative to an ultra function, *Arch. Math.*, Vol. 4, pp. 181-186.
- Pai, D. V. and Veermani, P. (1982). Applications of fixed point theorems in optimization and best approximation, *Nonlinear Anal. Appl. Marcel Dekker, Int. New York*, Edited by S.P. Singh and J. H. Barry, pp. 393-400.
- Rezapour, Sh. (2003). ϵ -weakly Chebyshev subspaces of Banach spaces, *Anal. Theory Appl.*, Vol. 19, No. 2, pp. 130-135.
- Rezapour, Sh. and Mohebi, H. (2003). ϵ -weakly Chebyshev subspaces and quotient spaces, *Bulletin of the Iranian Math. Soc.*, Vol. 29, No. 2, pp. 27-33.
- Saidi, F., Hussein, D. and Khalil, R. (2002). Best simultaneous approximation in $L^p(I, X)$, *Approx. Theory J.*, Vol. 116, pp. 369-379.
- Singer, I. (1970). *Best approximation in normed linear spaces by elements of linear subspaces*, Springer-Verlag.