




12-2018

Simplifying Coefficients in a Family of Ordinary Differential Equations Related to the Generating Function of the Laguerre Polynomials

Feng Qi

Tianjin Polytechnic University

Follow this and additional works at: <https://digitalcommons.pvamu.edu/aam>

 Part of the [Discrete Mathematics and Combinatorics Commons](#), [Harmonic Analysis and Representation Commons](#), [Number Theory Commons](#), and the [Ordinary Differential Equations and Applied Dynamics Commons](#)

Recommended Citation

Qi, Feng (2018). Simplifying Coefficients in a Family of Ordinary Differential Equations Related to the Generating Function of the Laguerre Polynomials, *Applications and Applied Mathematics: An International Journal (AAM)*, Vol. 13, Iss. 2, Article 9.

Available at: <https://digitalcommons.pvamu.edu/aam/vol13/iss2/9>

This Article is brought to you for free and open access by Digital Commons @PVAMU. It has been accepted for inclusion in *Applications and Applied Mathematics: An International Journal (AAM)* by an authorized editor of Digital Commons @PVAMU. For more information, please contact hvkoshy@pvamu.edu.



Simplifying Coefficients in a Family of Ordinary Differential Equations Related to the Generating Function of the Laguerre Polynomials

Feng Qi

Department of Mathematics
College of Science
Tianjin Polytechnic University
Tianjin 300387, China

College of Mathematics
Inner Mongolia University for Nationalities
Tongliao 028043, China

Institute of Mathematics
Henan Polytechnic University
Jiaozuo 454010, China

qifeng618@gmail.com, qifeng618@hotmail.com

Received: June 10, 2018; Accepted: September 27, 2018

Abstract

In the paper, by virtue of the Faà di Bruno formula, properties of the Bell polynomials of the second kind, and the Lah inversion formula, the author simplifies coefficients in a family of ordinary differential equations related to the generating function of the Laguerre polynomials.

Keywords: Simplifying; coefficient; Laguerre polynomial; generating function; Faà di Bruno formula; ordinary differential equation; Bell polynomial of the second kind; Lah inversion formula

MSC 2010 No.: Primary 34A05; Secondary 05A15, 11A25, 11B73, 11B83, 11C08, 42C10

1. Introduction

In (Kim et al., 2016, Theorem 1), it was established inductively and recursively that the family of differential equations

$$F^{(n)}(t) = F(t) \sum_{i=n}^{2n} \frac{a_{i-n}(n, x)}{(1-t)^i}, \quad n \in \mathbb{N}, \quad (1)$$

has a solution

$$F(t) = \frac{1}{1-t} \exp\left(-\frac{xt}{1-t}\right), \quad (2)$$

where $a_0(n, x) = n!$, $a_n(n, x) = (-x)^n$,

$$a_j(n, x) = (-x)^j \sum_{i_j=0}^{n-j} \sum_{i_{j-1}=0}^{n-i_j-j} \cdots \sum_{i_1=0}^{n-i_j-\cdots-i_2-j} \langle n+j \rangle_{i_j} \\ \times \prod_{k=2}^j \langle n-i_j-\cdots-i_k-[j-(2k-2)] \rangle_{i_{k-1}} (n-i_j-\cdots-i_1-j)!, \quad 2 \leq j \leq n-1, \quad (3)$$

the falling factorial

$$\langle x \rangle_n = \prod_{k=0}^{n-1} (x-k) = \begin{cases} x(x-1)\cdots(x-n+1), & n \geq 1, \\ 1, & n = 0, \end{cases}$$

and the function $F(t)$ in (2) can be used to generate the Laguerre polynomials $L_n(x)$ by

$$F(t) = \frac{1}{1-t} \exp\left(-\frac{xt}{1-t}\right) = \sum_{n=0}^{\infty} L_n(x) t^n.$$

Hereafter, the expression (3) was employed in (Kim et al., 2016, Theorem 2).

It is not difficult to see that

- (1) the expression (3) is too complicated to be remembered, understood, and computed easily;
- (2) the original proof of (Kim et al., 2016, Theorem 1) is long and tedious.

In this paper, we will provide a nice and standard proof for (Kim et al., 2016, Theorem 1) and, more importantly, discover a simple, meaningful, and significant form for $a_i(n, x)$.

2. Main results

Our main results can be stated as the following theorem.

Theorem 2.1.

For $n \geq 0$, the function $F(t)$ defined by (2) and its derivatives satisfy

$$F^{(n)}(t) = \frac{n!}{(1-t)^n} \left[\sum_{k=0}^n (-1)^k \binom{n-1}{k-1} \frac{x^{k-1} x - k(1-t)}{k! (1-t)^k} \right] F(t), \quad (4)$$

and

$$F(t) = \frac{n!(1-t)^n}{x^{n-1}[x-n(1-t)]} \sum_{k=0}^n \frac{(-1)^k}{k!} \binom{n-1}{k-1} (1-t)^k F^{(k)}(t), \tag{5}$$

where $\binom{-1}{-1} = 1$ and $\binom{k}{-1} = 0$ if $k \geq 0$.

Proof:

The famous Faà di Bruno formula reads that

$$\frac{d^n}{dt^n} f \circ h(t) = \sum_{k=0}^n f^{(k)}(h(t)) B_{n,k}(h'(t), h''(t), \dots, h^{(n-k+1)}(t)), \tag{6}$$

where $n \geq 0$ and the Bell polynomials of the second kind $B_{n,k}(x_1, x_2, \dots, x_{n-k+1})$ for $n \geq k \geq 0$ are defined (Comtet, 1974, p. 134, Theorem A) and (Comtet, 1974, p. 139, Theorem C) by

$$B_{n,k}(x_1, x_2, \dots, x_{n-k+1}) = \sum_{\substack{1 \leq i \leq n-k+1 \\ \ell_i \in \{0\} \cup \mathbb{N} \\ \sum_{i=1}^{n-k+1} i \ell_i = n \\ \sum_{i=1}^{n-k+1} \ell_i = k}} \frac{n!}{\prod_{i=1}^{n-k+1} \ell_i!} \prod_{i=1}^{n-k+1} \left(\frac{x_i}{i!}\right)^{\ell_i}.$$

The function $F(t)$ in (2) can be rearranged as

$$F(t) = \frac{e^x}{x} \frac{x}{1-t} \exp\left(-\frac{x}{1-t}\right).$$

Applying $u = h(t) = \frac{x}{1-t}$ and $f(u) = \frac{u}{e^u}$ to (6) gives

$$\begin{aligned} \frac{x}{e^x} F^{(n)}(t) &= \sum_{k=0}^n \frac{d^k}{du^k} \left(\frac{u}{e^u}\right) B_{n,k}\left(\frac{1!x}{(1-t)^2}, \frac{2!x}{(1-t)^3}, \dots, \frac{(n-k+1)!x}{(1-t)^{n-k+2}}\right) \\ &= \sum_{k=0}^n \frac{d^k}{du^k} \left(\frac{u}{e^u}\right) x^k \left(\frac{1}{1-t}\right)^{n+k} B_{n,k}(1!, 2!, \dots, (n-k+1)!) \\ &= \sum_{k=0}^n (-1)^k \frac{u-k}{e^u} x^k \left(\frac{1}{1-t}\right)^{n+k} \frac{n!}{k!} \binom{n-1}{k-1} \\ &= \left(\frac{1}{1-t}\right)^n \exp\left(-\frac{x}{1-t}\right) \sum_{k=0}^n (-1)^k \binom{n-1}{k-1} \frac{n!}{k!} x^k \left(\frac{x}{1-t} - k\right) \left(\frac{1}{1-t}\right)^k, \end{aligned}$$

where we used the identities

$$B_{n,k}(abx_1, ab^2x_2, \dots, ab^{n-k+1}x_{n-k+1}) = a^k b^n B_{n,k}(x_1, x_2, \dots, x_{n-k+1}),$$

and

$$B_{n,k}(1!, 2!, \dots, (n-k+1)!) = \frac{n!}{k!} \binom{n-1}{k-1},$$

in (Comtet, 1974, p. 135) and (Qi, 2016, Remark 3.5). The formula (4) is thus proved.

The Lah inversion theorem in (Aigner, 1979, p. 96, Corollary 3.38 (iii)) and (Aigner, 2007, pp. 60–61, Exercise 2.9) reads that

$$(-1)^n a_n = \sum_{k=0}^n \frac{n!}{k!} \binom{n-1}{k-1} b_k \iff (-1)^n b_n = \sum_{k=0}^n \frac{n!}{k!} \binom{n-1}{k-1} a_k. \tag{7}$$

Combining (7) with (4) arrives at

$$\frac{x^{n-1}[x - n(1-t)]}{(1-t)^n} = \sum_{k=0}^n (-1)^k \binom{n-1}{k-1} \frac{n!}{k!} \frac{(1-t)^k}{F(t)} F^{(k)}(t),$$

which can be rewritten as (5). The proof of Theorem 2.1 is complete. ■

3. Remarks

Finally, we list several remarks on our main results and closely related things.

Remark.

The equations (4) and (5) can be regarded as inversion formulas each other.

Remark.

The equation (5) is a new one which cannot be obtained inductively and recursively, as was in Kim et al. (2016).

Remark.

The equation (4) can be rewritten as

$$\begin{aligned} F^{(n)}(t) &= \frac{F(t)}{(1-t)^n} \sum_{k=0}^n (-1)^k \binom{n-1}{k-1} \frac{n!}{k!} \left[\frac{x^k}{(1-t)^k} - \frac{kx^{k-1}}{(1-t)^{k-1}} \right] \\ &= \frac{F(t)}{(1-t)^n} \left[\sum_{k=0}^n (-1)^k \binom{n-1}{k-1} \frac{n!}{k!} \frac{x^k}{(1-t)^k} - \sum_{k=1}^n (-1)^k \binom{n-1}{k-1} \frac{n!}{k!} \frac{kx^{k-1}}{(1-t)^{k-1}} \right] \\ &= \frac{F(t)}{(1-t)^n} \left[\sum_{k=0}^n (-1)^k \binom{n-1}{k-1} \frac{n!}{k!} \frac{x^k}{(1-t)^k} + \sum_{k=0}^{n-1} (-1)^k \binom{n}{k+1} \frac{(n-1)!(k+1)x^k}{k! (1-t)^k} \right] \quad (8) \\ &= \frac{F(t)}{(1-t)^n} \left[\frac{(-1)^n x^n}{(1-t)^n} + \sum_{k=0}^{n-1} (-1)^k \binom{n}{k} \frac{n!}{k!} \frac{x^k}{(1-t)^k} \right] \\ &= \frac{F(t)}{(1-t)^n} \sum_{k=0}^n (-1)^k \binom{n}{k} \frac{n!}{k!} \frac{x^k}{(1-t)^k}. \end{aligned}$$

The equation (1) can be reformulated as

$$F^{(n)}(t) = F(t) \sum_{i=0}^n \frac{a_i(n, x)}{(1-t)^{n+i}}, \quad n \in \mathbb{N}. \quad (9)$$

Remark.

The motivations in the papers Qi (2018a,b,c), Qi and Guo (2018, 2017), Qi et al. (2019a, 2018a, 2019b, 2018b,c), Qi and Zhao (2018), Zhao et al. (2018) are same as the one in this paper.

Remark.

This paper is a slightly modified version of the preprint Qi (2017).

4. Conclusions

Comparing two equalities (8) and (9) reveals that

$$a_k(n, x) = (-1)^k \binom{n}{k} \frac{n!}{k!} x^k,$$

for $n \geq k \geq 0$. This form for $a_k(n, x)$ is apparently simpler, more meaningful, and more significant than the one (3) obtained in (Kim et al., 2016, Theorem 1).

Acknowledgement:

The author thanks the anonymous referees for their careful comments on and valuable suggestions to the original version of this paper.

REFERENCES

- Aigner, M. (1979). *Combinatorial Theory*, Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], 234, Springer-Verlag, Berlin-New York.
- Aigner, M. (2007). *Discrete Mathematics*, translated from the 2004 German original by David Kramer, American Mathematical Society, Providence, RI.
- Comtet, L. (1974). *Advanced Combinatorics: The Art of Finite and Infinite Expansions*, Revised and Enlarged Edition, D. Reidel Publishing Co., Dordrecht and Boston. Available online at <https://doi.org/10.1007/978-94-010-2196-8>.
- Kim, T., Kim, D.S., Hwang, K.-W., and Seo, J.J. (2016). Some identities of Laguerre polynomials arising from differential equations, *Advances in Difference Equations*, **2016**, 2016:159, 9 pages. Available online at <https://doi.org/10.1186/s13662-016-0896-1>.
- Qi, F. (2016). Diagonal recurrence relations for the Stirling numbers of the first kind, *Contributions to Discrete Mathematics*, Vol. 11, No. 1, pp. 22–30. Available online at <http://hdl.handle.net/10515/sy5wh2dx6> and <https://doi.org/10515/sy5wh2dx6>.
- Qi, F. (2017). Simplifying coefficients in a family of ordinary differential equations related to the generating function of the Laguerre polynomials, *ResearchGate Preprint*, available online at <https://doi.org/10.13140/RG.2.2.13602.53448>.
- Qi, F. (2018a). A simple form for coefficients in a family of nonlinear ordinary differential equations, *Advances and Applications in Mathematical Sciences*, Vol. 17, No. 8, pp. 555–561.
- Qi, F. (2018b). Notes on several families of differential equations related to the generating function for the Bernoulli numbers of the second kind, *Turkish Journal of Analysis and Number Theory*, Vol. 6, No. 2, pp. 40–42. Available online at <https://doi.org/10.12691/tjant-6-2-1>.
- Qi, F. (2018c). Simple forms for coefficients in two families of ordinary differential equations, *Global Journal of Mathematical Analysis*, Vol. 6, No. 1, pp. 7–9. Available online at <https://doi.org/10.14419/gjma.v6i1.9778>.
- Qi, F. and Guo, B.-N. (2017). Explicit formulas and recurrence relations for higher order Eulerian

- polynomials, *Indagationes Mathematicae*, Vol. 28, No. 4, pp. 884–891. Available online at <https://doi.org/10.1016/j.indag.2017.06.010>.
- Qi, F. and Guo, B.-N. (2018). A diagonal recurrence relation for the Stirling numbers of the first kind, *Applicable Analysis and Discrete Mathematics*, Vol. 12, No. 1, pp. 153–165. Available online at <https://doi.org/10.2298/AADM170405004Q>.
- Qi, F., Lim, D. and Guo, B.-N. (2019a). Explicit formulas and identities for the Bell polynomials and a sequence of polynomials applied to differential equations, *Revista de la Real Academia de Ciencias Exactas, Físicas y Naturales–Serie A: Matemáticas*, Vol. 113, in press. Available online at <https://doi.org/10.1007/s13398-017-0427-2>.
- Qi, F., Lim, D. and Guo, B.-N. (2018a). Some identities related to Eulerian polynomials and involving the Stirling numbers, *Applicable Analysis and Discrete Mathematics*, Vol. 12, No. 2, in press. Available online at <https://doi.org/10.2298/AADM171008014Q>.
- Qi, F., Niu, D.-W. and Guo, B.-N. (2019b). Some identities for a sequence of unnamed polynomials connected with the Bell polynomials, *Revista de la Real Academia de Ciencias Exactas, Físicas y Naturales–Serie A: Matemáticas*, Vol. 113, in press. Available online at <https://doi.org/10.1007/s13398-018-0494-z>.
- Qi, F., Wang, J.-L. and Guo, B.-N. (2018b). Notes on a family of inhomogeneous linear ordinary differential equations, *Advances and Applications in Mathematical Sciences*, Vol. 17, No. 4, pp. 361–368. Available online at <https://doi.org/10.1007/s13398-018-0494-z>.
- Qi, F., Wang, J.-L. and Guo, B.-N. (2018c). Simplifying differential equations concerning degenerate Bernoulli and Euler numbers, *Transactions of A. Razmadze Mathematical Institute*, Vol. 172, No. 1, pp. 90–94. Available online at <https://doi.org/10.1016/j.trmi.2017.08.001>.
- Qi, F. and Zhao, J.-L. (2018). Some properties of the Bernoulli numbers of the second kind and their generating function, *Bulletin of the Korean Mathematical Society*, Vol. 55, in press. Available online at <https://doi.org/10.4134/BKMS.b180039>.
- Zhao, J.-L. and Qi, F. (2018). Derivative polynomials of a function related to the Apostol–Euler and Frobenius–Euler numbers, *Journal of Nonlinear Sciences and Applications*, Vol. 10, No. 4, pp. 1345–1349. Available online at <https://doi.org/10.22436/jnsa.010.04.06>.