



12-2018

A de Casteljaou Algorithm for Bernstein type Polynomials based on $(p; q)$ -integers

Khalid Khan
SC & SS, J.N.U.

D. K. Lobiyal
SC & SS, J.N.U.

Adem Kilicman
University of Putra Malaysia

Follow this and additional works at: <https://digitalcommons.pvamu.edu/aam>



Part of the [Numerical Analysis and Computation Commons](#)

Recommended Citation

Khan, Khalid; Lobiyal, D. K.; and Kilicman, Adem (2018). A de Casteljaou Algorithm for Bernstein type Polynomials based on $(p; q)$ -integers, *Applications and Applied Mathematics: An International Journal (AAM)*, Vol. 13, Iss. 2, Article 25.

Available at: <https://digitalcommons.pvamu.edu/aam/vol13/iss2/25>

This Article is brought to you for free and open access by Digital Commons @PVAMU. It has been accepted for inclusion in *Applications and Applied Mathematics: An International Journal (AAM)* by an authorized editor of Digital Commons @PVAMU. For more information, please contact hvkoshy@pvamu.edu.



A de Casteljaou Algorithm for Bernstein type Polynomials based on (p, q) -integers

¹Khalid Khan, ²D. K. Lobiyal and ³Adem Kilicman

^{1,2} School of Computer and System Science
 SC & SS, J.N.U.
 New Delhi, India

¹khalidga1517@gmail.com; ²dklobiyal@gmail.com;

³Department of Mathematics
 Faculty of Science
 University of Putra Malaysia
 Malaysia

akilicman@putra.upm.edu.my

Received: March 9, 2017; Accepted: October 12, 2017

Abstract

In this paper, a de Casteljaou algorithm to compute (p, q) -Bernstein Bézier curves based on (p, q) -integers are introduced. The nature of degree elevation and degree reduction for (p, q) -Bézier Bernstein functions are studied. The new curves have some properties similar to q -Bézier curves. Moreover, we construct the corresponding tensor product surfaces over the rectangular domain $(u, v) \in [0, 1] \times [0, 1]$ depending on four parameters. De Casteljaou algorithm and degree evaluation properties of the surfaces for these generalization over the rectangular domain are investigated. Furthermore, some fundamental properties for (p, q) -Bernstein Bézier curves are discussed. We get q -Bézier curves and surfaces for $(u, v) \in [0, 1] \times [0, 1]$ when we set the parameter $p_1 = p_2 = 1$. In comparison to q -Bézier curves based on q -Bernstein polynomials, this generalization gives us more flexibility in controlling the shapes of curves.

Keywords: (p, q) -integers; (p, q) -Bernstein polynomials; Degree elevation; Degree reduction; de Casteljaou algorithm; Tensor product; (p, q) -Bézier curve; (p, q) -Bézier surface; Shape preserving.

2010 MSC No.: 65D17, 41A10, 41A25, 41A36

1. Introduction

Recently, Mursaleen et al. (2015, 2016) applied the concept of (p, q) -calculus (which is considered as an extension of quantum-calculus) in Approximation theory and introduced the (p, q) -analogue of Bernstein operators based on (p, q) -integers. Apart from various generalisation of Bernstein operators, another paper by Mursaleen et al. (2016) which catches attention of researchers working in Approximation theory is Bleimann-Butzer-Hahn operators defined by (p, q) -integers. After this initiation, many researchers started working in this area. For similar works based on (p, q) -integers, one can refer [Acar (2016); Acar et al. (2016); Cai and Zhou (2016); Kadak (2016); Kadak (2017); Khan and Lobiyal (2017); Mursaleen et al. (2015); Mursaleen et al. (2016); Mursaleen et al. (2015); Mursaleen et al. (2016); Mishra and Pandey (2016); Mishra and Pandey (2017); Wafi and Rao (2016); Wafi and Rao (2017)].

Motivated by the above mentioned work, the idea of (p, q) -calculus (post quantum calculus) and its importance, we construct (p, q) -Bézier curves and surfaces based on (p, q) -integers which is further generalization of q -Bézier curves and surfaces.

Some of the advantages of using the extra parameter p has been discussed in [Khan and Lobiyal (2017); Mursaleen et al. (2016)].

It was S.N. Bernstein [Bernstein (1912)], who first introduced his famous operators $B_n : C[0, 1] \rightarrow C[0, 1]$ defined for any $n \in \mathbb{N}$ and for any function $f \in C[0, 1]$

$$B_n(f; x) = \sum_{k=0}^n \binom{n}{k} x^k (1-x)^{n-k} f\left(\frac{k}{n}\right), \quad x \in [0, 1], \quad (1)$$

and named it Bernstein polynomials to provide a constructive proof of the Weierstrass theorem [Korovkin (1960)]. Later, it was found that Bernstein polynomials possess many remarkable properties and has various applications in areas such as approximation theory, numerical analysis, computer-aided geometric design, and solutions of differential equations due to its fine properties of approximation [Oruk and Phillips (2003)].

In Computer Aided Geometric Design (CAGD), Bernstein polynomials and its variants are used in order to preserve the shape of the curves or surfaces. One of the most important curve in CAGD [Sederberg (2014); Farouki and Rajan (1988)] is the classical Bézier curve [Bézier (1972)] constructed with the help of Bernstein basis functions.

In recent years, generalization of the Bézier curve with shape parameters has received continuous attention. Several authors were concerned with the problem of changing the shape of curves and surfaces, while keeping the control polygon unchanged and thus, they generalized the Bézier curves in [Farouki and Rajan (1988); Hana et al. (2014); Oruk and Phillips (2003); Rababah and Manna (2011)].

Lupaş (1987) introduced the first q -analogue of Bernstein operators (rational) as follows:

$$L_{n,q}(f; x) = \sum_{k=0}^n \frac{f\left(\frac{[k]_q}{[n]_q}\right) \begin{bmatrix} n \\ k \end{bmatrix}_q q^{\frac{k(k-1)}{2}} x^k (1-x)^{n-k}}{\prod_{j=1}^n \{(1-x) + q^{j-1}x\}}, \quad (2)$$

and investigated its approximating and shape-preserving properties.

In 1996, Phillips (1997) proposed another q -variant of the classical Bernstein operators, the so-called Phillips q -Bernstein operators which attracted lots of investigations.

$$B_{n,q}(f; x) = \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q x^k \prod_{s=0}^{n-k-1} (1 - q^s x) f\left(\frac{[k]_q}{[n]_q}\right), \quad x \in [0, 1], \quad (3)$$

where $B_{n,q} : C[0, 1] \rightarrow C[0, 1]$ defined for any $n \in \mathbb{N}$ and any function $f \in C[0, 1]$.

The q -variants of Bernstein polynomials provide one shape parameter for constructing free-form curves and surfaces, Phillips q -Bernstein operator was applied well in this area.

In 2003, Oruk and Phillips [Oruk and Phillips (2003)] used the basis functions of Phillips q -Bernstein operator for construction of q -Bézier curves, which they call Phillips q -Bézier curves, and studied the properties of degree reduction and elevation.

Thus with the development of (p, q) -analogue of Bernstein operators and its variants, one natural question arises, how it can be used in order to preserve the shape of the curves or surfaces. In this way, it opens a new research direction which requires further investigations.

Recently, (p, q) -analogue of Lupaş Bernstein operators are defined as follows in [Khan and Lobiyal (2017)]:

For any $p > 0$ and $q > 0$, the linear operators $L_{p,q}^n : C[0, 1] \rightarrow C[0, 1]$

$$L_{p,q}^n(f; x) = \sum_{k=0}^n \frac{f\left(\frac{p^{n-k} [k]_{p,q}}{[n]_{p,q}}\right) \begin{bmatrix} n \\ k \end{bmatrix}_{p,q} p^{\frac{(n-k)(n-k-1)}{2}} q^{\frac{k(k-1)}{2}} x^k (1-x)^{n-k}}{\prod_{j=1}^n \{p^{j-1}(1-x) + q^{j-1}x\}}, \quad (4)$$

is (p, q) -analogue of Lupaş Bernstein operators.

Again when $p = 1$, Lupaş (p, q) -Bernstein operators turns out to be Lupaş q -Bernstein operators as given in [Mahmudov and Sabancıgil (2010)].

When $p = q = 1$, Lupaş (p, q) -Bernstein operators turns out to be classical Bernstein operators.

Let us recall certain notations of (p, q) -calculus .

For any $p > 0$ and $q > 0$, the (p, q) integers $[n]_{p,q}$ are defined by

$$[n]_{p,q} = p^{n-1} + p^{n-2}q + p^{n-3}q^2 + \dots + pq^{n-2} + q^{n-1} = \begin{cases} \frac{p^n - q^n}{p - q}, & \text{when } p \neq q \neq 1, \\ n p^{n-1}, & \text{when } p = q \neq 1, \\ [n]_q, & \text{when } p = 1, \\ n, & \text{when } p = q = 1, \end{cases}$$

where $[n]_q$ denotes the q -integers and $n = 0, 1, 2, \dots$.

It is obvious that $[n]_{p,q} = p^{n-1} [n]_{\frac{q}{p}}$.

The formula for (p, q) -binomial expansion is as follows:

$$(ax + by)_{p,q}^n := \sum_{k=0}^n p^{\frac{(n-k)(n-k-1)}{2}} q^{\frac{k(k-1)}{2}} \begin{bmatrix} n \\ k \end{bmatrix}_{p,q} a^{n-k} b^k x^{n-k} y^k.$$

For $a = b = 1$,

$$(x + y)_{p,q}^n := \sum_{k=0}^n p^{\frac{(n-k)(n-k-1)}{2}} q^{\frac{k(k-1)}{2}} \begin{bmatrix} n \\ k \end{bmatrix}_{p,q} x^{n-k} y^k.$$

On using,

$$(x + y)_{p,q}^n = (x + y)(px + qy)(p^2x + q^2y) \cdots (p^{n-1}x + q^{n-1}y),$$

one can write,

$$(x)_{p,q}^n = (x)(px)(p^2x) \cdots (p^{n-1}x) = p^{\frac{n(n-1)}{2}} x^n,$$

which implies

$$(1)_{p,q}^n = (1)(p)(p^2) \cdots (p^{n-1}) = p^{\frac{n(n-1)}{2}}.$$

The (p, q) -analogue of Euler’s identity is as follows:

$$\begin{aligned} (1-x)_{p,q}^n &= \prod_{s=0}^{n-1} (p^s - q^s x) = (1-x)(p - qx)(p^2 - q^2 x) \dots (p^{n-1} - q^{n-1} x) \\ &= \sum_{k=0}^n (-1)^k p^{\frac{(n-k)(n-k-1)}{2}} q^{\frac{k(k-1)}{2}} \begin{bmatrix} n \\ k \end{bmatrix}_{p,q} x^k. \end{aligned}$$

From (p, q) -binomial expansion, or by using induction on n , one can easily verify that

$$\sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_{p,q} p^{\frac{k(k-1)}{2}} x^k \prod_{s=0}^{n-k-1} (p^s - q^s x) = p^{\frac{n(n-1)}{2}}, \quad x \in [0, 1]. \tag{5}$$

Again by some simple calculations and using the property of (p, q) -integers, one can find (p, q) -analogue of Pascal’s relation as follows:

$$\begin{bmatrix} n \\ k \end{bmatrix}_{p,q} = q^{n-k} \begin{bmatrix} n-1 \\ k-1 \end{bmatrix}_{p,q} + p^k \begin{bmatrix} n-1 \\ k \end{bmatrix}_{p,q} \tag{6}$$

$$\begin{bmatrix} n \\ k \end{bmatrix}_{p,q} = p^{n-k} \begin{bmatrix} n-1 \\ k-1 \end{bmatrix}_{p,q} + q^k \begin{bmatrix} n-1 \\ k \end{bmatrix}_{p,q}. \tag{7}$$

Details on (p, q) -calculus can be found in [Hounkonnou et al. (2013); Jagannathan and Srinivasa (2005)].

The (p, q) -Bernstein Operators introduced by Mursaleen et al. (2015, 2016) for $0 < q < p \leq 1$ and $x \in [0, 1]$ are as follows:

$$B_{n,p,q}(f; x) = \frac{1}{p^{\frac{n(n-1)}{2}}} \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_{p,q} p^{\frac{k(k-1)}{2}} x^k \prod_{s=0}^{n-k-1} (p^s - q^s x) f\left(\frac{[k]_{p,q}}{p^{k-n} [n]_{p,q}}\right). \tag{8}$$

Note when $p = 1$, (p, q) -Bernstein Operators given by (8) turns out to be Phillips q -Bernstein Operators.

The operators given by (4) and (8) are respective generalization of q -Lupaş and q -Phillips operators.

For other works relevant to this field, one can see [Agarwal et al. (2013); Aral et al. (2013); Mishra and Deepmala (2013); Mursaleen and Khan (2013); Mursaleen and Nasiruzzaman (2017); Victor and Pokman (2002)].

We apply post-quantum calculus to introduce the (p, q) -Bézier curves and surfaces based on (p, q) -integers constructed with the help of Bernstein basis of (8) which is further generalization of q -Bézier curves and surfaces, for example, [Lupaş (1987); Hana et al. (2014); Oruk and Phillips (2003)].

The outline of this paper are as follows: Section 2 introduces a (p, q) -analogue of the Phillips q -Bernstein functions $B_{p,q}^{k,n}$ and its Properties. Section 3 introduces degree elevation and degree reduction properties for (p, q) -analogue of the Bernstein functions. Section 4 introduces a de Casteljau type algorithm for $B_{p,q}^{k,n}$. In Section 5, we define a tensor product patch based on algorithm 1 and its geometric properties as well as a degree elevation technique are investigated. Furthermore, tensor product of (p, q) -Bézier surfaces on $[0, 1] \times [0, 1]$ for (p, q) -Bernstein polynomials are introduced and its properties that is inherited from the univariate case are being discussed. In section 6, we discuss Shape control of (p, q) -Bernstein curves.

2. (p, q) -Bernstein functions

For any $p > 0$ and $q > 0$ and $t \in [0, 1]$, the (p, q) -Bernstein functions are defined as follows:

$$B_{p,q}^{k,n}(t) = \frac{1}{p^{\frac{n(n-1)}{2}}} \begin{bmatrix} n \\ k \end{bmatrix}_{p,q} p^{\frac{k(k-1)}{2}} t^k (1-t)_{p,q}^{n-k}, \quad (9)$$

where

$$(1-t)_{p,q}^{n-k} = \prod_{s=0}^{n-k-1} (p^s - q^s t).$$

Note: The (p, q) -analogue of Bernstein operators introduced by Mursaleen et al. generate positive linear operators only if $0 < q < p \leq 1$.

2.1. Properties of the (p, q) -analogue of the Bernstein functions

Theorem 2.1.

The (p, q) -analogue of the Bernstein functions possess the following properties:

(1). Non-negativity: $B_{p,q}^{k,n}(t) \geq 0$ $k = 0, 1, \dots, n$, $t \in [0, 1]$.

(2). Partition of unity:

$$\sum_{k=0}^n B_{p,q}^{k,n}(t) = 1, \quad \text{for every } t \in [0, 1].$$




Figure 1. Cubic Bezier blending functions

(3). Both sided end-point property:

$$B_{p,q}^{k,n}(0) = \begin{cases} 1, & \text{if } k = 0, \\ 0, & k \neq 0, \end{cases}$$

$$B_{p,q}^{k,n}(1) = \begin{cases} 1, & \text{if } k = n, \\ 0, & k \neq n, \end{cases}$$

when $p = 1$, then, both side end point interpolation property holds.

(4). Reducibility: when $p = 1$, formula (2.1) reduces to the q -Bernstein bases.

Proof:

All these property can be deduced easily from equation (9).

Figure 2 shows the (p, q) -analogues of the Bernstein basis functions of degree 3 with $q = 0.5$, $p = 0.8$. Here we can observe that sum of blending fuctions is always unity and also end point interpolation property holds, when we put $p = 1$, it turns out to be q -Bernstein basis which is shown in Figure 1.

Apart from the basic properties above, the (p, q) -analogue of the Bernstein functions also satisfy some recurrence relations, as for the classical Bernstein basis.




Figure 2. Cubic Bezier blending functions

3. Degree elevation and reduction for (p, q) -Bernstein functions

Technique of degree elevation has been used to increase the flexibility of a given curve. A degree elevation algorithm calculates a new set of control points by choosing a convex combination of the old set of control points which retains the old end points. For this purpose, the identities (12),(13) and Theorem (3.2) are useful.

Theorem 3.1.

Each (p, q) -Bernstein functions of degree n is a linear combination of two (p, q) -Bernstein functions of degree $n - 1$:

$$B_{p,q}^{k,n}(t) = q^{n-k} p^{k-1} t B_{p,q}^{k-1,n-1}(t) + (p^{n-1} - p^k q^{n-k-1} t) B_{p,q}^{k,n-1}(t), \quad (10)$$

$$B_{p,q}^{k,n}(t) = p^{n-1} t B_{p,q}^{k-1,n-1}(t) + (q^k p^{n-k-1} - q^{n-1} t) B_{p,q}^{k,n-1}(t). \quad (11)$$

Proof: On using Pascal’s type relation based on (p, q) -integers i.e (6), we get

$$\begin{aligned}
 B_{p,q}^{k,n}(t) &= \frac{1}{p^{\frac{n(n-1)}{2}}} \left(q^{n-k} \begin{bmatrix} n-1 \\ k-1 \end{bmatrix}_{p,q} + p^k \begin{bmatrix} n-1 \\ k \end{bmatrix}_{p,q} \right) p^{\frac{k(k-1)}{2}} t^k (1-t)_{p,q}^{n-k} \\
 &= \frac{1}{p^{\frac{n(n-1)}{2}}} q^{n-k} \begin{bmatrix} n-1 \\ k-1 \end{bmatrix}_{p,q} p^{\frac{k(k-1)}{2}} t^k (1-t)_{p,q}^{n-k} \\
 &\quad + \frac{1}{p^{\frac{n(n-1)}{2}}} p^k \begin{bmatrix} n-1 \\ k \end{bmatrix}_{p,q} p^{\frac{k(k-1)}{2}} t^k (1-t)_{p,q}^{n-k} \\
 &= \frac{1}{p^{\frac{n(n-1)}{2}}} q^{n-k} p^{k-1} t \begin{bmatrix} n-1 \\ k-1 \end{bmatrix}_{p,q} p^{\frac{(k-1)(k-2)}{2}} t^{k-1} (1-t)_{p,q}^{n-k} \\
 &\quad + \frac{1}{p^{\frac{n(n-1)}{2}}} p^k (p^{n-k-1} - q^{n-k-1}t) \begin{bmatrix} n-1 \\ k \end{bmatrix}_{p,q} p^{\frac{k(k-1)}{2}} t^k (1-t)_{p,q}^{n-k-1} \\
 &= q^{n-k} p^{k-1} t B_{p,q}^{k-1,n-1}(t) + (p^{n-1} - p^k q^{n-k-1}t) B_{p,q}^{k,n-1}(t).
 \end{aligned}$$

Thus,

$$B_{p,q}^{k,n}(t) = q^{n-k} p^{k-1} t B_{p,q}^{k-1,n-1}(t) + (p^{n-1} - p^k q^{n-k-1}t) B_{p,q}^{k,n-1}(t).$$

Similarly, if we use (7), we have

$$B_{p,q}^{k,n}(t) = p^{n-1} t B_{p,q}^{k-1,n-1}(t) + (q^k p^{n-k-1} - q^{n-1}t) B_{p,q}^{k,n-1}(t).$$

3.1. Degree elevation

$$q^{n-k} p^k t B_{p,q}^{k,n}(t) = \left(1 - \frac{p^{k+1} [n-k]_{p,q}}{[n+1]_{p,q}} \right) B_{p,q}^{k+1,n+1}(t), \tag{12}$$

$$(p^n - p^k q^{n-k} t) B_{p,q}^{k,n}(t) = \left(\frac{p^k [n+1-k]_{p,q}}{[n+1]_{p,q}} \right) B_{p,q}^{k,n+1}(t). \tag{13}$$

Proof:

$$\begin{aligned}
 q^{n-k} p^k t B_{p,q}^{k,n}(t) &= \frac{1}{p^{\frac{n(n-1)}{2}}} q^{n-k} p^k t \left(\begin{bmatrix} n \\ k \end{bmatrix}_{p,q} p^{\frac{k(k-1)}{2}} t^k (1-t)_{p,q}^{n-k} \right) \\
 &= q^{n-k} \frac{[k+1]_{p,q}}{[n+1]_{p,q}} \frac{1}{p^{\frac{n(n-1)}{2}}} \left(\begin{bmatrix} n+1 \\ k+1 \end{bmatrix}_{p,q} p^{\frac{(k+1)k}{2}} t^{k+1} (1-t)_{p,q}^{n-k} \right) \\
 &= q^{n-k} \frac{[k+1]_{p,q}}{[n+1]_{p,q}} B_{p,q}^{k+1,n+1}(t).
 \end{aligned}$$

By some simple calculation, we have

$$q^{n-k} \frac{[k+1]_{p,q}}{[n+1]_{p,q}} = 1 - \frac{p^{k+1} [n-k]_{p,q}}{[n+1]_{p,q}},$$

using this result, we get

$$q^{n-k} p^k t B_{p,q}^{k,n}(t) = \left(1 - \frac{p^{k+1} [n-k]_{p,q}}{[n+1]_{p,q}}\right) B_{p,q}^{k+1,n+1}(t).$$

Similarly on considering,

$$\begin{aligned} (p^n - p^k q^{n-k} t) B_{p,q}^{k,n}(t) &= (p^n - p^k q^{n-k} t) \left(\frac{1}{p^{\frac{n(n-1)}{2}}} \begin{bmatrix} n \\ k \end{bmatrix}_{p,q} p^{\frac{k(k-1)}{2}} t^k (1-t)_{p,q}^{n-k} \right) \\ &= \frac{(p^n - p^k q^{n-k} t)}{(p^{n-k} - q^{n-k} t)} \frac{1}{p^{\frac{n(n-1)}{2}}} \frac{[n+1-k]_{p,q}}{[n+1]_{p,q}} \\ &\times \left(\begin{bmatrix} n+1 \\ k \end{bmatrix}_{p,q} p^{\frac{k(k-1)}{2}} t^k (1-t)_{p,q}^{n+1-k} \right), \end{aligned}$$

finally we get

$$(p^n - p^k q^{n-k} t) B_{p,q}^{k,n}(t) = \left(\frac{p^k [n+1-k]_{p,q}}{[n+1]_{p,q}} \right) B_{p,q}^{k,n+1}(t).$$

Theorem 3.2.

Each (p, q) -Bernstein function of degree n is a linear combination of two (p, q) -Bernstein functions of degree $n+1$.

$$B_{p,q}^{k,n}(t) = \left(\frac{p^{k-n} [n+1-k]_{p,q}}{[n+1]_{p,q}} \right) B_{p,q}^{k,n+1}(t) + p^{-n} \left(1 - \frac{p^{k+1} [n-k]_{p,q}}{[n+1]_{p,q}} \right) B_{p,q}^{k+1,n+1}(t). \quad (14)$$

Proof:

From equation (12) and (13), we have

$$B_{p,q}^{k,n}(t) = \left(\frac{p^{k-n} [n+1-k]_{p,q}}{[n+1]_{p,q}} \right) B_{p,q}^{k,n+1}(t) + p^{-n} \left(1 - \frac{p^{k+1} [n-k]_{p,q}}{[n+1]_{p,q}} \right) B_{p,q}^{k+1,n+1}(t).$$

For $n=3$, the blending functions are given by:

$$B_{p,q}^{0,3} = \frac{1}{p^3} \begin{bmatrix} 3 \\ 0 \end{bmatrix}_{p,q} (1-t)(p-qt)(p^2-q^2t),$$

$$B_{p,q}^{1,3} = \frac{1}{p^3} \begin{bmatrix} 3 \\ 1 \end{bmatrix}_{p,q} t(1-t)(p-qt),$$

$$B_{p,q}^{2,3} = \frac{1}{p^3} \begin{bmatrix} 3 \\ 2 \end{bmatrix}_{p,q} pt^2(1-t),$$

$$B_{p,q}^{3,3} = \begin{bmatrix} 3 \\ 3 \end{bmatrix}_{p,q} t^3.$$

We observed that the both sided end point interpolation property and partition of unity property always holds in case of (p, q) -Bernstein functions.

The de Casteljau algorithm describes how to subdivide a Bézier curve, when a Bézier curve is repeatedly subdivided, the collection of control polygons converge to the curve. Thus, the way of computing a Bézier curve is to simply subdivide it an appropriate number of times and compute the control polygons.

4. (p, q) -Bernstein Bézier curves:

Let us define the (p, q) -Bézier curves of degree n using the (p, q) -analogues of the Bernstein functions as follows:

$$\mathbf{P}(t; p, q) = \sum_{i=0}^n \mathbf{P}_i B_{p,q}^{i,n}(t), \quad (15)$$

where $P_i \in R^3$ ($i = 0, 1, \dots, n$) and $p > q > 0$. P_i are control points. Joining up adjacent points P_i , $i = 0, 1, 2, \dots, n$ to obtain a polygon which is called the control polygon of (p, q) -Bézier curves.

4.1. Some basic properties of (p, q) -Bézier curves.

Theorem 4.1.

From the definition, we can derive some basic properties of (p, q) -Bézier curves:

1. (p, q) -Bézier curves have geometric and affine invariance.
2. (p, q) -Bézier curves lie inside the convex hull of its control polygon.
3. The end-point interpolation property: $\mathbf{P}(0; p, q) = \mathbf{P}_0$, $\mathbf{P}(1; p, q) = \mathbf{P}_n$.
4. Reducibility: when $p = 1$, formula 15 gives the q -Bézier curves.

Proof:

These properties of (p, q) -Bézier curves can be easily deduced from corresponding properties of the (p, q) -analogue of the Bernstein functions.

4.2. Degree elevation for (p, q) -Bézier curves

(p, q) -Bézier curves have a degree elevation algorithm that is similar to that possessed by the classical Bézier curves. Using the technique of degree elevation, we can increase the flexibility of a given curve.

$$\mathbf{P}(t; p, q) = \sum_{k=0}^n \mathbf{P}_k B_{p,q}^{k,n}(t),$$

$$\mathbf{P}(t; p, q) = \sum_{k=0}^{n+1} \mathbf{P}'_k B_{p,q}^{k+1,n}(t),$$

where

$$\mathbf{P}_k^* = p^n \mathbf{P}'_k = \left(1 - \frac{p^k [n+1-k]_{p,q}}{[n+1]_{p,q}}\right) \mathbf{P}_{k-1} + p^k \left(\frac{[n+1-k]_{p,q}}{[n+1]_{p,q}}\right) \mathbf{P}_k. \quad (16)$$

The statement above can be derived using the identities (12) and (13). Consider

$$p^n \mathbf{P}(t; p, q) = (p^n - p^k q^{n-k} t) \mathbf{P}(t; p, q) + p^k q^{n-k} t \mathbf{P}(t; p, q).$$

We obtain

$$\begin{aligned} p^n \mathbf{P}(t; p, q) &= \sum_{k=0}^n \left(p^k \frac{[n+1-k]_{p,q}}{[n+1]_{p,q}} \right) \mathbf{P}_k^0 B_{p,q}^{k,n+1}(t) \\ &+ \sum_{k=0}^n \left(1 - \frac{p^{k+1} [n-k]_{p,q}}{[n+1]_{p,q}} \right) \mathbf{P}_k^0 B_{p,q}^{k+1,n+1}(t). \end{aligned}$$

Now by shifting the limits, we have

$$\begin{aligned} p^n \mathbf{P}(t; p, q) &= \sum_{k=0}^{n+1} \left(p^k \frac{[n+1-k]_{p,q}}{[n+1]_{p,q}} \right) \mathbf{P}_k^0 B_{p,q}^{k,n+1}(t) \\ &+ \sum_{k=0}^{n+1} \left(1 - \frac{p^k [n+1-k]_{p,q}}{[n+1]_{p,q}} \right) \mathbf{P}_{k-1}^0 B_{p,q}^{k,n+1}(t), \end{aligned}$$

where \mathbf{P}_{-1}^0 is defined as the zero vector. Comparing coefficients on both side, we have

$$\mathbf{P}_k^* = p^n \mathbf{P}'_k = \left(1 - \frac{p^k [n+1-k]_{p,q}}{[n+1]_{p,q}}\right) \mathbf{P}_{k-1} + p^k \left(\frac{[n+1-k]_{p,q}}{[n+1]_{p,q}}\right) \mathbf{P}_k,$$

where $k = 0, 1, 2, \dots, n + 1$ and $\mathbf{P}_{-1} = \mathbf{P}_{n+1} = 0$.

When $p = 1$, formula (16) reduces to the degree evaluation formula of the q -Bézier curves. Let $P = (P_0, P_1, \dots, P_n)^T$ denote the vector of control points of the initial (p, q) -Bézier curve of degree n , and $\mathbf{P}^{(1)} = (P_0^*, P_1^*, \dots, P_{n+1}^*)$ the vector of control points of the degree elevated (p, q) -Bézier curve of degree $n + 1$, then, we can represent the degree elevation procedure as:

$$\mathbf{P}^{(1)} = T_{n+1}\mathbf{P},$$

where

$$T_{n+1} = \frac{1}{[n+1]_{p,q}} \times \begin{bmatrix} [n+1]_{p,q} & 0 & \dots & 0 & 0 \\ [n+1]_{p,q} - p[n]_{p,q} & p[n]_{p,q} & \dots & 0 & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & \dots & [n+1]_{p,q} - p^{n-1}[2]_{p,q} & p^{n-1}[2]_{p,q} & 0 \\ 0 & 0 & \dots & [n+1]_{p,q} - p^n[1]_{p,q} & p^n[1]_{p,q} \\ 0 & 0 & \dots & 0 & [n+1]_{p,q} \end{bmatrix}_{(n+2) \times (n+1)}.$$

For any $l \in \mathbb{N}$, the vector of control points of the degree elevated (p, q) -Bézier curves of degree $n + l$ is: $\mathbf{P}^{(l)} = T_{n+l} T_{n+2}, \dots, T_{n+1}\mathbf{P}$. As $l \rightarrow \infty$, the control polygon $\mathbf{P}^{(l)}$ converges to a (p, q) -Bézier curve.

4.3. de Casteljau algorithm:

(p, q) -Bézier curves of degree n can be written as two kinds of linear combination of two (p, q) -Bézier curves of degree $n - 1$, and we can get the two selectable algorithms to evaluate (p, q) -Bézier curves. The algorithms can be expressed as:

Algorithm 1.

$$\begin{cases} \mathbf{P}_i^0(t; p, q) \equiv \mathbf{P}_i^0 \equiv \mathbf{P}_i & i = 0, 1, 2, \dots, n \\ \mathbf{P}_i^r(t; p, q) = p^{r-1}t \mathbf{P}_{i+1}^{r-1}(t; p, q) + (q^k p^{r-i-1} - q^{r-1}t) \mathbf{P}_i^{r-1}(t; p, q) \\ r = 1, \dots, n, \quad i = 0, 1, 2, \dots, n - r, \end{cases} \tag{17}$$

or

$$\begin{cases} \mathbf{P}_i^0(t; p, q) \equiv \mathbf{P}_i^0 \equiv \mathbf{P}_i & i = 0, 1, 2, \dots, n \\ \mathbf{P}_i^r(t; p, q) = p^i q^{r-i-1} t \mathbf{P}_{i+1}^{r-1}(t; p, q) + (p^{r-1} - p^i q^{r-i-1} t) \mathbf{P}_i^{r-1}(t; p, q). \\ r = 1, \dots, n, \quad i = 0, 1, 2, \dots, n - r. \end{cases} \quad (18)$$

Then,

$$\mathbf{P}(t; p, q) = \sum_{i=0}^{n-1} \mathbf{P}_i^1(t; p, q) = \dots = \sum \mathbf{P}_i^r(t; p, q) B_{p,q}^{i,n-r}(t) = \dots = \mathbf{P}_0^n(t; p, q). \quad (19)$$

It is clear that the results can be obtained from Theorem (3.2). When $p = 1$, formula (17) and (18) recover the de Casteljau algorithms of classical q -Bézier curves. Let $P^0 = (P_0, P_1, \dots, P_n)^T$, $P^r = (P_0^r, P_1^r, \dots, P_{n-r}^r)^T$, then, de Casteljau algorithm can be expressed as:

Algorithm 2.

$$\mathbf{P}^r(t; p, q) = M_r(t; p, q) \dots M_2(t; p, q) M_1(t; p, q) \mathbf{P}^0, \quad (20)$$

where $M_r(t; p, q)$ is a $(n - r + 1) \times (n - r + 2)$ matrix and

$$M_r(t; p, q) =$$

$$\begin{bmatrix} (p^{r-1} - q^{r-1}t) & p^{r-1}t & \dots & 0 & 0 \\ 0 & (qp^{r-2} - q^{r-1}t) & p^{r-1}t & 0 & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & \dots & (q^{n-r-1}p^{2r-n-2} - q^{r-1}t) & p^{r-1}t & 0 \\ 0 & 0 & \dots & (q^{n-r}p^{2r-n-1} - q^{r-1}t) & p^{r-1}t \end{bmatrix}$$

or

$$M_r(t; p, q) = \begin{bmatrix} (p^{r-1} - q^{r-1}t) & q^{r-1}t & \dots & 0 & 0 \\ 0 & (p^{r-1} - pq^{r-2}t) & pq^{r-2}t & 0 & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & \dots & \dots & 0 & 0 \\ 0 & 0 & \dots & (p^{r-1} - p^{n-r}q^{-n-1}t) & p^{n-r}q^{-n-1}t \end{bmatrix}.$$

5. Tensor product (p, q) -Bernstein Bézier surfaces on $[0, 1] \times [0, 1]$

We define a two-parameter family $\mathbf{P}(u, v)$ of tensor product surfaces of degree $m \times n$ as follows:

$$\mathbf{P}(u, v) = \sum_{i=0}^m \sum_{j=0}^n \mathbf{P}_{i,j} B_{p_1, q_1}^{i,m}(u) B_{p_2, q_2}^{j,n}(v), \quad (u, v) \in [0, 1] \times [0, 1], \quad (21)$$

where $\mathbf{P}_{i,j} \in \mathbb{R}^3$ ($i = 0, 1, \dots, m, j = 0, 1, \dots, n$) and two real numbers $p_1 > q_1 > 0, p_2 > q_2 > 0$, $B_{p_1, q_1}^{i,m}(u), B_{p_2, q_2}^{j,n}(v)$ are (p, q) -analogue of Bernstein functions respectively with the parameter p_1, q_1 and p_2, q_2 . We call the parameter surface tensor product (p, q) -Bézier surface with degree $m \times n$. We refer to the $\mathbf{P}_{i,j}$ as the control points. By joining up adjacent points in the same row or column to obtain a net which is called the control net of tensor product (p, q) -Bézier surface.

5.1. Properties

1. Geometric invariance and affine invariance property:

Since

$$\sum_{i=0}^m \sum_{j=0}^n B_{p_1, q_1}^{i,m}(u) B_{p_2, q_2}^{j,n}(v) = 1, \quad (22)$$

$\mathbf{P}(u, v)$ is an affine combination of its control points.

2. Convex hull property: $\mathbf{P}(u, v)$ is a convex combination of $\mathbf{P}_{i,j}$ and lies in the convex hull of its control net.

3. Isoparametric curves property: The isoparametric curves $v = v^*$ and $u = u^*$ of a tensor product (p, q) -Bézier surface are respectively the (p, q) -Bézier curves of degree m and degree n , namely,

$$\mathbf{P}(u, v^*) = \sum_{i=0}^m \left(\sum_{j=0}^n \mathbf{P}_{i,j} B_{p_2, q_2}^{j,n}(v^*) \right) B_{p_1, q_1}^{i,m}(u), \quad u \in [0, 1],$$

$$\mathbf{P}(u^*, v) = \sum_{j=0}^n \left(\sum_{i=0}^m \mathbf{P}_{i,j} B_{p_1, q_1}^{i,m}(u^*) \right) B_{p_2, q_2}^{j,n}(v), \quad v \in [0, 1].$$

The boundary curves of $\mathbf{P}(u, v)$ are evaluated by $\mathbf{P}(u, 0), \mathbf{P}(u, 1), \mathbf{P}(0, v)$ and $\mathbf{P}(1, v)$.

4. Corner point interpolation property: The corner control net coincide with the four corners of the surface. Namely, $\mathbf{P}(0, 0) = \mathbf{P}_{0,0}, \mathbf{P}(0, 1) = \mathbf{P}_{0,n}, \mathbf{P}(1, 0) = \mathbf{P}_{m,0}, \mathbf{P}(1, 1) = \mathbf{P}_{m,n}$.

5. Reducibility: When $p_1 = p_2 = 1$, formula (21) reduces to a tensor product q -Bézier patch.

5.2. Degree elevation and de Casteljau algorithm

Let $\mathbf{P}(u, v)$ be a tensor product (p, q) -Bézier surface of degree $m \times n$. As an example, let us take obtaining the same surface as a surface of degree $(m + 1) \times (n + 1)$. Hence we need to find new control points $\mathbf{P}_{i,j}^*$ such that

$$\begin{aligned} \mathbf{P}(u, v) &= \sum_{i=0}^m \sum_{j=0}^n \mathbf{P}_{i,j} B_{p_1, q_1}^{i,m}(u) B_{p_2, q_2}^{j,n}(v) \\ &= \sum_{i=0}^{m+1} \sum_{j=0}^{n+1} \mathbf{P}_{i,j}^* B_{p_1, q_1}^{i,m+1}(u) B_{p_2, q_2}^{j,n+1}(v). \end{aligned} \quad (23)$$

$$\text{Let } \alpha_i = 1 - \frac{p_1^i [m+1-i]_{p_1, q_1}}{[m+1]_{p_1, q_1}}, \quad \beta_j = 1 - \frac{p_2^j [n+1-j]_{p_2, q_2}}{[n+1]_{p_2, q_2}}.$$

Then,

$$\mathbf{P}_{i,j}^* = \alpha_i \beta_j \mathbf{P}_{i-1, j-1} + \alpha_i (1 - \beta_j) \mathbf{P}_{i-1, j} + (1 - \alpha_i) (1 - \beta_j) \mathbf{P}_{i, j}, \quad (24)$$

which can be written in matrix form as

$$\begin{bmatrix} 1 - \frac{p_1^i [m+1-i]_{p_1, q_1}}{[m+1]_{p_1, q_1}} & \frac{p_1^i [m+1-i]_{p_1, q_1}}{[m+1]_{p_1, q_1}} \end{bmatrix} \begin{bmatrix} \mathbf{P}_{i-1, j-1} & \mathbf{P}_{i-1, j} \\ \mathbf{P}_{i, j-1} & \mathbf{P}_{i, j} \end{bmatrix} \begin{bmatrix} 1 - \frac{p_2^j [n+1-j]_{p_2, q_2}}{[n+1]_{p_2, q_2}} \\ \frac{p_2^j [n+1-j]_{p_2, q_2}}{[n+1]_{p_2, q_2}} \end{bmatrix}.$$

The de Casteljau algorithms are also easily extended to evaluate points on a (p, q) -Bézier surface. Given the control net $\mathbf{P}_{i,j} \in \mathbb{R}^3, i = 0, 1, \dots, m, j = 0, 1, \dots, n$, we have

$$\left\{ \begin{array}{l} \mathbf{P}_{i,j}^{0,0}(u, v) \equiv \mathbf{P}_{i,j}^{0,0} \equiv \mathbf{P}_{i,j} \quad i = 0, 1, 2, \dots, m; \quad j = 0, 1, 2, \dots, n. \\ \mathbf{P}_{i,j}^{r,r}(u, v) = \begin{bmatrix} (q_1^j p_1^{r-i-1} - q_1^{r-1} u) & p_1^{r-1} u \end{bmatrix} \begin{bmatrix} \mathbf{P}_{i,j}^{r-1, r-1} & \mathbf{P}_{i, j+1}^{r-1, r-1} \\ \mathbf{P}_{i+1, j}^{r-1, r-1} & \mathbf{P}_{i+1, j+1}^{r-1, r-1} \end{bmatrix} \begin{bmatrix} (q_2^j p_2^{r-j-1} - q_2^{r-1} v) \\ p_2^{r-1} v \end{bmatrix} \\ r = 1, \dots, k, \quad k = \min(m, n) \quad i = 0, 1, 2, \dots, m-r; \quad j = 0, 1, \dots, n-r \end{array} \right. \quad (25)$$

or

$$\left\{ \begin{array}{l} \mathbf{P}_{i,j}^{0,0}(u, v) \equiv \mathbf{P}_{i,j}^{0,0} \equiv \mathbf{P}_{i,j} \quad i = 0, 1, 2, \dots, m; \quad j = 0, 1, 2, \dots, n. \\ \mathbf{P}_{i,j}^{r,r}(u, v) = \begin{bmatrix} (p_1^{r-1} - p_1^i q_1^{r-i-1} u) & p_1^i q_1^{r-i-1} u \end{bmatrix} \begin{bmatrix} \mathbf{P}_{i,j}^{r-1, r-1} & \mathbf{P}_{i, j+1}^{r-1, r-1} \\ \mathbf{P}_{i+1, j}^{r-1, r-1} & \mathbf{P}_{i+1, j+1}^{r-1, r-1} \end{bmatrix} \begin{bmatrix} (p_2^{r-1} - p_2^j q_2^{r-j-1} v) \\ p_2^j q_2^{r-j-1} v \end{bmatrix} \\ r = 1, \dots, k, \quad k = \min(m, n) \quad i = 0, 1, 2, \dots, m-r; \quad j = 0, 1, \dots, n-r. \end{array} \right. \quad (26)$$

When $m = n$, one can directly use the algorithms above to get a point on the surface. When $m \neq n$, to get a point on the surface after k applications of formula (25) or (26), we perform formula (20) for the intermediate point $\mathbf{P}_{i,j}^{k,k}$.

Note: We get q -Bézier curves and surfaces for $(u, v) \in [0, 1] \times [0, 1]$ when we set the parameter $p_1 = p_2 = 1$ as proved in [Disibuyuk and Oruc (2008)] and [Disibuyuk (2009)].




Figure 3. The effect of the shape of (p, q) -Bézier curves.




Figure 4. The effect of the shape of (p, q) -Bézier curves.

6. Shape control of (p, q) -Bernstein curves

We have constructed (p, q) -Bernstein functions which holds both the end point interpolation property as shown in Figure 1 and 2. Parameter p and q have been used to control the shape of curves which can be observed through Figures 3-7.




Figure 5. The effect of the shape of (p, q) -Bézier curves.




Figure 6. The effect of the shape of (p, q) -Bézier curves.

7. Conclusion

In this paper, we have further studied various results for (p, q) -Bernstein basis function used in (p, q) -Bernstein operators (8). In comparison to q -Bézier curves based on q -Bernstein polynomials, this generalization gives us more flexibility in controlling the shapes of curves.




Figure 7. The effect of the shape of (p, q) -Bézier curves.

REFERENCES

- Acar, T. (2016). (p, q) -Generalization of Szász-Mirakyan operators. *Math. Methods Appl. Sci.* 39(10), 2685-2695.
- Acar, T. Aral, A. Mohiuddine, S.A. (2016). On Kantorovich modification of (p, q) -Baskakov operators, *Journal of Inequalities and Applications*, 98: DOI:10.1186/s13660-016-1045-9.
- Agarwal, P. N. Gupta, V., Kumar, S. (2013). On q -analogue of Bernstein-Schurer-Stancu operators, *Appl. Math. Comput.*, 219, 7754-7764.
- Aral, A. Gupta, V. Agarwal, R.P. (2013). Applications of q -Calculus in Operator Theory, *Springer*, New York.
- Bernstein, S. N. (1912-1913). Constructive proof of Weierstrass approximation theorem, *Comm. Kharkov Math. Soc.* 13(1-2).
- Bézier, P.E. (1972). Numerical Control-Mathematics and applications, *John Wiley and Sons*, London.
- Cai, Q. B. Zhou, G. (2016). On (p, q) -analogue of Kantorovich type Bernstein Stancu Schurer operators, *Appl. Math. Comput.*, 276 12-20.
- Disibuyuk, C. and Oruc, H. (2008). Tensor Product q -Bernstein Polynomials, *BIT Numerical Mathematics*, Springer 48, 689-700.
- Disibuyuk, C. (2009). "Tensor Product q -Bernstein Bézier Patches", *Lecture Notes in Computer Science*.
- Farouki, R. T. Rajan, V. T. (1988). Algorithms for polynomials in Bernstein form, *Computer Aided Geometric Design*, Volume 5, Issue 1.
- Hana, L.W. Chu, Y. Qiu, Z. Y. (2014). Generalized Bézier curves and surfaces based on Lupaş q -analogue of Bernstein operator, *Journal of Computational and Applied Mathematics* 261,

352-363.

- Hounkonnou, M. N. Désiré J. Kyemba, B. (2013). $\mathcal{R}(p, q)$ -calculus: differentiation and integration, *SUT Journal of Mathematics*, Vol. 49, No. 2, 145-167.
- Jagannathan, R. Srinivasa, R. K. (2005). Two-parameter quantum algebras, twin-basic numbers, and associated generalized hypergeometric series, *Proceedings of the International Conference on Number Theory and Mathematical Physics*, 20-21.
- Kadak, U. (2016). On weighted statistical convergence based on (p, q) -integers and related approximation theorems for functions of two variables, *Journal of Mathematical Analysis and Applications*, DOI: 10.1016/j.jmaa.2016.05.062.
- Kadak, U. (2017). Weighted statistical convergence based on generalized difference operator involving (p, q) -Gamma function and its applications to approximation theorems, *Journal of Mathematical Analysis and Applications*, Volume 448, Issue 2, Pages 1633-1650.
- Khan, K. Lobiyal, D.K. (2017). Bézier curves based on Lupas (p, q) -analogue of Bernstein functions in CAGD, *Journal of Computational and Applied Mathematics* Volume 317, Pages 458-477.
- Khan, K. Lobiyal, D.K. (2015). Bézier curves and surfaces based on modified Bernstein polynomials, *arXiv:1511.06594v1*. .
- Korovkin, P. P. (1960). Linear operators and approximation theory, *Hindustan Publishing Corporation, Delhi*.
- Lupaş, A. (1987). A q -analogue of the Bernstein operator, *Seminar on Numerical and Statistical Calculus, University of Cluj-Napoca*, 9,85-92.
- Oruk, H. Phillips, G. M. (2003). q -Bernstein polynomials and Bézier curves, *Journal of Computational and Applied Mathematics* 151, 1-12.
- Phillips, G.M. (1997). Bernstein polynomials based on the q -integers, *The heritage of P.L. Chebyshev, Ann. Numer. Math.*, 4, 511–518.
- Mahmudov N. I. and Sabancigil (2010). Some approximation properties of Lupaş q -analogue of Bernstein operators, *arXiv:1012.4245v1 [math.FA]*.
- Mishra, V.N. Pandey, S. (2016). On Chlodowsky variant of (p, q) -Kantorovich-Stancu-Schurer operators, *International Journal of Analysis and Applications*, Vol. 11, No. 1, pp. 28-39.
- Mishra, V.N. Pandey, S. (2017). On (p, q) Baskakov-Durrmeyer-Stancu Operators, *Advances in Applied Clifford Algebras*, Vol. 27, Issue 2, pp. 1633-164 DOI:10.1007/s00006-016-0738-y.
- Mishra, V.N. Khatri, K. Mishra, L.N. Deepmala (2013). Inverse result in simultaneous approximation by Baskakov-Durrmeyer-Stancu operators, *Journal of Inequalities and Applications*, 2013:586. doi:10.1186/1029-242X-2013-586.
- Mursaleen, M. Ansari, K. J. Khan, A. (2015). On (p, q) -analogue of Bernstein Operators, *Appl. Math. Comput.*, 266, 874-882, [Erratum: *Appl. Math. Comput.*(2016), 278, 70-71.]
- Mursaleen, M. and Nasiruzzaman M. and Nurgali, A. (2015). Some approximation results on Bernstein-Schurer operators defined by (p, q) -integers, *Journal of Inequalities and Applications* , 249, DOI 10.1186/s13660-015-0767-4.
- Mursaleen, M. Ansari, K. J. and Khan, A. (2015). Some Approximation Results by (p, q) -analogue of Bernstein-Stancu Operators, *Appl. Math. Comput.*, 264, 392-402.
- Mursaleen, M. Ansari, K. J. and Khan, A. (2016). Some approximation results for Bernstein-Kantorovich operators based on (p, q) -calculus, *U.P.B. Sci. Bull., Series A* Vol. 78, Iss. 4,

ISSN 1223-7027.

- Mursaleen, M. Khan, F and Khan, A. (2016). Approximation by (p, q) -Lorentz polynomials on a compact disk, *Complex Anal. Oper. Theory*, 10(8), 1725-1740.
- Mursaleen, M. Nasiruzzaman, M. Khan, A. and Ansari, K. J. (2016). Some approximation results on Bleimann-Butzer-Hahn operators defined by (p, q) -integers, *Filomat* 30:3 , 639-648, DOI 10.2298/FIL1603639M.
- Mursaleen, M. Khan, A. (2013). Generalized q -Bernstein-Schurer Operators and Some Approximation Theorems, *Journal of Function Spaces and Applications*, Article ID 719834, 7 pages <http://dx.doi.org/10.1155/2013/719834>
- Mursaleen, M. Alotaibi, A. Ansari, K.J. (2016). On a Kantorovich variant of (p, q) -Szász-Mirakjan operators, *Jour. Func. Spaces*, Article ID 1035253, 9 pages.
- Mursaleen, M. and Nasiruzzaman M. (2017). Some approximation properties of bivariate Bleimann-Butzer-Hahn operators based on (p, q) -integers, *M. Boll Unione Mat Ital* 10, 271.
- Mursaleen, M. Sarsenbi, A.M. Khan, T. (2016). On (p, q) -analogue of two parametric Stancu-Beta operators, *J. Ineq. Appl.*, 2016:190.
- Rababah, A. Manna, S. (2011). Iterative process for G2-multi degree reduction of Bézier curves, *Appl. Math. Comput.* 217, 8126-8133.
- Sederberg, T. W. (2014). Computer Aided Geometric Design Course Notes, Department of Computer Science *Brigham Young University*, October.
- Victor, K. Pokman, C. (2002). Quantum Calculus, *Springer-Verlag*, New York Berlin Heidelberg.
- Wafi, A. Rao, N. (2017). Bivariate-Schurer-Stancu operators based on (p, q) -integers, *Filomat*, (In press).
- Wafi, A. Rao, N. (2016). (p, q) -Bivariate-Bernstein-Chlowdosky Operators, *Filomat*, <http://journal.pmf.ni.ac.rs/filomat/index.php/filomat/author/index>.