



12-2018

Generalized Sylvester Polynomials of in Several Variables

Nejla Özmen
Düzce University

Sule Soytürk
Düzce University

Follow this and additional works at: <https://digitalcommons.pvamu.edu/aam>

 Part of the [Special Functions Commons](#)

Recommended Citation

Özmen, Nejla and Soytürk, Sule (2018). Generalized Sylvester Polynomials of in Several Variables, Applications and Applied Mathematics: An International Journal (AAM), Vol. 13, Iss. 2, Article 29. Available at: <https://digitalcommons.pvamu.edu/aam/vol13/iss2/29>

This Article is brought to you for free and open access by Digital Commons @PVAMU. It has been accepted for inclusion in Applications and Applied Mathematics: An International Journal (AAM) by an authorized editor of Digital Commons @PVAMU. For more information, please contact hvkoshy@pvamu.edu.



Generalized Sylvester Polynomials of in Several Variables

¹Nejla Özmen and ²Şule Soytürk

Department of Mathematics
 Faculty of Art and Science
 Düzce University
 Konuralp, TR-81620
 Düzce, Turkey

¹nejlaozmen06@gmail.com; ¹nejlaozmen@duzce.edu.tr;

²sule.soyturk61@gmail.com

Received: June 26, 2018; Accepted: October 22, 2018

Abstract

This study deals with some new properties for the Generalized Sylvester polynomials in several variables. Some properties of these polynomials were given. We also derive an application giving certain families of bilateral generating functions for the Generalized Sylvester polynomials in several variables. At the end, we discuss some special cases.

Keywords: Generalized Sylvester polynomials; generating function; recurrence relation; hypergeometric function.

MSC 2010 No.: 33C45, 33D65

1. Introduction

The Sylvester polynomials $\phi_n(x)$ are defined by (see, [Srivastava and Manocha (1984)])

$$\sum_{n=0}^{\infty} \phi_n(x)t^n = (1-t)^{-x}e^{xt}, \quad (1)$$

$$|t| < 1.$$

In Srivastava and Manocha (1984), from (1), we have

$$\phi_n(x) = \frac{x^n}{n!} {}_2F_0[-n, x; -; -x^{-1}],$$

where ${}_2F_0$ is Gauss's hypergeometric series. The generalized hypergeometric series ${}_pF_q$ defined and by

$${}_pF_q \left[\begin{matrix} \alpha_1, \dots, \alpha_p; \\ \beta_1, \dots, \beta_q; \end{matrix} z \right] = \sum_{n=0}^{\infty} \frac{(\alpha_1)_n \dots (\alpha_p)_n z^n}{(\beta_1)_n \dots (\beta_q)_n n!} \\ = {}_pF_q(\alpha_1, \dots, \alpha_p; \beta_1, \dots, \beta_q; z).$$

Here, $(\lambda)_\nu$ denotes the Pochhammer symbol defined by

$$(\lambda)_\nu = \begin{cases} 1, & \text{if } \nu = 0; \lambda \in \mathbb{C} \setminus \{0\} \\ \lambda(\lambda+1)\dots(\lambda+n-1), & \text{if } \nu = n \in \mathbb{N}; \lambda \in \mathbb{C}. \end{cases}$$

In Choi et al. (2017), introduced the generalized Sylvester polynomials of three variables denoted by $f_n(x, y, z; a, b, c, d, e, h)$ as follows:

$$f_n(x, y, z; a, b, c, d, e, h) = \frac{(dx)^n (ey)^n (hz)^n}{n!} F^{(3)} \left[\begin{matrix} -n :: -; -; - : ax; by; cz; \\ - :: -; -; - : -; -; -; \end{matrix} \begin{matrix} -\frac{1}{dx}, -\frac{1}{ey}, -\frac{1}{hz} \end{matrix} \right], \quad (2)$$

where $F^{(3)}[x, y, z]$ is the general triple hypergeometric series in Srivastava and Karlsson (1985). From (2), we have

$$f_n(x, y, z; a, b, c, d, e, h) = \frac{(dx)^n (ey)^n (hz)^n}{n!} \\ \times \sum_{r,s,k=0}^{\infty} \frac{(-n)_{r+s+k} (ax)_r (by)_s (cz)_k}{r!s!k!} \left(-\frac{1}{dx}\right)^r \left(-\frac{1}{ey}\right)^s \left(-\frac{1}{hz}\right)^k \\ = \sum_{r=0}^n \sum_{s=0}^{n-r} \sum_{k=0}^{n-r-s} \frac{(ax)_r (by)_s (cz)_k (dx)^n (ey)^n (hz)^n}{r!s!k!(n-r-s-k)! (dx)^r (ey)^s (hz)^k}.$$

The generalized Sylvester polynomials of three variables $f_n(x, y, z; a, b, c, d, e, h)$ have the following generating function in Choi et al. (2017):

$$\sum_{n=0}^{\infty} f_n(x, y, z; a, b, c, d, e, h) t^n \quad (3) \\ = e^{dxyhzt} (1 - eyhzt)^{-ax} (1 - dxhzt)^{-by} (1 - dxyt)^{-cz}.$$

In Choi et al. (2017), we have the following generating function

$$\sum_{n=0}^{\infty} \binom{n+k}{n} f_{n+k}(x, y, z; a, b, c, d, e, h) t^n \tag{4}$$

$$= e^{dehxyzt} (1 - ehzyt)^{-ax-k} (1 - dhxzt)^{-by-k} (1 - dexyt)^{-cz-k}$$

$$\times f_k(x, y, z; a, b, c, d(1 - ehzyt), e(1 - dhxzt), h(1 - dexyt)).$$

In this study, various families of multilinear and multilateral generating functions, miscellaneous properties and also some special cases for these polynomials are obtained. In addition, we derive a theorem giving certain families of bilateral generating functions for the generalized Sylvester polynomials of three variables and the Appell functions.

2. Summation Formula

Lemma 2.1.

The following addition expression holds for the generalized Sylvester polynomials of three variables:

$$f_n(x_1 + x_2, y, z; a, b, c, d, e, h) \tag{5}$$

$$= \sum_{m=0}^n \frac{1}{(1 - dx_1hzt)^m (1 - dx_1eyt)^m} f_{n-m}(x_1, y, z; a, b, c, d, e, h)$$

$$\times f_m(x_2, y(1 - dx_1hzt), z(1 - dx_1eyt); a, b, c, d, e, h).$$

Proof:

Replacing x by $x_1 + x_2$ in (3), we obtain

$$\sum_{n=0}^{\infty} f_n(x_1 + x_2, y, z; a, b, c, d, e, h) t^n$$

$$= e^{(x_1+x_2)deyhzt} (1 - eyhzt)^{-a(x_1+x_2)}$$

$$\times (1 - d(x_1 + x_2)hzt)^{-by} (1 - d(x_1 + x_2)eyt)^{-cz}$$

$$= (1 - eyhzt)^{-ax_1} e^{dx_1eyhzt} (1 - eyhzt)^{-ax_2} e^{dx_2eyhzt}$$

$$\times (1 - dx_1hzt - dx_2hzt)^{-by} (1 - dx_1eyt - dx_2eyt)^{-cz}$$

$$= e^{dx_1eyhzt} (1 - eyhzt)^{-ax_1} (1 - dx_1hzt)^{-by} (1 - dx_1eyt)^{-cz}$$

$$\times e^{dx_2eyhzt} (1 - eyhzt)^{-ax_2} \left(1 - \frac{dx_2hzt}{1 - dx_1hzt}\right)^{-by} \left(1 - \frac{dx_2eyt}{1 - dx_1eyt}\right)^{-cz}$$

$$\begin{aligned}
&= \sum_{n=0}^{\infty} f_n(x_1, y, z; a, b, c, d, e, h) t^n \left(e^{\frac{dx_2 eyhzt(1-dx_1 hzt)(1-dx_1 eyt)}{(1-dx_1 hzt)(1-dx_1 eyt)}} \right) \\
&\quad \times \left(1 - eyhzt \frac{(1-dx_1 hzt)(1-dx_1 eyt)}{(1-dx_1 hzt)(1-dx_1 eyt)} \right)^{-ax_2} \\
&\quad \times \left(1 - dx_2 hzt \frac{(1-dx_1 eyt)}{(1-dx_1 hzt)(1-dx_1 eyt)} \right)^{-by} \\
&\quad \times \left(1 - dx_2 eyt \frac{(1-dx_1 hzt)}{(1-dx_1 eyt)(1-dx_1 hzt)} \right)^{-cz} \\
&= \sum_{n=0}^{\infty} f_n(x_1, y, z; a, b, c, d, e, h) t^n \\
&\quad \times \sum_{m=0}^{\infty} f_m(x_2, y(1-dx_1 hzt), z(1-dx_1 eyt); a, b, c, d, e, h) \frac{t^m}{(1-dx_1 hzt)^m (1-dx_1 eyt)^m} \\
&= \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} f_n(x_1, y, z; a, b, c, d, e, h) f_m(x_2, y(1-dx_1 hzt), z(1-dx_1 eyt); a, b, c, d, e, h) \\
&\quad \times \frac{t^{n+m}}{(1-dx_1 hzt)^m (1-dx_1 eyt)^m} \\
&= \sum_{n=0}^{\infty} \sum_{m=0}^n f_{n-m}(x_1, y, z; a, b, c, d, e, h) f_m(x_2, y(1-dx_1 hzt), z(1-dx_1 eyt); a, b, c, d, e, h) \\
&\quad \times \frac{t^n}{(1-dx_1 hzt)^m (1-dx_1 eyt)^m}.
\end{aligned}$$

Comparing of the coefficients of t^n , lemma is proved. ■

3. Bilinear and Bilateral Generating Functions

In this section, with the help of the similar method as considered in Ozmen et al. (2018) and Ozmen (2015), we derive several families of bilinear and bilateral generating functions for the generalized Sylvester polynomials of three variables given by (3).

Theorem 3.1.

For a non-vanishing function $\Omega_{\mu}(y_1, \dots, y_r)$ of r complex variables y_1, \dots, y_r ($r \in \mathbb{N}$) and of complex order μ, ψ , let

$$\Lambda_{\mu, \psi}(y_1, \dots, y_r; \zeta) = \sum_{k=0}^{\infty} a_k \Omega_{\mu+\psi k}(y_1, \dots, y_r) \zeta^k \quad (a_k \neq 0),$$

and

$$\Theta_{n,p}^{\mu,\psi}(x; c; y_1, \dots, y_r; \xi) = \sum_{k=0}^{[n/p]} a_k f_{n-pk}(x, y, z; a, b, c, d, e, h) \Omega_{\mu+\psi k}(y_1, \dots, y_r) \xi^k. \tag{6}$$

Then, for $p \in \mathbb{N}$, we have

$$\begin{aligned} & \sum_{n=0}^{\infty} \Theta_{n,p}^{\mu,\psi} \left(x, y, z; a, b, c, d, e, h; y_1, \dots, y_r; \frac{\eta}{t^p} \right) t^n \\ &= e^{dxyhzt} (1 - eyhzt)^{-ax} (1 - dxhzt)^{-by} (1 - dxyt)^{-cz} \\ & \quad \times \Lambda_{\mu,\psi}(y_1, \dots, y_r; \eta). \end{aligned} \tag{7}$$

Proof:

For convenience, let S denote the first member of the assertion (7) of Theorem 3.1. Then, plugging the polynomials

$$\Theta_{n,p}^{\mu,\psi} \left(x, y, z; a, b, c, d, e, h; y_1, \dots, y_r; \frac{\eta}{t^p} \right),$$

which comes from (6) into the left-hand side of (7), we obtain

$$S = \sum_{n=0}^{\infty} \sum_{k=0}^{[n/p]} a_k f_{n-pk}(x, y, z; a, b, c, d, e, h) \Omega_{\mu+\psi k}(y_1, \dots, y_r) \eta^k t^{n-pk}. \tag{8}$$

Upon changing the order of summation in (8), if we replace n by $n + pk$, we can write

$$\begin{aligned} S &= \sum_{n=0}^{\infty} \sum_{k=0}^{\infty} a_k f_n(x, y, z; a, b, c, d, e, h) \Omega_{\mu+\psi k}(y_1, \dots, y_r) \eta^k t^n \\ &= \sum_{n=0}^{\infty} f_n(x, y, z; a, b, c, d, e, h) t^n \sum_{k=0}^{\infty} a_k \Omega_{\mu+\psi k}(y_1, \dots, y_r) \eta^k \\ &= e^{dexyzt} (1 - eyhzt)^{-ax} (1 - dxhzt)^{-by} (1 - dxyt)^{-cz} \\ & \quad \times \Lambda_{\mu,\psi}(y_1, \dots, y_r; \eta), \end{aligned}$$

the proof is completed. ■

Theorem 3.2.

Corresponding to an identically non-vanishing function $\Omega_\mu(y_1, \dots, y_r)$ of r complex variables y_1, \dots, y_r ($r \in \mathbb{N}$) and of complex order μ, ψ , let

$$\begin{aligned} & \Lambda_{\mu, \psi}^{n,p}(x_1 + x_2; c; y_1, \dots, y_r; t) \\ &= \sum_{k=0}^{[n/p]} a_k f_{n-pk}(x_1 + x_2, y, z; a, b, c, d, e, h) \\ & \quad \times \Omega_{\mu+\psi k}(y_1, \dots, y_r) t^k, \end{aligned}$$

where $a_k \neq 0$, $n, p \in \mathbb{N}$. Then, we have

$$\begin{aligned} & \sum_{k=0}^n \sum_{l=0}^{[k/p]} a_l \frac{1}{[(1-dx_1hzt)(1-dx_1eyt)]^{k-pl}} f_{n-k}(x_1, y, z; a, b, c, d, e, h) \\ & \quad \times f_{k-pl}(x_2, y(1-dx_1hzt), z(1-dx_1eyt); a, b, c, d, e, h) \\ & \quad \times \Omega_{\mu+\psi l}(y_1, \dots, y_r) t^l \\ &= \Lambda_{\mu, \psi}^{n,p}(x_1 + x_2, y, z; a, b, c, d, e, h; y_1, \dots, y_r; t), \end{aligned} \tag{9}$$

provided that each member of (9) exists.

Proof:

For convenience, let T denote the first member of the assertion (9) of Theorem 3.2. Then, upon substituting for the polynomials $f_{n-pk}(x_1 + x_2, y, z; a, b, c, d, e, h)$ from the (5) into the left-hand side of (9), we obtain

$$\begin{aligned} T &= \sum_{l=0}^{[n/p]} \sum_{k=0}^{n-pl} a_l \frac{1}{[(1-dx_1hzt)(1-dx_1eyt)]^k} f_{n-k-pl}(x_1, y, z; a, b, c, d, e, h) \\ & \quad \times f_k(x_2, y(1-dx_1hzt), z(1-dx_1eyt); a, b, c, d, e, h) \Omega_{\mu+\psi l}(y_1, \dots, y_r) t^l \\ &= \sum_{l=0}^{[n/p]} a_l \left(\sum_{k=0}^{n-pl} \frac{1}{[(1-dx_1hzt)(1-dx_1eyt)]^k} f_{n-k-pl}(x_1, y, z; a, b, c, d, e, h) \right) \\ & \quad \times f_k(x_2, y(1-dx_1hzt), z(1-dx_1eyt); a, b, c, d, e, h) \\ & \quad \times \Omega_{\mu+\psi l}(y_1, \dots, y_r) t^l \\ &= \sum_{l=0}^{[n/p]} a_l f_{n-pl}(x_1 + x_2, y, z; a, b, c, d, e, h) \Omega_{\mu+\psi l}(y_1, \dots, y_r) t^l \\ &= \Lambda_{\mu, \psi}^{n,p}(x_1 + x_2, y, z; a, b, c, d, e, h; y_1, \dots, y_r; t). \quad \blacksquare \end{aligned}$$

Theorem 3.3.

Corresponding to an identically non-vanishing function $\Omega_\mu(y_1, \dots, y_r)$ of r complex variables y_1, \dots, y_r ($r \in \mathbb{N}$) and of complex order μ , let

$$\begin{aligned} &\Lambda_{\mu,p,q} [x, y, z; a, b, c, d, e, h; y_1, \dots, y_r; t] \\ &= \sum_{k=0}^{\infty} a_k f_{m+qk}(x, y, z; a, b, c, d, e, h) \Omega_{\mu+pk}(y_1, \dots, y_r) t^k, \end{aligned}$$

where $a_k \neq 0$ and

$$\theta_{n,p,q}(y_1, \dots, y_r; z) = \sum_{k=0}^{[n/q]} \binom{m+n}{n-qk} a_k \Omega_{\mu+pk}(y_1, \dots, y_r) z^k.$$

Then, for $p, q \in \mathbb{N}$; we have

$$\begin{aligned} &\sum_{n=0}^{\infty} f_{m+n}(x, y, z; a, b, c, d, e, h) \theta_{n,p,q}(y_1, \dots, y_r; z) t^n \tag{10} \\ &= e^{dehxyt} (1 - ehzyt)^{-ax} (1 - dhxzt)^{-by} (1 - dexty)^{-cz} \\ &\times \Lambda_{\mu,p,q} \left(\begin{matrix} x, y, z; a, b, c, d(1 - ehzyt), e(1 - dhxzt), h(1 - dexty) \\ ; y_1, \dots, y_r; z \left(\frac{t}{(1 - ehzyt)(1 - dhxzt)(1 - dexty)} \right)^q \end{matrix} \right), \end{aligned}$$

provided that each member of (10) exists.

Proof:

For convenience, let T denote the first member of the assertion (10) of Theorem 3.3. Then, we get

$$\begin{aligned} T &= \sum_{n=0}^{\infty} f_{m+n}(x, y, z; a, b, c, d, e, h) \\ &\times \sum_{k=0}^{[n/q]} \binom{m+n}{n-qk} a_k \Omega_{\mu+pk}(y_1, \dots, y_r) z^k t^n. \end{aligned}$$

Replacing n by $n + qk$ and then using (4), we may write that

$$\begin{aligned} T &= \sum_{n=0}^{\infty} \sum_{k=0}^{\infty} \binom{m+n+qk}{n} f_{m+n+qk}(x, y, z; a, b, c, d, e, h) \Omega_{\mu+pk}(y_1, \dots, y_r) z^k t^{n+qk} \\ &= \sum_{k=0}^{\infty} \left(\sum_{n=0}^{\infty} \binom{m+n+qk}{n} f_{m+n+qk}(x, y, z; a, b, c, d, e, h) t^n \right) \Omega_{\mu+pk}(y_1, \dots, y_r) (zt^q)^k \\ &= \sum_{k=0}^{\infty} a_k e^{dehxyzt} (1 - ehzyt)^{-ax-m-qk} (1 - dhxzt)^{-by-m-qk} (1 - dexty)^{-cz-m-qk} \\ &\times f_{m+qk}(x, y, z; a, b, c, d(1 - ehzyt), e(1 - dhxzt), h(1 - dexty)) \Omega_{\mu+pk}(y_1, \dots, y_r) (zt^q)^k \\ &= e^{dehxyzt} (1 - ehzyt)^{-ax-m} (1 - dhxzt)^{-by-m} (1 - dexty)^{-cz-m} \end{aligned}$$

$$\begin{aligned}
& \times \sum_{k=0}^{\infty} a_k f_{m+qk}(x, y, z; a, b, c, d(1-ehyzt), e(1-dhxzt), h(1-dexyt)) \\
& \quad \times \Omega_{\mu+pk}(y_1, \dots, y_r) \left(\frac{zt^q}{(1-ehyzt)^q(1-dhxzt)^q(1-dexyt)^q} \right)^k \\
& = e^{dehxyzt} (1-ehyzt)^{-ax-m} (1-dhxzt)^{-by-m} (1-dexyt)^{-cz-m} \\
& \quad \times \Lambda_{\mu,p,q} \left(x, y, z; a, b, c, d(1-ehyzt), e(1-dhxzt), h(1-dexyt); \right. \\
& \quad \quad \left. y_1, \dots, y_r; z \left(\frac{t}{(1-ehyzt)(1-dhxzt)(1-dexyt)} \right)^q \right).
\end{aligned}$$

The proof is completed. ■

4. Special Cases

When the multivariable function $\Omega_{\mu+\psi k}(y_1, \dots, y_r)$, $k \in \mathbb{N}_0$, $r \in \mathbb{N}$, is expressed in terms of simpler functions of one and more variables, then we can give further applications of the above theorems. We first set

$$\Omega_{\mu+\psi k}(y_1, \dots, y_r) = h_{\mu+\psi k}^{(\alpha_1, \dots, \alpha_r)}(y_1, \dots, y_r),$$

in Theorem 3.1, where the multivariable extension of the Lagrange-Hermite polynomials $h_{\mu+\psi k}^{(\alpha_1, \alpha_2, \dots, \alpha_r)}(x_1, \dots, x_r)$ [Altin et al. (2006)], generated by

$$\begin{aligned}
\prod_{j=1}^r \{(1-x_j t^j)^{-\alpha_j}\} &= \sum_{n=0}^{\infty} h_n^{(\alpha_1, \dots, \alpha_r)}(x_1, \dots, x_r) t^n, \\
\alpha &\in \mathbb{C}, \quad |t| < \min \left\{ |x_1|^{-1}, |x_2|^{-1/2}, \dots, |x_r|^{-1/r} \right\}.
\end{aligned} \tag{11}$$

We are thus led to the following result which provides a class of bilateral generating functions for the multivariable extension of the Lagrange-Hermite polynomials $h_{\mu+\psi k}^{(\alpha_1, \dots, \alpha_r)}(y_1, \dots, y_r)$ and the generalized Sylvester polynomials of three variables.

Corollary 4.1.

If

$$\begin{aligned}
& \Lambda_{\mu,\psi}(y_1, \dots, y_r; \zeta) \\
& = \sum_{k=0}^{\infty} a_k h_{\mu+\psi k}^{(\alpha_1, \dots, \alpha_r)}(y_1, \dots, y_r) \zeta^k, \quad (a_k \neq 0, \mu, \psi \in \mathbb{C}),
\end{aligned}$$

then we have

$$\sum_{n=0}^{\infty} \sum_{k=0}^{\lfloor n/p \rfloor} a_k f_{n-pk}(x, y, z; a, b, c, d, e, h) h_{\mu+\psi k}^{(\alpha_1, \dots, \alpha_r)}(y_1, \dots, y_r) \frac{\eta^k}{t^{pk}} t^n \tag{12}$$

$$= e^{dxyhzt}(1 - eyhzt)^{-ax}(1 - dxhzt)^{-by}(1 - dxeyt)^{-cz} \Lambda_{\mu,\psi}(y_1, \dots, y_r; \eta).$$

Remark 4.2.

Using the generating relation (12) for the multivariable polynomials $h_{\mu+\psi k}^{(\alpha_1, \dots, \alpha_r)}(x_1, \dots, x_r)$ and getting $a_k = 1, \mu = 0, \psi = 1$, we find that

$$\begin{aligned} & \sum_{n=0}^{\infty} \sum_{k=0}^{[n/p]} f_{n-pk}(x, y, z; a, b, c, d, e, h) h_k^{(\alpha_1, \dots, \alpha_r)}(y_1, \dots, y_r) \eta^k t^{n-pk} \\ &= e^{dxyhzt}(1 - eyhzt)^{-ax}(1 - dxhzt)^{-by} \\ & \quad \times (1 - dxeyt)^{-cz} \prod_{j=1}^r \{(1 - y_j \eta^j)^{-\alpha_j}\}, \end{aligned}$$

$$|\zeta| < \min \left\{ |y_1|^{-1}, |y_2|^{-1/2}, \dots, |y_r|^{-1/r} \right\}, \quad |\eta| < 1.$$

If we set $r = 3$ and

$$\Omega_{\mu+\psi k}(y_1, y_2, y_3) = f_{\mu+\psi k}(x_3, y, z; a, b, c, d, e, h),$$

in Theorem 3.2, we have the following bilinear generating functions for the generalized Sylvester polynomials of three variables.

Corollary 4.3.

If

$$\begin{aligned} & \Lambda_{\mu,\psi}^{n,p}(x_1 + x_2, y, z; a, b, c, d, e, h; x_3, y, z; a, b, c, d, e, h; t) \\ &= \sum_{k=0}^{[n/p]} a_k f_{n-pk}(x_1 + x_2, y, z; a, b, c, d, e, h) \\ & \quad \times f_{\mu+\psi k}(x_3, y, z; a, b, c, d, e, h) t^k, \quad (a_k \neq 0, \mu, \psi \in \mathbb{C}), \end{aligned}$$

then we get

$$\begin{aligned} & \sum_{k=0}^n \sum_{l=0}^{[k/p]} a_l \frac{1}{[(1 - dx_1 hzt)(1 - dx_1 eyt)]^{k-pl}} f_{n-k}(x_1, y, z; a, b, c, d, e, h) \\ & \quad \times f_{k-pl}(x_2, y(1 - dx_1 hzt), z(1 - dx_1 eyt); a, b, c, d, e, h) f_{\mu+\psi l}(x_3, y, z; a, b, c, d, e, h) t^l \\ &= \Lambda_{\mu,\psi}^{n,p}(x_1 + x_2, y, z; a, b, c, d, e, h; x_3, y, z; a, b, c, d, e, h; t). \end{aligned} \tag{13}$$

If we set

$$s = 1 \text{ and } \Omega_{\mu+\psi k}(y_1) = P_{\mu+\psi k}^{(\alpha,\beta)}(\omega),$$

in Theorem 3.3, where the classical Jacobi polynomials $P_n^{(\alpha,\beta)}(y)$ is generated by [Erdélyi et al. (1955)],

$$\sum_{n=0}^{\infty} P_n^{(\alpha,\beta)}(x) t^n = \frac{2^{\alpha+\beta}}{\rho} (1-t+\rho)^{-\alpha} (1+t+\rho)^{-\beta},$$

$$\left\{ \rho = (1-2xt+t^2)^{1/2} \right\}.$$

We get a family of the bilateral generating functions for the classical Jacobi polynomials and the generalized Sylvester polynomials of three variables as follows:

Corollary 4.4.

If

$$\Lambda_{\mu,p,q}[x, y, z; a, b, c, d, e, h; \omega; t]$$

$$= \sum_{n=0}^{\infty} a_n f_{m+qn}(x, y, z; a, b, c, d, e, h) P_{\mu+pn}^{(\alpha,\beta)}(\omega) t^k,$$

$$(a_n \neq 0, m \in \mathbb{N}_0, k \neq 0, \mu, \psi \in \mathbb{C}),$$

and

$$\theta_{n,p,q}(\omega; z) := \sum_{k=0}^{[n/q]} \binom{m+n}{n-qn} a_k P_{\mu+pn}^{(\alpha,\beta)}(\omega) z^k,$$

where $n, p \in \mathbb{N}$, then we get that

$$\sum_{n=0}^{\infty} f_{m+qn}(x, y, z; a, b, c, d, e, h) \theta_{n,p,q}(\omega; z) t^n \tag{14}$$

$$= e^{dehxyzt} (1-ehyzt)^{-ax-m} (1-dhxyz)^{-by-m} (1-dexyt)^{-cz-m}$$

$$\times \Lambda_{\mu,p,q} \left(\begin{matrix} x, y, z; a, b, c, d(1-ehyzt), e(1-dhxyz), h(1-dexyt) \\ \omega; z \left(\frac{t}{(1-ehyzt)(1-dhxyz)(1-dexyt)} \right)^q \end{matrix} \right).$$

Remark 4.5.

Furthermore, for every suitable choice of the coefficients a_k ($k \in \mathbb{N}_0$), if the multivariable functions $\Omega_{\mu+\psi k}(y_1, \dots, y_r)$, $r \in \mathbb{N}$, are expressed as an appropriate product of several simpler functions, the assertions of Theorem 3.1, Theorem 3.2, Theorem 3.3 can be applied in order to derive various families of multilinear and multilateral generating functions for the family of the generalized Sylvester polynomials of three variables given explicitly by (3).

5. Miscellaneous Properties

In this section, we give some properties for the generalized Sylvester polynomials of three variables $f_n(x, y, z; a, b, c, d, e, h)$ given by (3).

Theorem 5.1.

The generalized Sylvester polynomials of three variables $f_n(x, y, z; a, b, c, d, e, h)$ have the following integral representation:

$$\begin{aligned}
 & f_n(x, y, z; a, b, c, d, e, h) \\
 &= \frac{1}{n! \Gamma(ax) \Gamma(by) \Gamma(cz)} \sum_{m=0}^n \binom{n}{m} \\
 & \times \int_0^\infty \int_0^\infty \int_0^\infty e^{-(u_1+u_2+u_3)} u_1^{ax-1} u_2^{by-1} u_3^{cz-1} (dehxyz)^{n-m} \\
 & \times (ehyzu_1 + dhxzu_2 + dxyu_3)^m du_1 du_2 du_3.
 \end{aligned}$$

Proof:

If we use the identity

$$a^{-v} = \frac{1}{\Gamma(v)} \int_0^\infty e^{-at} t^{v-1} dt, \quad Re(v) > 0,$$

on the left-hand side of the generating function (3), we have

$$\begin{aligned}
 & \sum_{n=0}^\infty f_n(x, y, z; a, b, c, d, e, h) t^n \\
 &= e^{dehxyz t} \frac{1}{\Gamma(ax)} \int_0^\infty e^{-(1-ehyz t)u_1} u_1^{ax-1} du_1
 \end{aligned}$$

$$\begin{aligned}
 & \times \frac{1}{\Gamma(by)} \int_0^\infty e^{-(1-dhxyz)t} u_2^{by-1} du_2 \frac{1}{\Gamma(cz)} \int_0^\infty e^{-(1-dexyt)u_3} u_3^{cz-1} du_3 \\
 & = \sum_{n=0}^\infty \frac{(dehxyzt)^n}{n!} \frac{1}{\Gamma(ax)\Gamma(by)\Gamma(cz)} \int_0^\infty \int_0^\infty \int_0^\infty e^{-u_1-u_2-u_3} \\
 & \quad \times e^{(ehyzu_1+dhxzu_2+dexyu_3)t} u_1^{ax-1} u_2^{by-1} u_3^{cz-1} du_1 du_2 du_3 \\
 & = \frac{1}{\Gamma(ax)\Gamma(by)\Gamma(cz)} \sum_{n=0}^\infty \int_0^\infty \int_0^\infty \int_0^\infty \frac{(dehxyz)^n t^n}{n!} e^{-(u_1+u_2+u_3)} \\
 & \quad \times \sum_{m=0}^\infty \frac{(ehyzu_1+dhxzu_2+dexyu_3)^m t^m}{m!} u_1^{ax-1} u_2^{by-1} u_3^{cz-1} du_1 du_2 du_3 \\
 & = \sum_{n=0}^\infty \left(\frac{1}{n! \Gamma(ax)\Gamma(by)\Gamma(cz)} \sum_{m=0}^n \binom{n}{m} \int_0^\infty \int_0^\infty \int_0^\infty e^{-(u_1+u_2+u_3)} u_1^{ax-1} u_2^{by-1} u_3^{cz-1} \right. \\
 & \quad \left. (dehxyz)^{n-m} (ehyzu_1 + dhxzu_2 + dexyu_3)^m du_1 du_2 du_3 \right) t^n.
 \end{aligned}$$

From the coefficients of t^n on the both sides of the last equality, one can get the desired result. ■

We now discuss some miscellaneous recurrence relations of the generalized Sylvester polynomials of three variables. By differentiating each member of the generating function relation (3) with respect to x, y, z and using

$$\sum_{n=0}^\infty \sum_{k=0}^\infty A(k, n) = \sum_{n=0}^\infty \sum_{k=0}^n A(k, n - k),$$

we arrive at the following (differential) recurrence relation for the generalized Sylvester polynomials:

On the other hand, by differentiating each member of the generating function relation (3) with respect to x, y, z , we have

$$\begin{aligned}
 & \frac{\partial}{\partial x} f_n(x, y, z; a, b, c, d, e, h) \\
 & = (dehyz) f_{n-1}(x, y, z; a, b, c, d, e, h) \\
 & + (a) \sum_{m=0}^{n-1} \frac{(ehyz)^{m+1}}{(m+1)} f_{n-m-1}(x, y, z; a, b, c, d, e, h) \\
 & + (bdhxyz) \sum_{m=0}^{n-1} (dhxz)^m f_{n-m-1}(x, y, z; a, b, c, d, e, h)
 \end{aligned}$$

$$\begin{aligned}
 &+(cdeyz) \sum_{m=0}^{n-1} (dexy)^m f_{n-m-1}(x, y, z; a, b, c, d, e, h). \\
 &\quad \frac{\partial}{\partial y} f_n(x, y, z; a, b, c, d, e, h) \\
 &= (dehxz) f_{n-1}(x, y, z; a, b, c, d, e, h) \\
 &+(axeHz) \sum_{m=0}^{n-1} (ehyz)^m f_{n-m-1}(x, y, z; a, b, c, d, e, h) \\
 &+(b) \sum_{m=0}^{n-1} \frac{(dhxz)^{m+1}}{(m+1)} f_{n-m-1}(x, y, z; a, b, c, d, e, h) \\
 &+(cdez) \sum_{m=0}^{n-1} (dexy)^m f_{n-m-1}(x, y, z; a, b, c, d, e, h). \\
 &\quad \frac{\partial}{\partial z} f_n(x, y, z; a, b, c, d, e, h) \\
 &= (dehxy) f_{n-1}(x, y, z; a, b, c, d, e, h) \\
 &+(axeHy) \sum_{m=0}^{n-1} (ehyz)^m f_{n-m-1}(x, y, z; a, b, c, d, e, h) \\
 &+(bydhx) \sum_{m=0}^{n-1} (dhxz)^m f_{n-m-1}(x, y, z; a, b, c, d, e, h) \\
 &+(c) \sum_{m=0}^{n-1} \frac{(dexy)^{m+1}}{m+1} f_{n-m-1}(x, y, z; a, b, c, d, e, h),
 \end{aligned}$$

respectively.

Besides, by differentiating each member of the generating function relation (3) with respect to t , we have the following another recurrence relation for these polynomials:

$$\begin{aligned}
 &(n+1) f_{n+1}(x, y, z; a, b, c, d, e, h) \\
 &= (dehxyz) f_n(x, y, z; a, b, c, d, e, h) \\
 &+(ax) \sum_{m=0}^n (ehyz)^{m+1} f_{n-m}(x, y, z; a, b, c, d, e, h) \\
 &+(by) \sum_{m=0}^n (dhxz)^{m+1} f_{n-m}(x, y, z; a, b, c, d, e, h) \\
 &+(cz) \sum_{m=0}^n (dexy)^{m+1} f_{n-m}(x, y, z; a, b, c, d, e, h).
 \end{aligned}$$

6. The Generalized Sylvester polynomials of three variables and Appell Functions

In 2012, Liu et al. derived bilateral generating functions for the Erkus-Srivastava polynomials and generalized Lauricella functions [Liu et al. (2012)]. In the present section, we derive various families of bilateral generating functions for the generalized Sylvester polynomials of three variables and the Appell functions. On the other hand, the four Appell functions, denoted by F_1 , F_2 , F_3 and F_4 , were generalized by Lauricella functions of n variables which are denoted by $F_A^{(n)}$, $F_B^{(n)}$, $F_C^{(n)}$, $F_D^{(n)}$ and

$$F_A^{(2)} = F_2, F_B^{(2)} = F_3, F_C^{(2)} = F_4, F_D^{(2)} = F_1,$$

where

$$F_1[a, b, b'; c; x, y] = \sum_{m,n=0}^{\infty} \frac{(a)_{m+n}(b)_m(b')_n x^m y^n}{(c)_{m+n} m! n!},$$

$$F_2[a, b, b'; c, c'; x, y] = \sum_{m,n=0}^{\infty} \frac{(a)_{m+n}(b)_m(b')_n x^m y^n}{(c)_m(c')_n m! n!},$$

$$F_3[a, a', b, b'; c; x, y] = \sum_{m,n=0}^{\infty} \frac{(a)_m(a')_n(b)_m(b')_n x^m y^n}{(c)_{m+n} m! n!},$$

$$F_4[a, b; c, c'; x, y] = \sum_{m,n=0}^{\infty} \frac{(a)_{m+n}(b)_{m+n} x^m y^n}{(c)_m(c')_n m! n!}.$$

Theorem 6.1.

The following bilateral generating function holds true:

$$\begin{aligned} & \sum_{n=0}^{\infty} f_n(x, y, z; a, b, c, d, e, h) F_1(a_1, -n, b_2; c_1; u_1, u_2) t^n \\ &= e^{dxyhzt} (1 - eyhzt)^{-ax} (1 - dxhzt)^{-by} (1 - dxeyt)^{-cz} \\ & \times F_D^{(5)} \left[\begin{matrix} a_1, -, ax, by, cz, b_2; c_1; \\ -u_1xyzhzt, \frac{-u_1teyhz}{1-ehzt}, \frac{-u_1tdxhz}{1-dhzt}, \frac{-u_1tdxey}{1-dxeyt}, u_2 \end{matrix} \right], \end{aligned}$$

where $F_D^{(s)}$ is the Lauricella function.

Proof:

By using the relationship (4), it is easily observed that

$$\begin{aligned}
 & \sum_{n=0}^{\infty} f_n(x, y, z; a, b, c, d, e, h) F_1(a_1, -n, b_2; c_1; u_1, u_2) t^n \\
 &= \sum_{n=0}^{\infty} f_n(x, y, z; a, b, c, d, e, h) \\
 & \quad \times \sum_{m=0}^n \sum_{p=0}^{\infty} \frac{(a_1)_{m+p} (-n)_m (b_2)_p}{(c_1)_{m+p}} \frac{u_1^m u_2^p}{m! p!} t^n \\
 &= \sum_{m,p=0}^{\infty} \left(\sum_{n=0}^{\infty} \binom{n+m}{m} f_{n+m}(x, y, z; a, b, c, d, e, h) t^n \right) \\
 & \quad \times \frac{(a_1)_{m+p} (b_2)_p}{(c_1)_{m+p}} (-u_1 t)^m \frac{u_2^p}{p!} \\
 &= e^{dxyhzt} (1 - eyhzt)^{-ax} (1 - dxhzt)^{-by} (1 - dxyt)^{-cz} \\
 & \times \sum_{m,p=0}^{\infty} f_m(x, y, z; a, b, c, d(1 - ehzt), e(1 - dhzt), h(1 - dxyt)) \\
 & \quad \times \frac{(a_1)_{m+p} (b_2)_p}{(c_1)_{m+p}} \left(\frac{-u_1 t}{(1 - ehzt)(1 - dhzt)(1 - dxyt)} \right)^m \frac{u_2^p}{p!} \\
 &= e^{dxyhzt} (1 - eyhzt)^{-ax} (1 - dxhzt)^{-by} (1 - dxyt)^{-cz} \\
 & \quad \times \sum_{m,r,s,k,p=0}^{\infty} \frac{(a_1)_{m+r+s+k+p} (ax)_r (by)_s (cz)_k (b_2)_p}{(c_1)_{m+r+s+k+p}} \\
 & \quad \times \frac{(-u_1 xyzdeht)^m}{m!} \frac{(-\frac{u_1 teyhz}{1-dhzt})^s}{s!} \frac{(-\frac{u_1 tdxhz}{1-ehzt})^r}{r!} \frac{(-\frac{u_1 tdxyt}{1-dxyt})^k}{k!} \frac{u_2^p}{p!} \\
 &= e^{dxyhzt} (1 - eyhzt)^{-ax} (1 - dxhzt)^{-by} (1 - dxyt)^{-cz} \\
 & \quad \times F_D^{(5)} \left[\begin{matrix} a_1, -, ax, by, cz, b_2; c_1; \\ -u_1 xyzdeht, \frac{-u_1 teyhz}{1-ehzt}, \frac{-u_1 tdxhz}{1-dhzt}, \frac{-u_1 tdxyt}{1-dxyt}, u_2 \end{matrix} \right]. \quad \blacksquare
 \end{aligned}$$

Theorem 6.2.

The following bilateral generating function holds true:

$$\begin{aligned} & \sum_{n=0}^{\infty} f_n(x, y, z; a, b, c, d, e, h) F_2(a_1, -n, b_1; c_1, c_2; u_1, u_2) t^n \\ &= e^{dxyhzt} (1 - eyhzt)^{-ax} (1 - dxhzt)^{-by} (1 - dxyt)^{-cz} \\ & \times F_A^{(5)} \left[a_1, -, ax, by, cz, b_1; c_1, c_2; -u_1xyzdeht, \frac{-u_1teyhz}{1-ehyzt}, \frac{-u_1tdxhz}{1-dhxzt}, \frac{-u_1tdxey}{1-dxyt}, u_2 \right], \end{aligned}$$

where $F_A^{(s)}$ is the Lauricella function.

Proof:

By using the relationship (4), it is easily observed that

$$\begin{aligned} & \sum_{n=0}^{\infty} f_n(x, y, z; a, b, c, d, e, h) F_2(a_1, -n, b_1; c_1, c_2; u_1, u_2) t^n \\ &= \sum_{n=0}^{\infty} f_n(x, y, z; a, b, c, d, e, h) \sum_{m=0}^n \sum_{p=0}^{\infty} \frac{(a_1)_{m+p} (-n)_m (b_1)_p}{(c_1)_m (c_2)_p} \frac{u_1^m u_2^p}{m! p!} t^n \\ &= \sum_{m,p=0}^{\infty} \left(\sum_{n=0}^{\infty} \binom{n+m}{m} f_{n+m}(x, y, z; a, b, c, d, e, h) t^n \right) \frac{(a_1)_{m+p} (b_1)_p}{(c_1)_m (c_2)_p} (-u_1 t)^m \frac{u_2^p}{p!} \\ &= e^{dxyhzt} (1 - eyhzt)^{-ax} (1 - dxhzt)^{-by} (1 - dxyt)^{-cz} \\ & \times \sum_{m,p=0}^{\infty} f_m(x, y, z; a, b, c, d(1 - ehzt), e(1 - dxhzt), h(1 - dxyt)) \\ & \times \frac{(a_1)_{m+p} (b_1)_p}{(c_1)_m (c_2)_p} \left(\frac{-u_1 t}{(1 - ehzt)(1 - dxhzt)(1 - dxyt)} \right)^m \frac{u_2^p}{p!} \\ &= e^{dxyhzt} (1 - eyhzt)^{-ax} (1 - dxhzt)^{-by} (1 - dxyt)^{-cz} \\ & \times \sum_{m,r,s,k,p=0}^{\infty} \frac{(a_1)_{m+r+s+k+p} (ax)_r (by)_s (cz)_k (b_1)_p}{(c_1)_{m+r+s+k} (c_2)_p} \\ & \times \frac{(-u_1xyzdeht)^m}{m!} \frac{\left(-\frac{u_1teyhz}{1-dhxzt}\right)^s}{s!} \frac{\left(-\frac{u_1tdxhz}{1-ehyzt}\right)^r}{r!} \frac{\left(-\frac{u_1tdxey}{1-dxyt}\right)^k}{k!} \frac{u_2^p}{p!} \\ &= e^{dxyhzt} (1 - eyhzt)^{-ax} (1 - dxhzt)^{-by} (1 - dxyt)^{-cz} \\ & \times F_A^{(5)} \left[a_1, -, ax, by, cz, b_1; c_1, -, -, -, c_2; -u_1xyzdeht, \frac{-u_1teyhz}{1-ehyzt}, \frac{-u_1tdxhz}{1-dhxzt}, \frac{-u_1tdxey}{1-dxyt}, u_2 \right]. \quad \blacksquare \end{aligned}$$

7. Conclusion

In this paper, we obtain some new properties for the Generalized Sylvester polynomials in several variables. Various families of multilinear and multilateral generating functions and their miscellaneous properties are obtained. We also derive an application giving certain families of bilateral generating functions for the Generalized Sylvester polynomials in several variables. With the method used here, it is possible to obtain bilinear and bilateral generating functions for other polynomials.

Acknowledgement:

This work is supported by Düzce University Project BAP-2018.05.04.799.

REFERENCES

- Altın A. and Erkuş, E. (2006). On a multivariable extension of the Lagrange-Hermite polynomials, *Integral Transforms and Special Functions*, Vol. 17, No. 4, 239-244.
- Choi, J. and Akhlaq, M.(2017). Note on generating functions of generalized Sylvester polynomials of several variables, *Far East Journal of Mathematical Sciences*, Vol. 12, No. 11, pp. 2877-2894.
- Erdélyi, A., Magnus, W., Oberhettinger, F., Tricomi, F. G. (1955). *Higher Transcendental Functions*, Vol. III, McGraw-Hill Book Company, New York, Toronto and London.
- Erkuş, E. and Srivastava, H. M. (2006). A unified presentation of some families of multivariable polynomials, *Integral Transform Spec. Funct.* Vol. 17, pp. 267-273.
- Liu, S.-J. , Lin, S.-D., Srivastava, H.M. and Wong, M.-M. (2012). Bilateral generating functions for the Erkuş-Srivastava polynomials and the generalized Lauricella functions, *App. Mathematicis and Comp.*, Vol. 218, pp. 7685-7693.
- Ozmen, N. (2017). Some new properties of the Meixner polynomials, *Sakarya University Journal of Science*, Vol. 21, No. 6, pp. 1454-1462.
- Ozmen, N., Erkuş-Duman, E. (2013). Some results for a family of multivariable polynomials, *AIP Conference Proceedings*, Vol. 1558, pp. 1124-1127.
- Ozmen, N. and Erkuş-Duman, E. (2015). On the Poisson-Charlier polynomials, *Serdica Math. J.* Vol. 41, No. 4, pp. 457-470.
- Ozmen, N. and Erkuş-Duman, E. (2018). Some families of generating functions for the generalized Cesáro polynomials, *J. Comput. Anal. Appl.*, Vol. 25, No. 4, pp. 670-683.
- Srivastava, H. M. and Karlsson, P.W. (2017). *Multiple Gaussian hypergeometric Series*, Ellis Horwood Series: Mathematics and its Applications, Ellis Horwood Ltd., Chichester, Halsted Press, John Wiley and Sons, Inc., New York.
- Srivastava, H.M. and Manocha, H.L. (1984). *A Treatise on Generating Functions*, Halsted Press (Ellis Horwood Limited, Chichester), John Wiley and Sons, New York.