



6-2019

## On the Lucas Difference Sequence Spaces Defined by Modulus Function

Murat Karakaş  
*Bitlis Eren University*

Tayfur Akbaş  
*Bitlis Eren University*

Ayşe M. Karakaş  
*Bitlis Eren University*

Follow this and additional works at: <https://digitalcommons.pvamu.edu/aam>



Part of the [Number Theory Commons](#), and the [Other Applied Mathematics Commons](#)

### Recommended Citation

Karakaş, Murat; Akbaş, Tayfur; and Karakaş, Ayşe M. (2019). On the Lucas Difference Sequence Spaces Defined by Modulus Function, *Applications and Applied Mathematics: An International Journal (AAM)*, Vol. 14, Iss. 1, Article 15.

Available at: <https://digitalcommons.pvamu.edu/aam/vol14/iss1/15>

This Article is brought to you for free and open access by Digital Commons @PVAMU. It has been accepted for inclusion in *Applications and Applied Mathematics: An International Journal (AAM)* by an authorized editor of Digital Commons @PVAMU. For more information, please contact [hvkoshy@pvamu.edu](mailto:hvkoshy@pvamu.edu).



## On the Lucas Difference Sequence Spaces Defined by Modulus Function

<sup>1</sup>Murat Karakaş, <sup>1</sup>Tayfur Akbaş and <sup>2</sup>Ayşe Metin Karakaş

<sup>1</sup>Department of Mathematics  
 Bitlis Eren University  
 Bitlis, Turkey

[m.karakas33@hotmail.com](mailto:m.karakas33@hotmail.com); [tayfurakbas@hotmail.com](mailto:tayfurakbas@hotmail.com)

<sup>2</sup>Department of Statistics  
 Bitlis Eren University  
 Bitlis, Turkey

[aysekarakas5767@gmail.com](mailto:aysekarakas5767@gmail.com);

Received: March 12, 2018; Accepted: October 3, 2018

### Abstract

In this paper, firstly, we define the Lucas difference sequence spaces by the help of Lucas sequence and a sequence of modulus function. Besides, we give some inclusion relations and examine geometrical properties such as Banach-Saks type  $p$ , weak fixed point property.

**Keywords:** Difference sequence space; Lucas numbers; Modulus function; Sequence space; Banach-Saks property; Weak fixed point property; Infinite matrix

**MSC 2010 No:** 11B39, 46A45, 46B45, 46B20

### 1. Introduction

Let  $w$  be the space of all real and complex valued sequences. Each linear subspace of  $w$  is called sequence space. Throughout the paper  $\ell_\infty$ ,  $\ell_p$  ( $1 \leq p < \infty$ ),  $c$  and  $c_0$  denote the spaces of all bounded,  $p$ -absolutely summable, convergent and null sequences, respectively. If  $X$  is a complete linear metric space, then a  $K$ -space  $X$  is called an  $FK$ -space. An  $FK$ -space whose topology is normable is called  $BK$ -space.

An infinite matrix is a double sequence  $A = (a_{nk})$  of real or complex numbers defined by a function  $A$  from the set  $\mathbb{N} \times \mathbb{N}$  into the complex field  $\mathbb{C}$  (or  $\mathbb{R}$ ) where  $\mathbb{N} = \{0, 1, 2, \dots\}$ . The treatment of infinite matrices is absolutely different from finite matrices. There are various reasons for this. In some instances, the most general linear operator among two sequence spaces

is presented by an infinite matrix. Let  $X$  and  $Y$  be any two sequence spaces.  $A$  defines a matrix mapping from  $X$  into  $Y$ , if  $Ax = \{(Ax)_n\} \in Y$  for every  $x = (x_k) \in X$  where

$$(Ax)_n = \sum_k a_{nk}x_k. \quad (1)$$

The class of all matrices  $A$  such that  $A: X \rightarrow Y$  is symbolized by  $(X:Y)$ . In this way,  $A \in (X:Y)$  iff the series on the right hand side of (1) converges for each  $n \in \mathbb{N}$  and every  $x \in X$ , and we get  $Ax = \{(Ax)_n\} \in Y$  for all  $x \in X$ .

The concept of matrix domain is important for our study. For an infinite matrix  $A$ , the matrix domain  $\phi_A$  in a sequence space  $\phi$  is defined by

$$\phi_A = \{x \in \omega: Ax \in \phi\}, \quad (2)$$

which is a sequence space [Başar (2011)].

Recently, so many authors have made use of the approach of constructing a new sequence space using matrix domain for a triangle infinite matrix, e.g., Başar and Altay (2003), Altay and Basar (2005), Kirisci and Basar (2010), Mursaleen and Noman (2010), Mursaleen and Noman (2011), Kara and Basarir (2012), Kara (2013), (Debnath and Saha 2014), (Debnath et al. 2015), Karakas (2015).

In the literature, the matrix domain  $\lambda_\Delta$  is called the difference sequence space if  $\lambda$  is a normed or paranormed sequence space where  $\Delta$  symbolizes the following backward difference matrix  $\Delta = (\Delta_{nk})$  and  $\Delta' = (\Delta'_{nk})$  symbolizes the following transpose of the matrix  $\Delta$ , the forward difference matrix. For  $\lambda = \ell_p$ , this space is called as the space of sequences of  $p$ -bounded variation, that is,  $bv_p$ . Also, it is clear that  $bv_p = (\ell_p)_\Delta$ .

$$\Delta_{nk} = \begin{cases} (-1)^{n-k}, & (n-1 \leq k \leq n), \\ 0, & (0 \leq k < n-1 \text{ or } k > n), \end{cases} \quad \text{and} \quad \Delta'_{nk} = \begin{cases} (-1)^{n-k}, & (n \leq k \leq n+1), \\ 0, & (0 \leq k < n \text{ or } k > n+1). \end{cases}$$

The notion of difference sequence spaces was first defined by Kizmaz (1981) in the form of  $X(\Delta) = \{x \in \omega: x_k - x_{k+1} \in X\}$  for  $X = \ell_\infty, c, c_0$ . These spaces were generalized by Et and Çolak (1995) as  $X(\Delta^r) = \{x \in \omega: \Delta^r x \in X\}$ ,  $X = \ell_\infty, c, c_0$ .

The difference sequence space  $bv_p$  which is examined by Altay and Başar (2007) contains sequences  $(x_k)$  such that  $(x_k - x_{k-1})$  for  $0 < p < 1$ . In the case  $1 \leq p \leq \infty$ , this space was studied by Çolak et al. (2004). Also, many authors analyzed the certain difference sequence spaces, see Ahmad (1987), Malkowsky (1989), Et (1993), Mursaleen (1996), Et and Basarir (1997), Tripathy (2003), Colak and Et (2005), Başar et al. (2008).

The main aim of this note is to introduce new sequence spaces  $\ell(\hat{E}(r, s), \mathfrak{F}, p, u)$  and  $\ell_\infty(\hat{E}(r, s), \mathfrak{F}, p, u)$  with the help of sequence of modulus functions, non-zero real numbers  $r$  and  $s$ , Lucas difference matrix and its matrix domain. In addition, we investigate some topological properties and also find out some inclusion relations concerning with these spaces. Finally, we work through geometrical properties of the space  $\ell(\hat{E}(r, s), \mathfrak{F}, p, u)$ .

## 2. Material and methods

In this part of our study, we inform about familiar concepts which are necessary for us later on the paper.

**Definition 2.1.**

A modulus  $f$  is a function from  $[0, \infty)$  to  $[0, \infty)$  such that  $f(x) = 0$  if and only if  $x = 0$ ;  $f(x + y) \leq f(x) + f(y)$  for  $x, y \geq 0$ ;  $f$  is increasing;  $f$  is continuous from the right at 0. [Nakano (1953)].

It follows that  $f$  must be continuous everywhere on  $[0, \infty)$ . To construct some sequence spaces, many authors used a modulus function, see Ruckle (1973), Maddox (2008), Pehlivan and Fisher (1994), Altin (2009), Raj et al. (2015).

**Definition 2.2.**

A Banach space  $X$  possess Banach-Saks property if any bounded sequence in  $X$  approves a subsequence whose arithmetic mean converges in norm. In a similar vein, a Banach space  $X$  has weak Banach-Saks property if any weakly null sequence in  $X$  admits a subsequence whose arithmetic mean strongly converges in norm.

**Definition 2.3.**

Let  $Y$  be a Banach space. The coefficient  $R(Y)$  was defined by

$$R(Y) = \sup \left( \liminf_{n \rightarrow \infty} \|y_n + y\| \right)$$

and also a Banach space  $Y$  with  $R(Y) < 2$  holds weak fixed point property. [Garcia-Falset (1994)].

**Definition 2.4.**

Let  $L_n$  be the  $n$ th term of a sequence such that  $L_0 = 2, L_1 = 1$  and  $L_n = L_{n-1} + L_{n-2}, n \geq 2$ . The resulting sequence 1,3,4,7,11,18,... is called Lucas sequence.

We can find so many fundamental properties about Lucas sequences in Koshy (2017) and Vajda (1989). Some of them are as follows:

$$\begin{aligned} \sum_{k=1}^n L_k &= L_{n+2} - 3; \quad \sum_{k=1}^n L_{2k-1} = L_{2n-2}; \quad n \geq 1, \\ \sum_{k=1}^n L_k^2 &= L_n L_{n+1} - 2; \quad \sum_{k=1}^{2n} L_k L_{k-1} = L_{2n}^2 - 4; \\ L_{n+1} L_{n-1} - L_n^2 &= (-5)(-1)^n. \end{aligned}$$

It can be easily derived by placing  $L_{n+1}$  in the last equality that

$$L_{n-1}^2 + L_n L_{n-1} - L_n^2 = -5(-1)^n.$$

By using above information, we define the following double band matrix

$$\hat{E} = (\hat{L}_{nk}) = \begin{cases} -\frac{L_n}{L_{n-1}}, & (k = n - 1), \\ \frac{L_{n-1}}{L_n}, & (k = n), \\ 0, & \text{other,} \end{cases} \quad ; \quad n, k \in \mathbb{N} - \{0\}. \tag{3}$$

Karakaş and Karakaş (2017). The inverse of this matrix is

$$\hat{E}^{-1} = \begin{cases} \frac{L_n^2}{L_{k-1}L_k}, & 0 < k \leq n, \\ 0, & \text{other,} \end{cases}$$

and the  $\hat{E}$  – transform of a sequence  $x = (x_n)$  is defined by

$$y_n = \hat{E}_n(x) = \frac{L_{n-1}}{L_n} x_n - \frac{L_n}{L_{n-1}} x_{n-1}; \quad n \geq 1. \quad (4)$$

Later, the above Lucas matrix was generalized and constructed the matrix  $E(r, s) = (L_{nk}(r, s))$  for non-zero real numbers  $r$  and  $s$  as follows:

$$\hat{E}(r, s) = \begin{cases} s \frac{L_n}{L_{n-1}}, & (k = n - 1), \\ r \frac{L_{n-1}}{L_n}, & (k = n), \\ 0, & (0 \leq k < n - 1 \text{ or } k > n), \end{cases}; \quad n, k \in \mathbb{N} - \{0\}.$$

Now, by using the matrix  $\hat{E}(r, s)$ , we introduce the following Lucas difference sequence spaces:

$$\ell(\hat{E}(r, s), \mathfrak{F}, p, u) = \left\{ x \in w: \sum_n \left[ u_n F_n \left( \left| r \frac{L_{n-1}}{L_n} x_n + s \frac{L_n}{L_{n-1}} x_{n-1} \right| \right) \right]^{p_n} < \infty \right\}$$

and

$$\ell_\infty(\hat{E}(r, s), \mathfrak{F}, p, u) = \left\{ x \in w: \sup_n \left[ u_n F_n \left( \left| r \frac{L_{n-1}}{L_n} x_n + s \frac{L_n}{L_{n-1}} x_{n-1} \right| \right) \right]^{p_n} < \infty \right\},$$

where  $\mathfrak{F} = (F_n)$  is a sequence of modulus functions,  $p = (p_n)$  is any bounded sequence of positive real numbers and  $u = (u_n)$  is a sequence of strictly positive real numbers. Also, for the inverse of the matrix  $\hat{E}(r, s)$  and the  $\hat{E}(r, s)$ -transform of the sequence  $x = (x_n)$ , we'll use the following equalities:

$$\hat{E}^{-1}(r, s) = \begin{cases} \frac{1}{r} \left( -\frac{s}{r} \right)^{n-k} \frac{L_n^2}{L_{k-1}L_k}, & 0 < k \leq n, \\ 0, & k > n, \end{cases}$$

and

$$y_n = \hat{E}_n(r, s)(x) = r \frac{L_{n-1}}{L_n} x_n + s \frac{L_n}{L_{n-1}} x_{n-1}; \quad n \geq 1. \quad (5)$$

In addition, we will require the inequality

$$|a_k + b_k|^{p_k} \leq C(|a_k|^{p_k} + |b_k|^{p_k}), \quad (6)$$

where  $C = \max\{1, 2^{H-1}\}$ ,  $H = \sup p_k$  and  $p = (p_k)$  is a sequence of positive real numbers.

### 3. Results and discussion

The proofs of the following two theorems are easy, so we give them without proof.

**Theorem 3.1.**

$\ell(\hat{E}(r, s), \mathfrak{F}, p, u)$  and  $\ell_\infty(\hat{E}(r, s), \mathfrak{F}, p, u)$  are linear spaces over  $\mathbb{C}$ .

**Theorem 3.2.**

$\ell(\hat{E}(r, s), \mathfrak{F}, p, u)$  is a paranormed space for  $M = \max(1, H)$ ,  $H = \sup p_n$  with paranorm

$$\varphi(x) = \sup p_n \left( \sum_n \left[ u_n F_n \left( \left| r \frac{L_{n-1}}{L_n} x_n + s \frac{L_n}{L_{n-1}} x_{n-1} \right| \right) \right]^{p_n} \right)^{1/M}.$$

**Theorem 3.3.**

Let  $\mathfrak{F} = (F_n)$  be a sequence of modulus functions,  $p = (p_n)$  and  $q = (q_n)$  be bounded sequences of positive real numbers. If  $0 \leq p_n \leq q_n < \infty$  for each  $n$ , then  $\ell(\hat{E}(r, s), \mathfrak{F}, p, u) \subseteq \ell(\hat{E}(r, s), \mathfrak{F}, q, u)$ .

**Proof:**

Let us take  $x \in \ell(\hat{E}(r, s), \mathfrak{F}, p, u)$ . So,  $\sum_n \left[ u_n F_n \left( \left| r \frac{L_{n-1}}{L_n} x_n + s \frac{L_n}{L_{n-1}} x_{n-1} \right| \right) \right]^{p_n} < \infty$ . It means that  $\left[ u_n F_n \left( \left| r \frac{L_{n-1}}{L_n} x_n + s \frac{L_n}{L_{n-1}} x_{n-1} \right| \right) \right]^{p_n} \leq 1$  for  $n \geq n_0$ , that is, sufficiently large values of  $n$ . Hence, we obtain

$$\begin{aligned} & \sum_{n \geq n_0} \left[ u_n F_n \left( \left| r \frac{L_{n-1}}{L_n} x_n + s \frac{L_n}{L_{n-1}} x_{n-1} \right| \right) \right]^{q_n} \\ & \leq \sum_{n \geq n_0} \left[ u_n F_n \left( \left| r \frac{L_{n-1}}{L_n} x_n + s \frac{L_n}{L_{n-1}} x_{n-1} \right| \right) \right]^{p_n} < \infty, \end{aligned}$$

which implies that  $x \in \ell(\hat{E}(r, s), \mathfrak{F}, q, u)$  because of  $p_n \leq q_n$  and  $F_n$  is increasing.

**Theorem 3.4.**

If  $\mathfrak{F} = (F_n)$  be a sequence of modulus functions and  $\alpha = \lim_{q \rightarrow \infty} \frac{F_n(q)}{q} > 0$ , then  $\ell(\hat{E}(r, s), \mathfrak{F}, p, u) \subseteq \ell(\hat{E}(r, s), p, u)$ .

**Proof:**

The definition of  $\alpha$  gives us  $F_n(q) \geq \alpha(q)$ , for all  $q > 0$ . Herefrom, we see that  $\frac{1}{\alpha} F_n(q) \geq q$ , for all  $q > 0$ . Now, for  $x \in \ell(\hat{E}(r, s), \mathfrak{F}, p, u)$  we have

$$\sum_n \left[ u_n \left( \left| r \frac{L_{n-1}}{L_n} x_n + s \frac{L_n}{L_{n-1}} x_{n-1} \right| \right) \right]^{p_n} \leq \frac{1}{\alpha} \sum_n \left[ u_n F_n \left( \left| r \frac{L_{n-1}}{L_n} x_n + s \frac{L_n}{L_{n-1}} x_{n-1} \right| \right) \right]^{p_n},$$

which shows that  $x \in \ell(\hat{E}(r, s), p, u)$ .

**Theorem 3.5.**

The inclusion  $\ell(\hat{E}(r, s), \mathfrak{F}', p, u) \cap \ell(\hat{E}(r, s), \mathfrak{F}'', p, u) \subseteq \ell(\hat{E}(r, s), \mathfrak{F}' + \mathfrak{F}'', p, u)$  holds for sequences of modulus functions  $\mathfrak{F}' = (F'_n)$  and  $\mathfrak{F}'' = (F''_n)$ .

**Proof:**

Let  $x \in \ell(\hat{E}(r, s), \mathfrak{F}', p, u) \cap \ell(\hat{E}(r, s), \mathfrak{F}'', p, u)$ . Then, it can be easily seen that

$$\sum_n \left[ u_n F'_n \left( \left| r \frac{L_{n-1}}{L_n} x_n + s \frac{L_n}{L_{n-1}} x_{n-1} \right| \right) \right]^{p_n} < \infty;$$

and

$$\sum_n \left[ u_n F''_n \left( \left| r \frac{L_{n-1}}{L_n} x_n + s \frac{L_n}{L_{n-1}} x_{n-1} \right| \right) \right]^{p_n} < \infty.$$

From here, we have

$$\begin{aligned} & \sum_n \left[ u_n (F'_n + F''_n) \left( \left| r \frac{L_{n-1}}{L_n} x_n + s \frac{L_n}{L_{n-1}} x_{n-1} \right| \right) \right]^{p_n} \\ & \leq M \sum_n \left[ u_n F'_n \left( \left| r \frac{L_{n-1}}{L_n} x_n + s \frac{L_n}{L_{n-1}} x_{n-1} \right| \right) \right]^{p_n} \\ & \quad + M \sum_n \left[ u_n F''_n \left( \left| r \frac{L_{n-1}}{L_n} x_n + s \frac{L_n}{L_{n-1}} x_{n-1} \right| \right) \right]^{p_n}, \end{aligned}$$

which means that  $x \in \ell(\hat{E}(r, s), \mathfrak{F}' + \mathfrak{F}'', p, u)$ .

**Theorem 3.6.**  $\ell(\hat{E}(r, s), \mathfrak{F}', p, u) \subseteq \ell(\hat{E}(r, s), \mathfrak{F} \circ \mathfrak{F}', p, u)$

**Proof:**

Let us take  $\varepsilon > 0, 0 < \delta < 1$  such that  $F_n(q) < \varepsilon$ , for  $0 \leq q \leq \delta$ . Choose  $y_n = u_n F'_n \left( \left| r \frac{L_{n-1}}{L_n} x_n + s \frac{L_n}{L_{n-1}} x_{n-1} \right| \right)$  and consider

$$\sum_n [F_n(y_n)]^{p_n} = \sum_{y_n \leq \delta} [F_n(y_n)]^{p_n} + \sum_{y_n > \delta} [F_n(y_n)]^{p_n}.$$

In view of the continuity of  $F_n$ , we have

$$\sum_{y_n \leq \delta} [F_n(y_n)]^{p_n} < \varepsilon^H, \quad (7)$$

and  $y_n > \delta \Rightarrow y_n < \frac{y_n}{\delta} \leq 1 + \frac{y_n}{\delta}$ . For  $y_n > \delta$ , we obtain by the definition that

$$F_n(y_n) < 2F_n(1) \frac{y_n}{\delta}$$

and so

$$\sum_{y_n > \delta} [F_n(y_n)]^{p_n} < \max\{1, (2F_n(1)\delta^{-1})^H\} \sum_n [y_n]^{p_n}. \tag{8}$$

Equations (7) and (8) give the fact that  $\ell(\hat{E}(r, s), \mathfrak{F}', p, u) \subseteq \ell(\hat{E}(r, s), \mathfrak{F} \circ \mathfrak{F}', p, u)$ .

**Theorem 3.7.**

$\ell(\hat{E}(r, s), \mathfrak{F}, p, u)$  and  $\ell_\infty(\hat{E}(r, s), \mathfrak{F}, p, u)$  are normed spaces with

$$\|x\|_{\ell(\hat{E}(r,s),\mathfrak{F},p,u)} = \left( \sum_n [u_k F_k(|\hat{E}_n(r, s)(x)|)]^{p_k} \right)^{1/M},$$

and

$$\|x\|_{\ell_\infty(\hat{E}(r,s),\mathfrak{F},p,u)} = \sup_n [u_k F_k(|\hat{E}_n(r, s)(x)|)]^{p_k},$$

for  $1 \leq p_k \leq H \leq \infty$  for all  $k \in \mathbb{N}$ .

**Proof:**

It can be proved by standard technic in Raj et al. (2015).

$\ell(\hat{E}(r, s), \mathfrak{F}, p, u)$  and  $\ell_\infty(\hat{E}(r, s), \mathfrak{F}, p, u)$  are sequence spaces of non-absolute type. Indeed,

$$\|x\|_{\ell(\hat{E}(r,s),\mathfrak{F},p,u)} \neq \| |x| \|_{\ell(\hat{E}(r,s),\mathfrak{F},p,u)} \text{ and } \|x\|_{\ell_\infty(\hat{E}(r,s),\mathfrak{F},p,u)} \neq \| |x| \|_{\ell_\infty(\hat{E}(r,s),\mathfrak{F},p,u)};$$

which means that the absolute property is not valid on the spaces  $\ell(\hat{E}(r, s), \mathfrak{F}, p, u)$  and  $\ell_\infty(\hat{E}(r, s), \mathfrak{F}, p, u)$  for at least one sequence  $x = (x_k)$ .

**Theorem 3.8.**

The sequence space  $\ell(\hat{E}(r, s), \mathfrak{F}, p, u)$  of non-absolute type is linearly isomorphic to the space  $\ell_p$  for  $1 \leq p_k \leq H < \infty$  for all  $k \in \mathbb{N}$ , that is,  $\ell(\hat{E}(r, s), \mathfrak{F}, p, u) \cong \ell_p$ .

**Proof:**

Let us take into consideration the transformation  $Z: \ell(\hat{E}(r, s), \mathfrak{F}, p, u) \rightarrow \ell_p$  defined by  $x \rightarrow y = Zx$ , with equality (5). Then, for  $x \in \ell(\hat{E}(r, s), \mathfrak{F}, p, u)$ , we have  $Zx = y = \hat{E}(r, s)x \in \ell_p$ . So, it is obvious that  $Z$  is linear. Also, it can be easily shown that  $Zx = 0 \Rightarrow x = 0$  and thus  $Z$  is injective. Now, let us consider  $y = (y_k) \in \ell_p$  and identify the sequence  $x = (x_k)$  for  $1 \leq p_k \leq H < \infty$  for all  $k \in \mathbb{N}$  as follows:

$$x_k = \frac{1}{r} \sum_{j=1}^k \left(-\frac{s}{r}\right)^{k-j} \frac{L_k^2}{L_{j-1}L_j} y_j.$$

Hence, we obtain for  $1 \leq p_k \leq H < \infty$  for all  $k \in \mathbb{N}$  and  $p = \infty$ ,



$$\begin{aligned}
\|x\|_{\ell(\hat{E}(r,s),\mathfrak{F},p,u)} &= \left( \sum_k \left[ u_k F_k \left( \left| r \frac{L_{k-1}}{L_k} x_k + s \frac{L_k}{L_{k-1}} x_{k-1} \right| \right) \right]^{p_k} \right)^{1/M} \\
&= \left( \sum_k \left[ u_k F_k \left( \left| r \frac{L_{k-1}}{L_k} \frac{1}{r} \sum_{j=1}^k \left( -\frac{s}{r} \right)^{k-j} \frac{L_k^2}{L_{j-1} L_j} y_j \right. \right. \right. \\
&\quad \left. \left. \left. + s \frac{L_k}{L_{k-1}} \frac{1}{r} \sum_{j=1}^{k-1} \left( -\frac{s}{r} \right)^{k-j} \frac{L_k^2}{L_{j-1} L_j} y_j \right| \right) \right]^{p_k} \right)^{1/M} \\
&= \|y\|_{\ell_p} < \infty,
\end{aligned}$$

and

$$\|x\|_{\ell_\infty(\hat{E}(r,s),\mathfrak{F},p,u)} = \sup_k [u_k F_k(|\hat{E}_k(r,s)(x)|)]^{p_k} = \|y\|_{\ell_\infty} < \infty.$$

This gives the fact that  $Z$  is linear bijection and so the proof is completed. Now, we'll investigate the geometric structure of the space  $\ell(\hat{E}(r,s), \mathfrak{F}, p, u)$ , that is, we'll examine whether the space  $\ell(\hat{E}(r,s), \mathfrak{F}, p, u)$  has Banach-Saks property of type  $p$  and the weak fixed point property or not.

### Theorem 3.9.

Let  $1 < p_k \leq H < \infty$  for all  $k \in \mathbb{N}$ . The space  $\ell(\hat{E}(r,s), \mathfrak{F}, p, u)$  has Banach-Saks property of type  $p$ .

#### *Proof:*

It can be demonstrated by standard technic which can be seen in Et et al. (2014).

Since the space  $\ell(\hat{E}(r,s), \mathfrak{F}, p, u)$  is linearly isomorphic to space  $\ell_p$ , we have  $R(\ell(\hat{E}(r,s), \mathfrak{F}, p, u)) = R(\ell_p) = 2^{1/p}$ . In view of  $R(\ell(\hat{E}(r,s), \mathfrak{F}, p, u)) < 2$ , we can give the following theorem with the help of definition 2.3.

### Theorem 3.10.

The space  $\ell(\hat{E}(r,s), \mathfrak{F}, p, u)$  has weak fixed point property in the case  $1 < p_k \leq H < \infty$  for all  $k \in \mathbb{N}$ .

## 4. Conclusion

Geometrical properties of Banach spaces have been studied by many authors. In recent years, one of the interesting topics is to examine topological and geometrical properties of difference sequence spaces defined by Fibonacci and Lucas numbers. In this paper, we apply the domain of infinite triangular matrix established by Lucas numbers and the modulus function to space  $\ell_p$ . Then, we investigate topological and geometric structure of the obtained space. Since some of the geometric properties of Banach spaces play an important role in the fixed point theory, our results are interesting. However, the Lucas numbers and its properties can be considered in different fields of summability theory such as statistical convergence and its applications.

## REFERENCES

- Ahmad, Z. (1987). Köthe-Toeplitz Duals of Some New Sequence Spaces and Their Matrix Maps, Publications de l'Institut Mathématique, Vol. 42, No. 62, pp. 57-61.
- Altay, Bilal and Başar, Feyzi (2005). On Some Euler Sequence Spaces of Nonabsolute Type, Ukrainian Mathematical Journal, Vol. 57, No. 1, pp. 1-17.
- Altay, Bilal and Başar, Feyzi (2007). The Fine Spectrum and The Matrix Domain of The Difference Operator  $\Delta$  on The Sequence Space  $\ell_p$ , ( $0 < p < 1$ ), Communications in Mathematical Analysis, Vol. 2, No. 2, pp. 1-11.
- Altin, Yavuz (2009). Properties of Some Sets of Sequences Defined by a Modulus Function, Acta Mathematica Scientia, Vol. 29, No. 2, pp. 427-434.
- Başar, F. (2011). *Summability Theory and Its Applications*, Bentham e books, Büyükçekmece, İstanbul.
- Başar, Feyzi and Altay, Bilal (2003). On the Space of Sequences of p-Bounded Variation and Related Matrix Mappings, Ukrainian Mathematical Journal, Vol. 55, No. 1, pp. 136-147.
- Başar, Feyzi, Altay, Bilal and Mursaleen, Muhammed (2008). Some Generalizations of the Space  $bvp$  of p-bounded Variation Sequences, Nonlinear Analysis: Theory, Methods & Applications, Vol. 68, No. 2, pp. 273-287.
- Colak, Rifat and Et, Mikail (2005). On Some Difference Sequence Sets and Their Topological Properties, Bulletin of the Malaysian Mathematical Sciences Society, Vol. 28, No. 2, pp. 125-130.
- Çolak, Rifat, Et, Mikail and Malkowsky, Eberhard (2004). *Some Topics of Sequence Spaces*, Lecture Notes in Mathematics, Firat University Press, Elazig, Turkey.
- Debnath, Shyamal, Datta, Amar J. and Saha, Subrata (2015). Regular Matrix of Interval Numbers Based on Fibonacci Numbers, Afrika Matematika, Vol. 26, No. 7-8, pp. 1379-1385.
- Debnath, Shyamal and Saha, Subrata (2014). Some Newly Defined Sequence Spaces Using Regular Matrix of Fibonacci Numbers, Afyon Kocatepe University Journal of Science & Engineering, Vol. 14, No. 1, pp. 1-3.
- Et, Mikail (1993). On Some Difference Sequence Spaces, Turkish Journal of Mathematics, Vol. 17, pp. 18-24.
- Et, Mikail and Basarir, Metin (1997). On Some New Generalized Difference Sequence Spaces, Periodica Mathematica Hungarica, Vol. 35, No. 3, pp. 169-175.
- Et, Mikail and Çolak, Rifat (1995). On Some Generalized Difference Sequence Spaces, Soochow Journal of Mathematics, Vol. 21, No. 4, pp. 377-386.
- Et, Mikail, Karakas, Murat and Karakaya, Vatan (2014). Some Geometric Properties of a New Difference Sequence Space Defined by De La Vallee-Poussin Mean, Applied Mathematics and Computation, Vol. 234, pp. 237-244.
- Garcia-Falset, Jesus (1994). Stability and Fixed Points for Nonexpansive Mappings, Houston Journal of Mathematics, Vol. 20, No. 3, pp. 495-506.
- Kara, Emrah Evren (2013). Some Topological and Geometrical Properties of New Banach Sequence Spaces, Journal of Inequalities and Applications, Vol. 2013, No. 38, pp. 15.
- Kara, Emrah Evren and Basarir, Metin (2012). An application of Fibonacci Numbers into Infinite Toeplitz Matrices, Caspian Journal of Mathematical Sciences, Vol. 1, No. 1, pp. 43-47.
- Karakas, Murat (2015). A New Regular Matrix Defined by Fibonacci Numbers and Its Applications, Bitlis Eren University Journal of Science, Vol. 4, No. 2, pp. 205-210.

- Karakaş, Murat and Metin Karakaş, Ayşe (2017). New Banach Sequence Spaces That Is Defined By The Aid Of Lucas Numbers, *Iğdır University Journal of Institute of Science and Technology*, Vol. 7, No. 4, pp. 103-111.
- Kirisci, Murat and Başar, Feyzi (2010). Some New Sequence Spaces Derived by the Domain of Generalized Difference Matrix, *Computers & Mathematics with Applications*, Vol. 60, No. 5, pp. 1299-1309.
- Kizmaz, Hüsni (1981). On Certain Sequence Spaces, *Canadian Mathematical Bulletin*, Vol. 25, No. 2, pp. 169-176.
- Koshy, T. (2017). *Fibonacci and Lucas Numbers with Applications*, Wiley & Sons, Hoboken, New Jersey.
- Maddox, Ivor J. (2008). Sequence Spaces Defined by a Modulus, *Mathematical Proceedings of the Cambridge Philosophical Society*, Vol. 100, No. 1, pp. 161-166.
- Malkowsky, Eberhard (1989). Absolute and Ordinary Köthe-Toeplitz Duals of Some Sets of Sequences and Matrix Transformations, *Publications De L'Institut Mathematique*, Vol. 46, No. 60, pp. 97-103.
- Mursaleen, Muhammed (1996). Generalized Spaces of Difference Sequences, *Journal of Mathematical Analysis and Applications*, Vol. 203, No. 3, pp. 738-745.
- Mursaleen, Muhammed and Noman, Abdullah K. (2010). On Some New Difference Sequence Spaces of Non-Absolute Type, *Mathematical and Computer Modelling*, Vol. 52, No. 3-4, pp. 603-617.
- Mursaleen, Muhammed and Noman, Abdullah K. (2011). On Some New Sequence Spaces of Non-Absolute Type Related to the Spaces  $l(p)$  and  $l(\infty)$  II, *Mathematical Communications*, Vol. 16, No. 2, pp. 383-398.
- Nakano, Hidegoro (1953). Concave Modulars, *Journal of the Mathematical Society of Japan*, Vol. 5, No. 1, pp. 29-49.
- Pehlivan, Serpil and Fisher, Brian (1994). On Some Sequence Spaces, *Indian Journal of Pure & Applied Mathematics*, Vol. 25, No. 10, pp. 1067-1071.
- Raj, Kuldip, Pandoh, Suruchi and Jamwal, Seema (2015). Fibonacci Difference Sequence Spaces for Modulus Functions, *Le Matematiche*, Vol. 70, No. 1, pp. 137-156.
- Ruckle, William H. (1973). FK-Spaces in which The Sequence of Coordinate Vectors is Bounded, *Canadian Journal of Mathematics*, Vol. 25, pp. 973-978.
- Tripathy, Binod Chandra (2003). On a Class of Difference Sequences Related to the  $p$ -normed Space  $l(p)$ , *Demonstratio Mathematica*, Vol. 36, No. 4, pp. 867-872.
- Vajda, S. (1989). *Fibonacci and Lucas Numbers, and the Golden Section: Theory and Applications*, Dover Publications, Inc., Mineola, New York.