



12-2020

## An Efficient Algorithm for Numerical Inversion of System of Generalized Abel Integral Equations

Sandeep Dixit

*University of Petroleum and Energy Studies Dehradun*

Follow this and additional works at: <https://digitalcommons.pvamu.edu/aam>



Part of the [Analysis Commons](#), and the [Numerical Analysis and Computation Commons](#)

### Recommended Citation

Dixit, Sandeep (2020). An Efficient Algorithm for Numerical Inversion of System of Generalized Abel Integral Equations, *Applications and Applied Mathematics: An International Journal (AAM)*, Vol. 15, Iss. 2, Article 33.

Available at: <https://digitalcommons.pvamu.edu/aam/vol15/iss2/33>

This Article is brought to you for free and open access by Digital Commons @PVAMU. It has been accepted for inclusion in *Applications and Applied Mathematics: An International Journal (AAM)* by an authorized editor of Digital Commons @PVAMU. For more information, please contact [hvkoshy@pvamu.edu](mailto:hvkoshy@pvamu.edu).



## An Efficient Algorithm for Numerical Inversion of System of Generalized Abel Integral Equations

**Sandeep Dixit**

Department of Mathematics  
University of Petroleum and Energy Studies  
Dehradun, India  
[sandydixit27@gmail.com](mailto:sandydixit27@gmail.com)

Received: June 26, 2020; Accepted: October 11, 2020

### Abstract

In this article a direct method is introduced, which is based on orthonormal Bernstein polynomials, to present an efficient and stable algorithm for numerical inversion of the system of singular integral equations of Abel type. The appropriateness of earlier numerical inversion methods was restricted to the one portion of singular integral equations of Abel type. The proposed method is absolutely accurate, and numerical illustrations are given to show the convergence and utilization of the suggested method and comparisons are made with some other existing numerical solution.

**Keywords:** Bernstein polynomials; Abel inversion; Operational matrix; System of generalized Abel integral equations

**MSC 2010 No.:** 26A33, 41A10, 45A05

### 1. Introduction

Many attention grabbing real life problems of physics and mechanics result in an integral equation in which the kernel of the integral equation is of convolution type. A similar type integral equation with singular kernel named singular integral equation of Abel type which emerges in many branches of science and engineering. Commonly, physical quantities available to assessment are actually often related to physically essential but experimentally unattainable ones by integral equations of generalized Abel type.

There are certain numerical approximation methods for solving Volterra integral equations and integral equations of Abel kernel and generalized Abel kernel like, homotopy analysis transform technique, Galerkin method, Taylor series method and Taylor polynomials Methods, (Kumar et al. (2014), Burton (2005), Maleknejad and Aghazadeh (2005) and Yalsinbas (2002)). In recent past a number of methods have been contemplated based on Chebyshev polynomials (Maleknejad et al. (2007)), Homotopy Perturbation method (Kumar and Singh O. P. (2010), Kumar et al. (2011)), Laplace transform method (Kumar et al. (2015)), Bernstein polynomials (Pandey et al. (2009)), Power Series method (Tahmasbi and Fard (2008)), Laguerre polynomial approximation (Singha and Nahak (2019)) and almost Bernstein operational matrix method (Dixit et al. (2011), Dixit et al. (2012)) to find the numerical inversion of integral equation with Abel type kernel.

Recently Mandal and Chakrabarti (2016) achieved a numerical inversion of the system of generalized singular integral equations of Abel type kernel using order 8 Bernstein polynomials but the proposed method is less accurate when the coefficients to equations are polynomial functions compared to the case when the coefficients are constants. Irfan et al. (2014) used the concept Bernstein operational matrices to solve integro-differential equation. Recently Kumar et al. (2020) used Bernstein wavelets to solve fractional predator – prey dynamical system and fractional SIR epidemic model respectively. This article is devoted to obtaining a class of almost operational matrices based on Bernstein type of orthogonal polynomials to find the numerical inversion of system of integral equation with generalized Abel type.

I have organized my paper as follows. In section 2, system of Abel integral equation with variable coefficients is given. Section 3 contains orthonormal Bernstein polynomials. In section 4 function approximation given to apply the technique. Section 5 contains method of solution to find inversion. Section 6 contains numerical examples to illustrate the stability and the accuracy of our algorithm

## 2. System of Abel Integral Equations

A system of singular integral equations of Abel type was studied by Lowengrub (1976) and Walton (1979). As told in J Walton (1979), a few mixed boundary value based problems arising in the classical elasticity theory diminish to the research problem of determining functions  $\mu_1$  and  $\mu_2$  gratifying generalized singular integral equations of the form,

$$q_{11}(y) \int_0^y \frac{\mu_1(z) dz}{(y^d - z^d)^b} + q_{12}(y) \int_y^1 \frac{\mu_2(z) dz}{(z^d - y^d)^b} = h_1(y),$$

$$q_{21}(y) \int_0^y \frac{\mu_1(z) dz}{(z^d - y^d)^b} + q_{22}(y) \int_y^1 \frac{\mu_2(z) dz}{(y^d - z^d)^b} = h_2(y), \quad (1)$$

where  $y \in (0,1)$ ,  $0 < b < 1$ ,  $q_{ij}(y)$  are continuous on  $[0,1]$ ,  $h_1$  and  $h_2$  are the known functions. Since only  $d=1,2$  occur in physics related problems, we impede our consideration to these values only.

Mandal *et al.* (1996), solved this physical problem analytically for the particular case  $d = 1, b = 1/2, q_{ij}(y) = 1$ , using the concept of fractional calculus. Recently Mandal and Chakrabarti (2016) achieved a numerical inversion of the given system (1) using order 8 Bernstein polynomials

$$\int_0^y \frac{\mu(z)dz}{(y-z)^{1/2}} = h(y) \quad \text{and} \quad \int_y^1 \frac{\mu(z)dz}{(z-y)^{1/2}} = h(y). \quad (2)$$

But the method proposed by Mandal and Chakrabarti (2016) is less accurate when the coefficients  $q_{ij}(y)$  are polynomial functions compared to the case when the coefficients are constants.

### 3. The Orthonormal Polynomials

The Bernstein polynomials (B polynomials) characterized over the interval  $[0,1]$  are given as;

$$B_{i,\alpha}(z) = \binom{\alpha}{i} z^i (1-z)^{\alpha-i}, \quad \forall i = 0, 1, 2, \dots, \alpha.$$

By using orthonormalization process given by Gram- Schmidt method on  $B_{i,\alpha}$ , we attain a family of orthonormal polynomials from B polynomials. We denote these polynomials by  $b_{0\alpha}, b_{1\alpha}, \dots, b_{\alpha\alpha}$ . After all, for  $\alpha = 5$ , the six orthonormal B polynomials were enumerated and given:

$$b_{05}(z) = \sqrt{11}(1-z)^5,$$

$$b_{15}(z) = 3(1-z)^4 [11z - 1],$$

$$b_{25}(z) = \sqrt{7}(1-z)^3 [55z^2 - 20z + 1],$$

$$b_{35}(z) = \frac{28}{\sqrt{5}} \left[ 10(1-z)^2 z^3 - 15(1-z)^3 z^2 + \frac{30}{7}(1-z)^4 z - \frac{5}{28}(1-z)^5 \right],$$

$$b_{45}(z) = 7\sqrt{3} \left[ 5(1-z)z^4 - 20(1-z)^2 z^3 + 18(1-z)^3 z^2 - 4(1-z)^4 z + \frac{1}{7}(1-z)^5 \right],$$

$$b_{55}(z) = 6 \left[ z^5 - \frac{25}{2}(1-z)z^4 + \frac{100}{3}(1-z)^2 z^3 - 25(1-z)^3 z^2 + 5(1-z)^4 z - \frac{1}{6}(1-z)^5 \right].$$

#### 4. Function Approximation

Any given function  $h \in L^2[0,1]$  be allowed to express as

$$h(z) = \lim_{\alpha \rightarrow \infty} \sum_{i=0}^{\alpha} c_{i\alpha} b_{i\alpha}(z), \quad (3)$$

where  $c_{i\alpha} = \langle h, b_{i\alpha} \rangle$  and  $\langle \cdot, \cdot \rangle$  is the notation for inner product on  $L^2[0,1]$ .

If the given series (3) is truncated at  $n = \alpha$ , then we get

$$h \cong \sum_{i=0}^{\alpha} c_{i\alpha} b_{i\alpha} = C^T B(z), \quad (4)$$

where  $C$  and  $B(z)$  are  $(\alpha + 1) \times 1$  matrices given by

$$C = [c_{0\alpha}, c_{1\alpha}, \dots, c_{\alpha\alpha}]^T, \quad (5)$$

and

$$B(z) = [b_{0\alpha}(z), b_{1\alpha}(z), \dots, b_{\alpha\alpha}(z)]^T. \quad (6)$$

#### 5. Method of Solution

Under this heading we solve integral equation of generalized Abel kernel by orthonormal B polynomials. Using Equation (4), we approximate  $\mu_1(z), \mu_2(z), h_1(y)$  and  $h_2(y)$  as

$$\tilde{\mu}_1(z) = C_1^T B(z), \tilde{\mu}_2(z) = C_2^T B(z), \tilde{h}_1(y) = H_1^T B(y), \tilde{h}_2(y) = H_2^T B(y). \quad (7)$$

The matrices  $H_1, H_2$  are known and the matrices  $C_1, C_2$  are to be found. From Equations (1) and (7), we have

$$\begin{aligned} q_{11}(y) \int_0^y \frac{C_1^T B(z) dz}{(y^d - z^d)^b} + q_{12}(y) \int_y^1 \frac{C_2^T B(z) dz}{(z^d - y^d)^b} &= H_1^T B(y), \\ q_{21}(y) \int_0^y \frac{C_1^T B(z) dz}{(z^d - y^d)^b} + q_{22}(y) \int_y^1 \frac{C_2^T B(z) dz}{(y^d - z^d)^b} &= H_2^T B(y), \end{aligned} \quad (8)$$

Using Equation (6) and the following formulae,

$$\int_0^y \frac{z^n dz}{(y^d - z^d)^b} = \frac{\pi y^n (d/y)^{b-1} (y^{d-1}d)^{-b} \operatorname{cosec}(\pi b) \Gamma(n+1/d)}{\Gamma(b) \Gamma(n+d-db+1/d)}, \quad (9)$$

$$\int_y^1 \frac{z^n dz}{(z^d - y^d)^b} = \left[ \frac{(y^{-d})^{\frac{n+1}{d}} (y^d)^{-b} [\Gamma(1-b) \Gamma(b-n+1/d)]}{d \Gamma(-n+d-1/d)} - \frac{(y^{-d})^{\frac{n+1}{d}} (y^d)^{-b} [B_{y^d}(b-n+1/d, 1-b) \Gamma(-n+d-1/d)]}{d \Gamma(-n+d-1/d)} \right], \quad (10)$$

where  $B_y(p, q) = \int_0^y u^{p-1} (1-u)^{q-1} du$  is incomplete beta function. Thus, we obtain

$$\int_0^y \frac{B(z)}{(y^d - z^d)^b} dz = U B(y), \quad \int_y^1 \frac{B(z)}{(z^d - y^d)^b} dz = V B(y), \quad (11)$$

where  $U$  and  $V$  are  $(\alpha+1) \times (\alpha+1)$  order matrices, which we call as AOMI for singular integral equation with generalized kernel.

Replacing the values of Equation (11) in Equation (8), we have

$$C_1^T = - \left[ \frac{H_2^T q_{12}(y)V - H_1^T q_{22}(y)U}{q_{11}(y)q_{22}(y)U^2 - q_{21}(y)q_{12}(y)V^2} \right], \quad (12)$$

$$C_2^T = - \left[ \frac{H_1^T q_{21}(y)V - H_2^T q_{11}(y)U}{q_{11}(y)q_{22}(y)U^2 - q_{21}(y)q_{12}(y)V^2} \right].$$

Hence, the approximate solutions  $\tilde{\mu}_1(z), \tilde{\mu}_2(z)$  for singular integral equation (1) are attained by placing the value of  $C_1^T, C_2^T$  from Equation (12) in Equation (1).

## 6. Illustrative Examples

Under this heading, we examine the utilization of our suggested method and examine its accuracy and stability by implementing it on test functions with known analytical result of Abel inversion. For it is always required to check the demeanor of a numerical method using given simulated data, for which the exact solutions are known and thus creating a resemblance between the inverted solutions and the theoretical data is possible. We have chosen two test functions having various shapes for this objective.

The five numerical illustrations are given with added noise terms and without added noise terms to understand the efficiency as well as stability of our given algorithm. Note that in all five

illustrated results to follow, the series in Equation (4) is truncated at level  $\alpha = 5$  and hence the AOMI in Equation (11) are of order  $6 \times 6$ .

The accuracy of given method is established by considering the parameters of absolute error  $\Delta\mu_j(z_i)$  and root mean square error (RMS)  $\sigma^j$  also known as average deviation. Both of them are calculated using the equations given below:

$$\begin{aligned} E_j(z_i) &= |\mu_j(z_i) - \tilde{\mu}_j(z_i)|, \quad \sigma^j_N = \left\{ \frac{1}{N} \sum_{i=1}^N [\mu_j(z_i) - \tilde{\mu}_j(z_i)]^2 \right\}^{1/2} \\ &= \left\{ \frac{1}{N} \sum_{i=1}^N \Delta\mu_j^2(z_i) \right\}^{1/2} = \|\Delta\mu_j\|_2, \end{aligned} \quad (13)$$

where  $\mu_j(z_i)$  represent the exact solution and  $\tilde{\mu}_j(z_i)$  represent approximate numerical result calculated at discrete point  $z_i$  corresponding to the point,  $j=1,2$ . For computational accessibility  $\sigma^j$  is stand for  $\sigma^j_N$ , the discrete  $l^2$ -norm of the absolute error  $\Delta\mu_j$  represented by  $\|\Delta\mu_j\|_2$ .

Note that the root mean square error results of  $\sigma^j_N$  is found by taking  $N=1000, 500$  in Equation (13). In all five numerical examples, the exact profile is represented by  $h_j(y)$  and the profiles with added noises are represented by  $h_j^\delta(y)$ , where  $h_j^\delta(y)$  is obtained by adding a noise term  $\delta$  to  $h_j(y)$  so that  $h_j^\delta(y_i) = h_j(y_i) + \delta\theta_i$ , where  $y_i = ih$ ,  $i=1, \dots, N$ ,  $Nh=1$  and  $\theta_i$  is the discrete random variable uniformly distributed with values in the interval  $[-1, 1]$  such that

$$\text{Max}_{1 \leq i \leq N} |h_{ji}^\delta(y) - h_{ji}(y)| \leq \delta.$$

The given five numerical illustrations are determined without noise and with noise to learn the efficiency and stability of our given algorithm by deciding two distinct values of the noises  $\delta_k$  as  $\delta_0 = 0$ ,  $\delta_1 = \sigma^j_N$ .

In all five numerical illustrations to follow, we measure the corresponding errors  $E_j(z)$  and express them as  $E1(z), E2(z)$  respectively. Then small perturbation (noise term)  $\delta_1$  is added in the known forcing term  $h_j(y)$  and for  $N=1000, 500$  the errors associated to  $\mu_1$  and  $\mu_2$  are labeled by  $E3(z), E4(z)$  and  $E5(z), E6(z)$ , respectively. For correct scaling some  $E_j(z)$  are multiplied by 10. Also, we listed the approximate and exact values for  $z = 0.0, 0.2, \dots, 0.8, 1.0$ .

In the first two numerical examples,  $d=1$  whereas we have taken  $d=2$  in the last three examples.

**Example 1:**

In the first example, take

$$q_{11}(y) = 1, q_{12}(y) = 1/4, q_{21}(y) = 1/2, q_{22}(y) = 3, b = 1/3$$

with

$$h_1(y) = \frac{243}{1760}(1-y)^{2/3}y^3 + \frac{27}{40}y^{8/3} + \frac{459}{1760}y^2(1-y)^{2/3} + \frac{333}{1760}y(1-y)^{2/3} + \frac{57}{352}(1-y)^{2/3},$$

and

$$h_2(y) = \frac{81}{440}(9y+11)(y)^{8/3} + \frac{3}{80}(1-y)^{2/3}(9y^2+6y+5).$$

This generalized system has the solution

$$\mu_1(z) = z^2 \text{ and } \mu_2(z) = z^2 + z^3.$$

The corresponding values of  $C_1^T, C_2^T, \sigma_{1000}^1, \sigma_{500}^1, \sigma_{1000}^2$  and  $\sigma_{500}^2$  are given as

$$C_1^T = [0.026317, 0.130948, 0.304484, 0.457853, 0.49879, 0.333334],$$

$$C_2^T = [0.019736, 0.089282, 0.185845, 0.252881, 0.255692, 0.166667],$$


$$\sigma_{1000}^1 = 1.8 \times 10^{-5}, \quad \sigma_{500}^1 = 1.804 \times 10^{-5},$$

$$\sigma_{1000}^2 = 1.846 \times 10^{-5}, \quad \sigma_{500}^2 = 1.851 \times 10^{-5}.$$


**Table 1.** Approximate and exact values of  $\mu$  for Example 1

$Z$	0.0	0.2	0.4	0.6	0.8	1.0
$\tilde{\mu}_1(z)$	0.000053	0.039999	0.160017	0.360003	0.640000	1.00003
$\mu_1(z)$	0.0	0.040000	0.16000	0.360000	0.640000	1.000000
$\mu_2(z)$	0.000061	0.047969	0.224018	0.576003	1.151998	2.000004
$\tilde{\mu}_2(z)$	0.0	0.048000	0.224000	0.576000	1.152000	2.00000





**Figure 1.** Absolute errors with added noise  $\delta_1(=\sigma_N)$  for  $N = 1000$



**Figure 2.** Absolute errors with added noise  $\delta_1(=\sigma_N)$  for  $N = 500$

**Example 2.**

In second example, take  $q_{11}(y) = y^2 + 1$ ,  $q_{12}(y) = (y + 1) / 4$ ,  $q_{21}(y) = y^2 / 2$ ,  $q_{22}(y) = y$ ,  $b = 1 / 2$  and

$$h_1(y) = \left[ 1/4(y + 1)(4\sqrt{y + 1} \sin^{-1}(\sqrt{1 - y} / \sqrt{2}) + \sqrt{1 - y}(-4 + \log 4)) + e^y \sqrt{\pi} \operatorname{erf}(\sqrt{y})(y^2 + 1) \right],$$

$$h_2(y) = 1/2 y \left( 8\sqrt{y + 1} \sinh^{-1}(\sqrt{y}) + \frac{e^y y \sqrt{\pi} \sqrt{1 - y} \operatorname{erf}(\sqrt{y - 1})}{\sqrt{y - 1}} - 8\sqrt{y} \right),$$

where  $\operatorname{erf}(y)$  is error function, defined by  $\operatorname{erf}(y) = 2/\sqrt{\pi} \int_0^y e^{-t^2} dt$ . This has the exact solution  $\mu_1(z) = e^z$  and  $\mu_2(z) = \log(z + 1)$ .

The corresponding values of  $C_1^T, C_2^T, \sigma_{1000}^1, \sigma_{500}^1, \sigma_{1000}^2$  and  $\sigma_{500}^2$  are given as

$$C_1^T = [0.030749, 0.078086, 0.097141, 0.097892, 0.083314, 0.050181],$$


$$C_2^T = [0.642824, 0.766876, 0.847149, 0.85023, 0.739863, 0.453045],$$

$$\sigma_{1000}^1 = 5.968 \times 10^{-6}, \quad \sigma_{500}^1 = 6.001 \times 10^{-6},$$


$$\sigma_{1000}^2 = 1.102 \times 10^{-5}, \quad \sigma_{500}^2 = 1.109 \times 10^{-5}.$$

**Table 2.** Approximate and exact values of  $\mu$  for Example 2

$Z$	0.0	0.2	0.4	0.6	0.8	1.0
$\tilde{\mu}_1(z)$	0.999978	1.221404	1.491823	1.822111	2.225541	2.718264
$\mu_1(z)$	1.000000	1.221403	1.491825	1.822119	2.225541	2.718271
$\mu_2(z)$	-0.000022	0.079194	0.146128	0.20413	0.255279	0.301079
$\tilde{\mu}_2(z)$	0.000000	0.079181	0.146128	0.20412	0.255273	0.30103



**Figure 3.** Absolute errors with added noise  $\delta_1(=\sigma_N)$  for  $N = 1000$



**Figure 4.** Absolute errors with added noise  $\delta_1(=\sigma_N)$  for  $N = 500$

**Example 3.**

This example presents the problem (1) with

$$q_{11}(y) = e^{2y} + 1, \quad q_{12}(y) = e^y, \quad q_{21}(y) = e^y / 2, \quad q_{22}(y) = e^{3y}, \quad \mu = 1/4 \text{ and}$$

$$h_1(y) = \left[ e^y \left( 2(3\sqrt{(1-y)y} + 4\sqrt{(1-y)y^3} + 8\sqrt{(1-y)y^5} / 15\sqrt{y}) \right) \right. \\ \left. + (1 + e^{2y})(e^y(\Gamma(3/4) - \Gamma(3/4, y)) + 4y^{3/4}/3) \right],$$

$$h_2(y) = \left[ e^{3y}(16/15 y^{5/2}) + \frac{e^{2y}(1-y)^{3/4}((\Gamma(3/4) - \Gamma(3/4, y-1))}{2(y-1)^{3/4}} \right].$$

This has the exact solution  $\mu_1(z) = e^z + 1$  and  $\mu_2(z) = z^2$ . The corresponding values of  $C_1^T, C_2^T, \sigma_{1000}^1, \sigma_{500}^1, \sigma_{1000}^2$  and  $\sigma_{500}^2$  are given as

$$C_1^T = [0.019755, 0.089266, 0.185858, 0.25288, 0.255705, 0.166674],$$


$$C_2^T = [1.195592, 1.266867, 1.288123, 1.222903, 1.028541, 0.619684],$$

$$\sigma_{1000}^1 = 3.259 \times 10^{-5}, \quad \sigma_{500}^1 = 3.284 \times 10^{-5},$$


$$\sigma_{1000}^2 = 4.651 \times 10^{-5}, \quad \sigma_{500}^2 = 4.692 \times 10^{-5}.$$

**Table 3.** Approximate and exact values of  $\mu$  for Example 3

$Z$	0.0	0.2	0.4	0.6	0.8	1.0
$\tilde{\mu}_1(z)$	2.000092	2.221366	2.491837	2.822111	3.225568	3.718126
$\mu_1(z)$	2.000000	2.221403	2.491825	2.822119	3.225541	3.718271
$\mu_2(z)$	0.000233	0.039951	0.160026	0.359993	0.640037	0.99986
$\tilde{\mu}_2(z)$	0.00000	0.040000	0.160000	0.360000	0.640000	1.0000



**Figure 5.** Absolute errors with added noise  $\delta_1(=\sigma_N)$  for  $N = 1000$



**Figure 6.** Absolute errors with added noise  $\delta_1(=\sigma_N)$  for  $N = 500$

**Example 4:**

Consider the system of generalized Abel integral equation with  $q_{11}(y) = \sin y$ ,

$$q_{12}(y) = \cos 2y, q_{21}(y) = 2 \sin y, q_{22}(y) = \sin y / 2, b = 1/2$$

and

$$h_1(y) = \left[ y \sin y + \frac{1}{2} \cos 2y \left( \ln \left( \frac{1 + \sqrt{1 - y^2}}{y} \right) y^2 + \sqrt{1 - y^2} \right) \right],$$

$$h_2(y) = \left[ \frac{1}{4} (\pi y^2 + 16(1-y^2)^{1/2}) \sin y \right].$$

This has the exact solution  $\mu_1(z) = z$  and  $\mu_2(z) = z^2$ . The corresponding values of  $C_1^T, C_2^T, \sigma_{1000}^1, \sigma_{500}^1, \sigma_{1000}^2$  and  $\sigma_{500}^2$  are given as

$$C_1^T = [0.01974, 0.089287, 0.185834, 0.252889, 0.255684, 0.166674],$$


$$C_2^T = [0.078967, 0.214286, 0.289776, 0.30879, 0.27218, 0.166680],$$

$$\sigma_{1000}^1 = 1.288 \times 10^{-5}, \quad \sigma_{500}^1 = 1.299 \times 10^{-5},$$


$$\sigma_{1000}^2 = 7.711 \times 10^{-5}, \quad \sigma_{500}^2 = 7.75 \times 10^{-5}.$$

**Table 4.** Approximate and exact values of  $\mu$  for example 4

$Z$	0.0	0.2	0.4	0.6	0.8	1.0
$\tilde{\mu}_1(z)$	-0.000004	0.199996	0.400001	0.600011	0.799993	1.000073
$\mu_1(z)$	0.0	0.20000	0.40000	0.60000	0.80000	1.0000
$\mu_2(z)$	-0.000016	0.039998	0.160001	0.360006	0.639996	1.000041
$\tilde{\mu}_2(z)$	0.0	0.04000	0.160000	0.360000	0.640000	1.00000



**Figure 7.** Absolute errors with added noise  $\delta_1 (= \sigma_N)$  for  $N = 1000$



**Figure 8.** Absolute errors with added noise  $\delta_1 (= \sigma_N)$  for  $N = 500$

### Example 5.

Consider the system of generalized Abel integral equation with  $q_{ij}(y) = 1$ , for  $i, j = 1, 2$ ,  $b = 1/3$  and

$$h_1(y) = \left[ \frac{1}{4} \left( y \Gamma\left(\frac{2}{3}\right) \left( \frac{2\Gamma\left(\frac{-17}{12}\right)\sqrt{y}}{\Gamma\left(\frac{-3}{4}\right)} + \frac{\sqrt{\pi}}{\Gamma\left(\frac{13}{6}\right)} \right) + 3 \right) y^{4/3} + \frac{16}{17} {}_2F_1\left(\frac{-17}{12}, \frac{1}{3}, -\frac{5}{12}, y^2\right) \right],$$

$$h_2(y) = \left[ \frac{1}{4} \left( \Gamma\left(\frac{2}{3}\right) \left( \frac{2\sqrt{y}\Gamma\left(\frac{7}{4}\right)}{\Gamma\left(\frac{29}{12}\right)} - \frac{\Gamma\left(\frac{-7}{6}\right)}{\sqrt{\pi}} \right) y^{7/3} + 3(1-y^2)^{2/3} \right) + \frac{3}{7} {}_2F_1\left(-\frac{7}{6}, \frac{1}{3}, -\frac{1}{6}, y^2\right) \right].$$

This has the exact solution  $\mu_1(z) = z + z^2$  and  $\mu_2(z) = z^{5/2}$ . The corresponding values of  $C_1^T, C_2^T, \sigma_{1000}^1, \sigma_{500}^1, \sigma_{1000}^2$  and  $\sigma_{500}^2$  are given as

$$C_1^T = [0.011095, 0.060171, 0.148163, 0.227998, 0.247406, 0.166718],$$


$$C_2^T = [0.098659, 0.303574, 0.475602, 0.56168, 0.527859, 0.333320],$$

$$\sigma_{1000}^1 = 2.091 \times 10^{-5}, \quad \sigma_{500}^1 = 2.104 \times 10^{-5},$$


$$\sigma_{1000}^2 = 5.503 \times 10^{-5}, \quad \sigma_{500}^2 = 5.65 \times 10^{-5}.$$

**Table 5.** Approximate and exact values of  $\mu$  for example 5

$Z$	0.0	0.2	0.4	0.6	0.8	1.0
$\tilde{\mu}_1(z)$	-0.00006	0.239993	0.559996	0.959993	1.440007	1.999917
$\mu_1(z)$	0.0	0.24	0.56	0.96	1.44	2.0
$\mu_2(z)$	0.000271	0.01792	0.101171	0.278885	0.572426	1.000307
$\tilde{\mu}_2(z)$	0.0	0.017889	0.101193	0.278855	0.572433	1.0



**Figure 9.** Absolute errors with added noise  $\delta_1(=\sigma_N)$  for  $N = 1000$



**Figure 10.** Absolute errors with added noise  $\delta_1(=\sigma_N)$  for  $N = 500$

## 7. Conclusion

We have composed an almost Bernstein polynomials operational matrices of integration (AOMI) and used them to present an efficient and stable method for numerical approximation of system of singular integral equation with generalized Abel type kernel. The stability of method with relevancy to the data is restored, and a favorable result is attained, even for highly added noise modest sample intervals in the data. The selection of a small number of orthonormal polynomials (as shown for  $\alpha = 5$ ) makes the method easy and straight forward to use. This algorithm can be applied to any system of integral equations to get the desired numerical inversion, which is one of the applications of the proposed method.

### ***Acknowledgment:***

*The author wish to thank the anonymous referees and Professor Aliakbar Montazer Haghighi for their careful review and valuable suggestions that led to considerable improvement in the presentation of this manuscript.*

## REFERENCES

- Burton, T. A. (2005). Volterra integral and differential equations, Elsevier B. V., Netherlands.
- Dixit, S., Pandey, R. K., Kumar, S. and Singh, O. P. (2011). Solution of the generalized Abel integral equation by using almost Bernstein operational matrix, American Journal of Computational Mathematics, Vol. 1, No. 4, pp. 226-234.
- Dixit, S., Singh, O. P. and Kumar, S. (2012). A stable numerical inversion of generalized Abel's integral equation, Applied Numerical Mathematics, Vol. 62, No. 5, pp.567-579.
- Irfan, N., Kumar, S. and Kapoor, S. (2014). Bernstein operational matrix approach for integro-differential equation arising in control theory, Nonlinear Engineering, Vol. 3, No. 2, pp. 117-123.
- Kumar, S., Kumar, R., Cattani, C. and Samet, B. (2020). Chaotic behavior of fractional predator – prey dynamical system, Chaos, Soliton & Fractals, Vol. 135, pp. 109811.
- Kumar, S., Ahmadian, A., Kumar, R., Kumar, D., Singh, J., Baleanu, D. and Salimi, M. (2020). An efficient numerical method for fractional SIR epidemic model of Infectious disease by using Bernstein wavelets, Mathematics, Vol. 8, pp. 558.
- Kumar, S. and Singh, O. P. (2010). Numerical inversion of the abel integral equation using homotopy perturbation method, Zeitschrift für Naturforschung A, Vol. 65, No. 8-9, pp. 677-682.
- Kumar, S., Singh, J., Kumar, D. and Kapoor, S. (2014). New homotopy analysis transform algorithm to solve Volterra integral equation, Ain Shams Engineering Journal, Vol. 5, No. 1, pp. 243-246.
- Kumar, S., Kumar, A., Kumar, D., Singh, J. and Singh, A. (2015). Analytical solution of Abel integral equation arising in astrophysics via Laplace transform, Journal of the Egyptian Mathematical Society, Vol. 23, No. 1, pp. 102-107.



- Kumar, S., Singh, O. P. and Dixit, S. (2011). Homotopy Perturbation Method for Solving System of Generalized Abel's Integral Equations, Applications and Applied Mathematics (AAM), Vol. 6, No. 1, pp. 268 – 283.
- Kumar, S., Singh, O. P. and Dixit, S. (2011). Generalized Abel inversion using Homotopy perturbation method, Applied Mathematics, Vol. 2, No. 2, pp. 254- 257.
- Lowengrub, M. (1976). Systems of Abel type integral equations in function theoretic methods in differential equations, In: Gilbert, R. P. and Weinacht, R. J. (eds), pp. 277-296.
- Maleknejad, K. and Aghazadeh, N. (2005). Numerical solution of Volterra integral equations of the second kind with convolution type kernel by using Taylors- series expansion method, Appl. Math. Computation, Vol. 161, pp. 915-922.
- Maleknejad, K., Sohrabi, S. and Rostami, S. Y. (2007). Numerical solution of nonlinear Volterra integral equation of the second kind by using Chebyshev polynomials, Appl. Math. Computation, Vol. 188, pp. 123-128.
- Mandal, N., Chakrabarti, A. and Mandal, B.N. (1996). Numerical solution of a system of generalized Abel's integral equation using fractional calculus, Appl. Math. Letters, Vol. 9, No. 5, pp.1-4.
- Mandal, B. N. and Chakrabarti, A. (2016). Applied singular integral equations, CRC press, pp.231-239.
- Pandey, R. K., Singh, O. P. and Singh, K. (2009). Numerical solution of system of Volterra integral equations using Bernstein polynomials, Nonl. Sci. and Simulation, Vol. 10, No. 7, pp. 891-895.
- Singha, N. and Nahak, C. (2019). Solution of Generalized Abel's integral equation using Laguerre orthogonal approximation, Applications and Applied Mathematics (AAM), Vol. 14, No. 2, pp. 1051 – 1066.
- Tahmasbi, A. and Fard, O. S. (2008). Numerical solution of linear Volterra integral equations system of the second kind, Appl. Math. Computation, Vol. 201, pp. 547-552.
- Walton, J. (1979). System of generalized Abel's integral equation with applications to simultaneous dual relations, SIAM J. Math. Analysis, Vol. 10, No. 4, pp. 808-822.
- Yalsinbas, S. (2002). Taylor polynomials solutions of nonlinear Volterra–Fredholm integral equations, Appl. Math. Computation, Vol. 127, pp. 195-206.