



4-2020

Some Quadratic Transformations and Reduction Formulas associated with Hypergeometric Functions

M. I. Qureshi
Jamia Millia Islamia (A Central University)

M. Kashif Khan
Jamia Millia Islamia (A Central University)

Follow this and additional works at: <https://digitalcommons.pvamu.edu/aam>



Part of the [Special Functions Commons](#)

Recommended Citation

Qureshi, M. I. and Khan, M. Kashif (2020). Some Quadratic Transformations and Reduction Formulas associated with Hypergeometric Functions, *Applications and Applied Mathematics: An International Journal (AAM)*, Vol. 15, Iss. 3, Article 6.

Available at: <https://digitalcommons.pvamu.edu/aam/vol15/iss3/6>

This Article is brought to you for free and open access by Digital Commons @PVAMU. It has been accepted for inclusion in *Applications and Applied Mathematics: An International Journal (AAM)* by an authorized editor of Digital Commons @PVAMU. For more information, please contact hvkoshy@pvamu.edu.



Some Quadratic Transformations and Reduction Formulas associated with Hypergeometric Functions

¹M.I. Qureshi and ^{2,*}M. Kashif Khan

Department of Applied Sciences and Humanities
Jamia Millia Islamia (A Central University)
Maulana Mohammad Ali Jauhar Marg, Jamia Nagar
New Delhi-110025, India

¹miqureshi_delhi@yahoo.co.in, ²md.kashifkhan85@gmail.com

*Corresponding Author

Received: February 29, 2020; Accepted: March 15, 2020

Abstract

In this paper, we construct four summation formulas for terminating Gauss' hypergeometric series having argument "two" and with the help of our summation formulas. We establish two quadratic transformations for Gauss' hypergeometric function in terms of finite summation of combination of two Clausen hypergeometric functions. Further, we have generalized our quadratic transformations in terms of general double series identities as well as in terms of reduction formulas for Kampé de Fériet's double hypergeometric function. Some results of Rathie-Nagar, Kim et al. and Choi-Rathie are also obtained as special cases of our findings.

Keywords: Generalized hypergeometric function; Hypergeometric summation theorem; Bounded sequence; Quadratic transformation; Reduction formulas

MSC 2010 No.: 33C05, 33C20

1. Introduction

Functions which are important enough to be given their own name are known as “special function”. These include the well known logarithmic, exponential, and trigonometric functions and extend to cover the gamma, beta, and zeta functions and the class of orthogonal polynomials, among many others. Special functions have broad applications in pure mathematics as well as in applied areas such as quantum mechanics, solutions of wave equations, moments of inertia, fluid dynamics, and heat conduction.

In 1656, John Wallis was the first mathematician who used the term “hypergeometric series” in his treatise “Arithmetica Infinitorum” to explicate the infinite series of the form $1 + a + a(a + 1) + a(a + 1)(a + 2) + \dots$. In 1730, Euler established nearly all of the significant properties of the Gamma functions. The series ${}_1F_1$ was introduced by Kümmer (1836) and the series ${}_3F_2$ was given by Clausen (1828). In the theory of hypergeometric series a major development was given (although Euler and Pfaff had been found certain important results) by Gauss (1866) on ${}_2F_1$ series. The Hypergeometric functions of two variables was introduced by Appell (1880) and Lauricella (1893) generalized them to several variables.

It is interesting to mention here that the results are very important for the application point of view, whenever hypergeometric functions reduce to gamma functions. Thus, the classical theorems such as Gauss, Kümmer and Bailey for the series ${}_2F_1$ and of Watson, Dixon, Whipple and Saalschütz for the series ${}_3F_2$ and others play an important role in the theory of hypergeometric function, generalized hypergeometric function and in other applicable sciences. In a series of papers Lavoie et al. (1992, 1994, 1996) studied generalizations of some of the above classical summation theorems. In the theory of special functions, summations, transformations and reduction formulas have received keen attention during the last few years (see Chen and Srivastava (2005), Chen et al. (2006), Chu (2011), Kim (2009), Kim et al. (2012) and Kim et al. (2010)). Recently, in Miller (2009), Miller and Paris (2011a, 2011b) and Miller and Srivastava (2010) examined the generalized hypergeometric series (together with integer parameter differences) and derived several transformations and summation formulas.

Influenced by the recent work of researchers Choi and Rathie (2014), Meethal et al. (2015), Ebisu (2017), Kim, Gaboury et al. (2018, 2018), Miller and Paris (2013), Qureshi et al. (2016), and Srivastava (2016), some new summation formulas, quadratic transformations, generalizations of quadratic transformations and reduction formulas for Kampé de Fériet’s double hypergeometric function are obtained in this article.

2. Preliminaries

In our present investigation, we shall make use of the following standard notations and results.

$$\mathbb{N} = \{1, 2, 3, \dots\}, \mathbb{N}_0 = \mathbb{N} \cup \{0\} \text{ and } \mathbb{Z}_0^- = \mathbb{Z}^- \cup \{0\} = \{0, -1, -2, -3, \dots\}.$$

As usual, the symbols \mathbb{C} , \mathbb{R} , \mathbb{N} , \mathbb{Z} , \mathbb{R}^+ and \mathbb{R}^- denote the sets of complex numbers, real numbers,

natural numbers, integers, positive and negative real numbers, respectively.

The well-known Pochhammer’s symbol $(\alpha)_p$ ($\alpha, p \in \mathbb{C}$) is defined by (see Rainville (1960)), Srivastava and Manocha (1984))

$$(\alpha)_p = \frac{\Gamma(\alpha + p)}{\Gamma(\alpha)} = \begin{cases} 1; & (p = 0; \alpha \in \mathbb{C} \setminus \{0\}), \\ \alpha(\alpha + 1)(\alpha + 2) \cdots (\alpha + n - 1); & (p = n \in \mathbb{N}; \alpha \in \mathbb{C}), \\ \frac{(-1)^k n!}{(n-k)!}; & (\alpha = -n; p = k; k, n \in \mathbb{N}_0; 0 \leq k \leq n), \\ 0; & (\alpha = -n; p = k; k, n \in \mathbb{N}_0; k > n), \\ \frac{(-1)^n}{(1-\alpha)_n}; & (p = -n; n \in \mathbb{N}; \alpha \in \mathbb{C} \setminus \mathbb{Z}), \end{cases} \quad (1)$$

it being understood usually that $(0)_0 = 1$ and concluded tacitly that the Gamma quotient exists.

Generalized ordinary hypergeometric function of one variable is defined by (see Srivastava and Manocha (1984))

$${}_A F_B \left[\begin{matrix} (a_A); \\ (b_B); \end{matrix} z \right] \equiv {}_A F_B \left[\begin{matrix} a_1, a_2, \dots, a_A; \\ b_1, b_2, \dots, b_B; \end{matrix} z \right] = \sum_{k=0}^{\infty} \frac{(a_1)_k (a_2)_k \cdots (a_A)_k z^k}{(b_1)_k (b_2)_k \cdots (b_B)_k k!}, \quad (2)$$

where denominator parameters b_1, b_2, \dots, b_B are neither zero nor negative integers, and A, B are non-negative integers and numerator parameters a_1, a_2, \dots, a_A may be zero or negative integers.

Convergence Conditions of Series (2):

Suppose that numerator parameters a_1, a_2, \dots, a_A are neither zero nor negative integers (otherwise the question of convergence will not arise).

- (i) If $A \leq B$, then series ${}_A F_B$ is always convergent for all finite values of z (real or complex), i.e., $|z| < \infty$.
- (ii) If $A = B + 1$ and $|z| < 1$, then series ${}_A F_B$ is convergent.
- (iii) If $A = B + 1$ and $|z| = 1$, then series ${}_A F_B$ is absolutely convergent, when

$$\Re \left\{ \sum_{m=1}^B b_m - \sum_{n=1}^A a_n \right\} > 0,$$

- (iv) If $A = B + 1$ and $|z| = 1$, but $z \neq 1$, then series ${}_A F_B$ is conditionally convergent, when

$$-1 < \Re \left\{ \sum_{m=1}^B b_m - \sum_{n=1}^A a_n \right\} \leq 0,$$

where \Re denotes the real part of complex number throughout this paper.

Kampé de Fériet’s Double Hypergeometric Function:

Kampé de Fériet (1921) combined and generalized Appell's four double hypergeometric functions F_1, F_2, F_3, F_4 (see Srivastava and Manocha (1984)) and their seven confluent forms $\Phi_1, \Phi_2, \Phi_3, \Psi_1, \Psi_2, \Xi_1, \Xi_2$ given by Humbert (1920-21). We reminisce here the definition of general double hypergeometric function of Kampé de Fériet (see Appell and Kampé de Fériet (1926)), Burchnall and Chaundy (1941)) in the slightly refined notation of Srivastava and Panda (1976), Srivastava and Pathan (1979). Thus, the appropriate generalization of the Kampé de Fériet function is given by

$$F_{E;G;H}^{A;B;D} \left[\begin{array}{l} (a_A) : (b_B) ; (d_D) ; \\ (e_E) : (g_G) ; (h_H) ; \end{array} ; x, y \right] = \sum_{m=0}^{\infty} \sum_{n=0}^{\infty} \frac{[(a_A)]_{m+n} [(b_B)]_m [(d_D)]_n x^m y^n}{[(e_E)]_{m+n} [(g_G)]_m [(h_H)]_n m! n!}, \quad (3)$$

where (a_A) denotes the array of A number of parameters a_1, a_2, \dots, a_A and

$$[(a_A)]_m = \prod_{j=1}^A (a_j)_m,$$

with similar interpretation for others.

Convergence Conditions of Double Series (3) (see Srivastava and Panda (1976)):

- (i) $A + B < E + G + 1, A + D < E + H + 1, |x| < \infty, |y| < \infty,$ or
(ii) $A + B = E + G + 1, A + D = E + H + 1,$ and

$$\left\{ \begin{array}{l} |x|^{\frac{1}{(A-E)}} + |y|^{\frac{1}{(A-E)}} < 1, \text{ if } A > E, \\ \max\{|x|, |y|\} < 1, \quad \text{if } A \leq E \end{array} \right\}.$$

For absolutely and conditionally convergence of double series (3), the reader can consult an article by Hàì et al. (1992).

The Series Decomposition Identity is given by (see Srivastava and Manocha (1984))

$$\sum_{n=0}^{\infty} \Lambda(n) = \sum_{n=0}^{\infty} \Lambda(2n) + \sum_{n=0}^{\infty} \Lambda(2n + 1). \quad (4)$$

The Reversal of the Order of Terms in Finite Summation is given by

$$\sum_{r=0}^n \Phi(r) = \sum_{r=0}^n \Phi(n - r), \quad (n \in \mathbb{N}_0). \quad (5)$$

The Series Rearrangement Formulas are given as follows (see Srivastava and Manocha (1984))

$$\sum_{n=0}^{\infty} \sum_{m=0}^{\infty} \Xi(m, n) = \sum_{n=0}^{\infty} \sum_{m=0}^n \Xi(m, n - m), \tag{6}$$

and

$$\sum_{m=0}^{\infty} \sum_{n=0}^{\infty} \Psi(m, n) = \sum_{m=0}^{\infty} \sum_{n=0}^{\lfloor \frac{m}{2} \rfloor} \Psi(m - 2n, n), \tag{7}$$

provided that involved double series are absolutely convergent and $[k]$ denotes the greatest integer for $k \in \mathbb{R}$.

The rest of the article is organized as follows. In Section 3, we obtain four summation formulas for terminating Gauss’ hypergeometric series having argument “two” by means of summation formula of Rakha and Rathie (2011), and summation formula recorded by Prudnikov et al. (1990). In Section 4, as an application of our four summation formulas, we established two new quadratic transformations for Gauss’ hypergeometric function. In Section 5, we have generalized our quadratic transformations in terms of general double series identities having bounded sequences. In Section 6, we obtain two results for the reducibility of Kampé de Fériet double hypergeometric functions. In Section 7, we discuss several special cases of Sections 4 and 6. Any values of parameters and variables leading to the results in following sections which do not make sense are tacitly excluded.

3. Four Summation Formulas

The following four summation formulas for terminating Gauss series will be derived in this section.

First Summation Formula.

$${}_2F_1 \left[\begin{matrix} -2n, a; \\ 2a + j; \end{matrix} \right]_2 = \frac{\Gamma(1 - a)}{2^{2a+j}(a)_j \Gamma(1 - 2a - j)} \sum_{r=0}^j \left\{ \binom{j}{r} (-1)^r \frac{\Gamma(\frac{1-2a-j+r}{2})(\frac{1-r+j}{2})_n}{\Gamma(\frac{1+r-j}{2})(\frac{1+2a-r+j}{2})_n} \right\}, \tag{8}$$

$$\left(a, 1 - a, 2a + j, 1 - 2a - j \in \mathbb{C} \setminus \mathbb{Z}_0^- ; n, j \in \mathbb{N}_0 \right).$$

Second Summation Formula.

$${}_2F_1 \left[\begin{matrix} -2n - 1, a; \\ 2a + j; \end{matrix} \right]_2 = \frac{-\Gamma(1 - a)}{2^{2a+j}(a)_j \Gamma(1 - 2a - j)} \sum_{r=0}^j \left\{ \binom{j}{r} (-1)^r \frac{\Gamma(\frac{-2a-j+r}{2})(\frac{2-r+j}{2})_n}{\Gamma(\frac{r-j}{2})(\frac{2+2a-r+j}{2})_n} \right\}, \tag{9}$$

$$\left(a, 1 - a, 2a + j, 1 - 2a - j \in \mathbb{C} \setminus \mathbb{Z}_0^- ; n, j \in \mathbb{N}_0 \right).$$

Third Summation Formula.

$${}_2F_1 \left[\begin{matrix} -2n, a; \\ 2a - j; \end{matrix} 2 \right] = \frac{\Gamma(1 - a)}{2^{2a-j}\Gamma(1 - 2a + j)} \sum_{r=0}^j \left\{ \binom{j}{r} \frac{\Gamma(\frac{1-2a+j+r}{2})(\frac{1-r+j}{2})_n}{\Gamma(\frac{1+r-j}{2})(\frac{1+2a-r-j}{2})_n} \right\}, \quad (10)$$

$$\left(a, 1 - a, 2a - j, 1 - 2a + j \in \mathbb{C} \setminus \mathbb{Z}_0^- ; n, j \in \mathbb{N}_0 \right).$$

Fourth Summation Formula.

$${}_2F_1 \left[\begin{matrix} -2n - 1, a; \\ 2a - j; \end{matrix} 2 \right] = \frac{-\Gamma(1 - a)}{2^{2a-j}\Gamma(1 - 2a + j)} \sum_{r=0}^j \left\{ \binom{j}{r} \frac{\Gamma(\frac{-2a+j+r}{2})(\frac{2-r+j}{2})_n}{\Gamma(\frac{r-j}{2})(\frac{2+2a-r-j}{2})_n} \right\}, \quad (11)$$

$$\left(a, 1 - a, 2a - j, 1 - 2a + j \in \mathbb{C} \setminus \mathbb{Z}_0^- ; n, j \in \mathbb{N}_0 \right).$$

where the notation $\binom{j}{r}$ is the binomial coefficient.

Remark 3.1.

We have verified above summation formulas for suitable numerical values of n , a and j .

In order to start the derivation of above summation formulas consider the series,

$${}_2F_1 \left[\begin{matrix} -n, a; \\ 2a + j; \end{matrix} 2 \right] = \sum_{r=0}^n \frac{(-n)_r (a)_r 2^r}{(2a + j)_r r!} = \sum_{r=0}^n \frac{n! (-1)^r (a)_r 2^r}{(2a + j)_r r! (n - r)!},$$

which, upon replacing r by $n - r$ and applying reversal of the order of summation (5) and some algebraic properties of Pochhammer's symbol (1), becomes

$${}_2F_1 \left[\begin{matrix} -n, a; \\ 2a + j; \end{matrix} 2 \right] = \frac{(-1)^n (a)_n 2^n}{(2a + j)_n} {}_2F_1 \left[\begin{matrix} -n, 1 - 2a - j - n; \\ 1 - a - n; \end{matrix} \frac{1}{2} \right]. \quad (12)$$

Proof:

In order to derive the first summation formula (8), we begin as follows.

Replacing n by $2n$ on both sides of equation (12), we obtain

$${}_2F_1 \left[\begin{matrix} -2n, a; \\ 2a + j; \end{matrix} 2 \right] = \frac{(-1)^{2n} (a)_{2n} 2^{2n}}{(2a + j)_{2n}} {}_2F_1 \left[\begin{matrix} -2n, 1 - 2a - j - 2n; \\ 1 - a - 2n; \end{matrix} \frac{1}{2} \right]. \quad (13)$$

We now applying the summation formula of Rakha and Rathie (2011) in the right-hand side of equation (13), after straightforward calculation we easily arrive at the right-hand side of equation (8).

Similarly, by using the result (12) in conjugation with the result of Rakha and Rathie (2011), we can prove the second summation formula (9). ■

Proof:

In order to find the third summation formula (10), we continue as follows.

Changing n by $2n$ and j by $-j$ on both sides of equation (12), to yield

$${}_2F_1 \left[\begin{matrix} -2n, a; \\ 2a - j; \end{matrix} 2 \right] = \frac{(-1)^{2n} (a)_{2n} 2^{2n}}{(2a - j)_{2n}} {}_2F_1 \left[\begin{matrix} -2n, 1 - 2a + j - 2n; \\ 1 - a - 2n; \end{matrix} \frac{1}{2} \right]. \quad (14)$$

By employing the summation formula recorded by Prudnikov et al. (1990) in the right-hand side of equation (14), after little algebra we obtain the right-hand side of equation (10).

Similarly, by utilizing the result (12) and summation formula recorded by Prudnikov et al. (1990), we can derive the fourth summation formula (11). ■

4. Two Quadratic Transformations

In this section, we shall establish the following two new quadratic transformations for Gauss series by means of series rearrangement technique and use of hypergeometric summation formulas (8) to (11).

First Transformation. The following transformation holds true:

$$\begin{aligned}
(1+z)^{-2b} {}_2F_1 \left[\begin{matrix} 2b, a; \\ 2a+j; \end{matrix} \frac{2z}{1+z} \right] &= \frac{\Gamma(1-a)}{2^{2a+j} (a)_j \Gamma(1-2a-j)} \sum_{r=0}^j \binom{j}{r} (-1)^r \left\{ \frac{\Gamma(\frac{1-2a-j+r}{2})}{\Gamma(\frac{1+r-j}{2})} \right. \\
&\times {}_3F_2 \left[\begin{matrix} b, b+\frac{1}{2}, \frac{1-r+j}{2}; \\ \frac{1+2a+j-r}{2}, \frac{1}{2}; \end{matrix} z^2 \right] + 2bz \frac{\Gamma(\frac{-2a-j+r}{2})}{\Gamma(\frac{r-j}{2})} \\
&\left. \times {}_3F_2 \left[\begin{matrix} b+\frac{1}{2}, b+1, \frac{2-r+j}{2}; \\ \frac{2+2a+j-r}{2}, \frac{3}{2}; \end{matrix} z^2 \right] \right\}, \tag{15}
\end{aligned}$$

$$\left(b, a, 1-a, 2a+j, 1-2a-j \in \mathbb{C} \setminus \mathbb{Z}_0^-; j \in \mathbb{N}_0 \right).$$

Second Transformation. The following transformation holds true:

$$\begin{aligned}
(1+z)^{-2b} {}_2F_1 \left[\begin{matrix} 2b, a; \\ 2a-j; \end{matrix} \frac{2z}{1+z} \right] &= \frac{\Gamma(1-a)}{2^{2a-j} \Gamma(1-2a+j)} \sum_{r=0}^j \binom{j}{r} \left\{ \frac{\Gamma(\frac{1-2a+j+r}{2})}{\Gamma(\frac{1+r-j}{2})} \right. \\
&\times {}_3F_2 \left[\begin{matrix} b, b+\frac{1}{2}, \frac{1-r+j}{2}; \\ \frac{1+2a-j-r}{2}, \frac{1}{2}; \end{matrix} z^2 \right] + 2bz \frac{\Gamma(\frac{r-2a+j}{2})}{\Gamma(\frac{r-j}{2})} \\
&\left. \times {}_3F_2 \left[\begin{matrix} b+\frac{1}{2}, b+1, \frac{2-r+j}{2}; \\ \frac{2+2a-j-r}{2}, \frac{3}{2}; \end{matrix} z^2 \right] \right\}, \tag{16}
\end{aligned}$$

$$\left(b, a, 1-a, 2a-j, 1-2a+j \in \mathbb{C} \setminus \mathbb{Z}_0^-; j \in \mathbb{N}_0 \right).$$

Proof:

In order to establish first transformation (15), we proceed as follows.

Denoting the series expansion of left-hand side of transformation (15) by ζ and after some simplifications, we have

$$\begin{aligned}
 \zeta &= \sum_{m=0}^{\infty} \frac{(2b)_m (a)_m 2^m z^m}{(2a+j)_m m!} {}_1F_0 \left[\begin{matrix} 2b+m; \\ \text{---}; \end{matrix} -z \right] \\
 &= \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} \frac{(2b)_{m+n} (a)_m 2^m (-1)^n z^{n+m}}{(2a+j)_m m! n!} \\
 &= \sum_{n=0}^{\infty} \frac{(2b)_n (-1)^n z^n}{n!} \sum_{m=0}^n \frac{(a)_m 2^m (-n)_m}{(2a+j)_m m!} \\
 &= \sum_{n=0}^{\infty} \frac{(2b)_n (-1)^n z^n}{n!} {}_2F_1 \left[\begin{matrix} -n, a; \\ 2a+j; \end{matrix} 2 \right]. \tag{17}
 \end{aligned}$$

Now, use series decomposition identity (4) in the right-hand side of Equation (17), to yield

$$\zeta = \sum_{n=0}^{\infty} \frac{(2b)_{2n} (-1)^{2n} z^{2n}}{(2n)!} {}_2F_1 \left[\begin{matrix} -2n, a; \\ 2a+j; \end{matrix} \right] + \sum_{n=0}^{\infty} \frac{(2b)_{2n+1} (-1)^{2n+1} z^{2n+1}}{(2n+1)!} {}_2F_1 \left[\begin{matrix} -2n-1, a; \\ 2a+j; \end{matrix} \right]. \tag{18}$$

Finally, by employing summation formulas (8) and (9) in the right-hand side of equation (18), after some simplifications we get the right-hand side of transformation (15). ■

Proof:

In order to establish second transformation (16), we proceed as follows.

Denoting the series expansion of left-hand side of transformation (16) by ω and after some simplifications, we have

$$\begin{aligned}
 \omega &= \sum_{m=0}^{\infty} \frac{(2b)_m (a)_m 2^m z^m}{(2a-j)_m m!} {}_1F_0 \left[\begin{matrix} 2b+m; \\ \text{---}; \end{matrix} -z \right] \\
 &= \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} \frac{(2b)_{m+n} (a)_m 2^m (-1)^n z^{n+m}}{(2a-j)_m m! n!} \\
 &= \sum_{n=0}^{\infty} \frac{(2b)_n (-1)^n z^n}{n!} \sum_{m=0}^n \frac{(a)_m 2^m (-n)_m}{(2a-j)_m m!} \\
 &= \sum_{n=0}^{\infty} \frac{(2b)_n (-1)^n z^n}{n!} {}_2F_1 \left[\begin{matrix} -n, a; \\ 2a-j; \end{matrix} 2 \right]. \tag{19}
 \end{aligned}$$

Applying series decomposition identity (4) in Equation (19), we find

$$\begin{aligned} \omega = & \sum_{n=0}^{\infty} \frac{(2b)_{2n} (-1)^{2n} z^{2n}}{(2n)!} {}_2F_1 \left[\begin{matrix} -2n, a; \\ 2a - j; \end{matrix} \right. \\ & \left. + \sum_{n=0}^{\infty} \frac{(2b)_{2n+1} (-1)^{2n+1} z^{2n+1}}{(2n+1)!} {}_2F_1 \left[\begin{matrix} -2n-1, a; \\ 2a - j; \end{matrix} \right. \right]. \end{aligned} \quad (20)$$

By using summation formulas (10) and (11) in the right-hand side of Equation (20), after some simplifications we easily get the right-hand side of transformation (16). ■

5. General Double Series Identities

Here we generalize our quadratic transformations (15) and (16) as in the following theorems.

Theorem 5.1.

Let $\{\Phi(n)\}_{n=1}^{\infty}$ be a bounded sequence of essentially arbitrary complex numbers such that $\Phi(0) \neq 0$. Then, we have

$$\begin{aligned} \sum_{m=0}^{\infty} \sum_{n=0}^{\infty} \Phi(m+n) \frac{(a)_m (2z)^m (-z)^n}{(2a+j)_m m! n!} = & \frac{\Gamma(1-a)}{2^{2a+j} (a)_j \Gamma(1-2a-j)} \sum_{r=0}^j \binom{j}{r} (-1)^r \left\{ \frac{\Gamma(\frac{1-2a-j+r}{2})}{\Gamma(\frac{1+r-j}{2})} \right. \\ & \times \sum_{n=0}^{\infty} \Phi(2n) \frac{(\frac{1-r+j}{2})_n z^{2n}}{(\frac{1+2a+j-r}{2})_n (2n)!} + \frac{\Gamma(\frac{-2a-j+r}{2})}{\Gamma(\frac{r-j}{2})} \\ & \left. \times \sum_{n=0}^{\infty} \Phi(2n+1) \frac{(\frac{2-r+j}{2})_n z^{2n+1}}{(\frac{2+2a+j-r}{2})_n (2n+1)!} \right\}, \end{aligned} \quad (21)$$

$$\left(a, 1-a, 2a+j, 1-2a-j \in \mathbb{C} \setminus \mathbb{Z}_0^- ; j \in \mathbb{N}_0 \right),$$

provided that single and double series involved are absolutely convergent.

Theorem 5.2.

Let $\{\Phi(n)\}_{n=1}^{\infty}$ be a bounded sequence of essentially arbitrary complex numbers such that $\Phi(0) \neq 0$. Then, we have

$$\sum_{m=0}^{\infty} \sum_{n=0}^{\infty} \Phi(m+n) \frac{(a)_m (2z)^m (-z)^n}{(2a-j)_m m! n!} = \frac{\Gamma(1-a)}{2^{2a-j} \Gamma(1-2a+j)} \sum_{r=0}^j \binom{j}{r} \left\{ \frac{\Gamma(\frac{1-2a+j+r}{2})}{\Gamma(\frac{1+r-j}{2})} \right. \\ \times \sum_{n=0}^{\infty} \Phi(2n) \frac{(\frac{1-r+j}{2})_n z^{2n}}{(\frac{1-r+2a-j}{2})_n (2n)!} + \frac{\Gamma(\frac{r-2a+j}{2})}{\Gamma(\frac{r-j}{2})} \\ \left. \times \sum_{n=0}^{\infty} \Phi(2n+1) \frac{(\frac{2-r+j}{2})_n z^{2n+1}}{(\frac{2-r+2a-j}{2})_n (2n+1)!} \right\}, \quad (22)$$

$$\left(a, 1-a, 2a-j, 1-2a+j \in \mathbb{C} \setminus \mathbb{Z}_0^- ; j \in \mathbb{N}_0 \right),$$

provided that single and double series involved are absolutely convergent.

Proof:

In order to prove Theorem 5.1, we proceed as follows.

Denote the left-hand side of double-series identity (21) by S and replacing n by $n - m$, after some simplifications, we have

$$S = \sum_{n=0}^{\infty} \Phi(n) \frac{(-1)^n (z)^n}{n!} {}_2F_1 \left[\begin{matrix} -n, a; \\ 2a+j; \end{matrix} \quad 2 \right]. \quad (23)$$

Using series decomposition identity (4) in Equation (23), we get

$$S = \sum_{n=0}^{\infty} \Phi(2n) \frac{(-1)^{2n} (z)^{2n}}{2n!} {}_2F_1 \left[\begin{matrix} -2n, a; \\ 2a+j; \end{matrix} \quad 2 \right] \\ + \sum_{n=0}^{\infty} \Phi(2n+1) \frac{(-1)^{2n+1} (z)^{2n+1}}{2n+1!} {}_2F_1 \left[\begin{matrix} -2n-1, a; \\ 2a+j; \end{matrix} \quad 2 \right]. \quad (24)$$

Finally, by applying summation formulas (8) and (9) in the right-hand side of Equation (24), after simplification we easily arrive at the right-hand side of Equation (21). ■

Proof:

In order to prove Theorem 5.2, we proceed as follows.

Denote the left-hand side of double-series identity (22) by T and replacing n by $n - m$, after some

simplifications, we have

$$T = \sum_{n=0}^{\infty} \Phi(n) \frac{(-1)^n (z)^n}{n!} {}_2F_1 \left[\begin{matrix} -n, a; \\ 2a - j; \end{matrix} \middle| 2 \right]. \quad (25)$$

Using series decomposition identity (4) in Equation (25), to yield

$$T = \sum_{n=0}^{\infty} \Phi(2n) \frac{(-1)^{2n} (z)^{2n}}{2n!} {}_2F_1 \left[\begin{matrix} -2n, a; \\ 2a - j; \end{matrix} \middle| 2 \right] \\ + \sum_{n=0}^{\infty} \Phi(2n+1) \frac{(-1)^{2n+1} (z)^{2n+1}}{2n+1!} {}_2F_1 \left[\begin{matrix} -2n-1, a; \\ 2a - j; \end{matrix} \middle| 2 \right]. \quad (26)$$

By employing summation formulas (10) and (11) in the right-hand side of Equation (26), after some simplifications we easily obtain the right-hand side of Equation (22). ■

6. Reducibility of Kampé de Fériet Functions

In the assertions (21) and (22), putting $\Phi(n) = \left(\prod_{j=1}^D (d_j)_n \right) \left(\prod_{j=1}^E (e_j)_n \right)^{-1}$; $n \in \mathbb{N}_0$ and after straightforward calculation we obtain presumably new hypergeometric reduction formulas for the Kampé de Fériet double hypergeometric functions in two variables. Under the common convergence conditions of Kampé de Fériet double hypergeometric function $F_{E:1;0}^{D:1;0}(2z, -z)$ and generalized hypergeometric function ${}_{2D+1}F_{2E+2}(z^2)$ given below

(i) If $2D \leq 2E + 1$, then $|z| < \infty$,

(ii) If $D = E + 1$, then $|z| < \frac{1}{3}$, the following hypergeometric reduction formulas hold true.

First Reduction Formula.

$$F_{E:1;0}^{D:1;0} \left[\begin{matrix} (d_D) : a; \text{---}; \\ (e_E) : 2a + j; \text{---}; \end{matrix} \middle| 2z, -z \right] = \frac{\Gamma(1-a)}{2^{2a+j} (a)_j \Gamma(1-2a-j)} \sum_{r=0}^j \binom{j}{r} (-1)^r \left\{ \frac{\Gamma(\frac{1-2a-j+r}{2})}{\Gamma(\frac{1+r-j}{2})} \right. \\ \times {}_{2D+1}F_{2E+2} \left[\begin{matrix} \frac{(d_D)}{2}, \frac{1+(d_D)}{2}, \frac{1-r+j}{2}; \\ \frac{(e_E)}{2}, \frac{1+(e_E)}{2}, \frac{1+2a-r+j}{2}, \frac{1}{2}; \end{matrix} \middle| 4^{(D-E-1)} z^2 \right] \\ \left. + \frac{z \Gamma(\frac{-2a-j+r}{2}) \prod_{i=1}^D (d_i)}{\Gamma(\frac{r-j}{2}) \prod_{i=1}^E (e_i)} {}_{2D+1}F_{2E+2} \left[\begin{matrix} \frac{1+(d_D)}{2}, \frac{2+(d_D)}{2}, \frac{2-r+j}{2}; \\ \frac{1+(e_E)}{2}, \frac{2+(e_E)}{2}, \frac{2+2a+j-r}{2}, \frac{3}{2}; \end{matrix} \middle| 4^{(D-E-1)} z^2 \right] \right\}, \quad (27)$$

$$\left(a, 1-a, 2a+j, 1-2a-j, d_1, d_2, \dots, d_D, e_1, e_2, \dots, e_E \in \mathbb{C} \setminus \mathbb{Z}_0^-; j \in \mathbb{N}_0 \right).$$

Second Reduction Formula.

$$\begin{aligned}
 F_{E:1;0}^{D:1;0} \left[\begin{matrix} (d_D) : a; \text{---}; \\ (e_E) : 2a - j; \text{---}; \end{matrix} \quad 2z, -z \right] &= \frac{\Gamma(1-a)}{2^{2a-j}\Gamma(1-2a+j)} \sum_{r=0}^j \binom{j}{r} \left\{ \frac{\Gamma(\frac{1-2a+j+r}{2})}{\Gamma(\frac{1+r-j}{2})} \right. \\
 &\quad \times {}_{2D+1}F_{2E+2} \left[\begin{matrix} \frac{(d_D)}{2}, \frac{1+(d_D)}{2}, \frac{1-r+j}{2}; \\ \frac{(e_E)}{2}, \frac{1+(e_E)}{2}, \frac{1+2a-r-j}{2}, \frac{1}{2}; \end{matrix} \quad 4^{(D-E-1)} z^2 \right] \\
 &\quad \left. + \frac{z\Gamma(\frac{r-2a+j}{2}) \prod_{i=1}^D (d_i)}{\Gamma(\frac{r-j}{2}) \prod_{i=1}^E (e_i)} {}_{2D+1}F_{2E+2} \left[\begin{matrix} \frac{1+(d_D)}{2}, \frac{2+(d_D)}{2}, \frac{2-r+j}{2}; \\ \frac{1+(e_E)}{2}, \frac{2+(e_E)}{2}, \frac{2+2a-j-r}{2}, \frac{3}{2}; \end{matrix} \quad 4^{(D-E-1)} z^2 \right] \right\}, \quad (28) \\
 &\quad \left(a, 1-a, 2a-j, 1-2a+j, d_1, d_2, \dots, d_D, e_1, e_2, \dots, e_E \in \mathbb{C} \setminus \mathbb{Z}_0^-; j \in \mathbb{N}_0 \right).
 \end{aligned}$$

7. Special Cases

(i) Set $D = E = 0$, in Equation (27). We get

$$\begin{aligned}
 e^{-z} {}_1F_1 \left[\begin{matrix} a; \\ 2a+j; \end{matrix} \quad 2z \right] &= \frac{\Gamma(1-a)}{2^{2a+j}(a)_j\Gamma(1-2a-j)} \sum_{r=0}^j \binom{j}{r} (-1)^r \left\{ \frac{\Gamma(\frac{1-2a-j+r}{2})}{\Gamma(\frac{1+r-j}{2})} \right. \\
 &\quad \times {}_1F_2 \left[\begin{matrix} \frac{1-r+j}{2}; \\ \frac{1+2a-r+j}{2}, \frac{1}{2}; \end{matrix} \quad \frac{z^2}{4} \right] + \frac{z\Gamma(\frac{-2a-j+r}{2})}{\Gamma(\frac{r-j}{2})} {}_1F_2 \left[\begin{matrix} \frac{2-r+j}{2}; \\ \frac{2+2a+j-r}{2}, \frac{3}{2}; \end{matrix} \quad \frac{z^2}{4} \right] \left. \right\}, \quad (29) \\
 &\quad \left(a, 1-a, 2a+j, 1-2a-j \in \mathbb{C} \setminus \mathbb{Z}_0^-; j \in \mathbb{N}_0 \right).
 \end{aligned}$$

(ii) Put $D = E = 0$, in Equation (28). We obtain

$$\begin{aligned}
 e^{-z} {}_1F_1 \left[\begin{matrix} a; \\ 2a-j; \end{matrix} \quad 2z \right] &= \frac{\Gamma(1-a)}{2^{2a-j}\Gamma(1-2a+j)} \sum_{r=0}^j \binom{j}{r} \left\{ \frac{\Gamma(\frac{1-2a+j+r}{2})}{\Gamma(\frac{1+r-j}{2})} \right. \\
 &\quad \times {}_1F_2 \left[\begin{matrix} \frac{1-r+j}{2}; \\ \frac{1+2a-r-j}{2}, \frac{1}{2}; \end{matrix} \quad \frac{z^2}{4} \right] + \frac{z\Gamma(\frac{r-2a+j}{2})}{\Gamma(\frac{r-j}{2})} {}_1F_2 \left[\begin{matrix} \frac{2-r+j}{2}; \\ \frac{2+2a-j-r}{2}, \frac{3}{2}; \end{matrix} \quad \frac{z^2}{4} \right] \left. \right\}, \quad (30)
 \end{aligned}$$

$$\left(a, 1 - a, 2a - j, 1 - 2a + j \in \mathbb{C} \setminus \mathbb{Z}_0^- ; j \in \mathbb{N}_0 \right).$$

(iii) Set $j = 0$ in Equations (15) and (16); we get Kümmer's transformation (see Kümmer (1836)).

(iv) Put $j = 0$, in Equation (29); we obtain another transformation of Kümmer (see Rainville (1960)).

(v) Take $j = 1$, in Equations (29) and (30); we have known results of Rathie and Nagar (see Rathie and Nagar (1995)).

(vi) Fix $j = 0, 1$, in Equations (27) and (28); we recover known results of Kim (see Kim (2009)).

(vii) Substitute $j = 2, 3$, in Equations (29) and (30); we get known results of Choi and Rathie (see Choi and Rathie (2014)) and see also Kim et al. (2010).

8. Conclusion

In this paper, from Section 3 to Section 7, some interesting results are obtained for single and double hypergeometric functions, which may be potentially useful to non-specialists who are interested in Applied Mathematics or Mathematical Physics. We conclude our present analysis by observing that several interesting summation formulas, quadratic transformations, reduction formulas, corresponding multiple series identities and their hypergeometric representations can be obtained with the help of some newly derived terminating Clausen series of argument "two" in an analogous manner.

Acknowledgment:

The authors are thankful to the referees for their valuable comments and suggestions in improving this research paper.

REFERENCES

- Appell, P. (1880). Sur une classe de polynômes, Ann Sci. École Norm. Sup., Vol. 9, No. 2, pp. 119–144.
- Appell, P. and Kampé de Fériet, J. (1926). *Fonctions Hypergéométriques et Hypersphériques: Polynômes d' Hermite*, Gauthier-Villars, Paris.
- Burchnall, J. L. and Chaundy, T. W. (1941). Expansions of Appell's double hypergeometric functions-II, Quart. J. Math. (Oxford) Ser., Vol. 12, pp. 112–128.

Chen, K.-Y., Liu, S.-J. and Srivastava, H. M. (2006). Some double-series identities and associated generating-function relationships, *Appl. Math. Lett.*, Vol. 19, pp. 887–893.

Chen, K.-Y. and Srivastava, H. M. (2005). Series identities and associated families of generating functions, *J. Math. Anal. Appl.*, Vol. 311, pp. 582–599.

Choi, J. and Rathie, A. K. (2014). On the reducibility of Kampé de Fériet function, *Honam Mathematical J.*, Vol. 36, No. 2, pp. 345–355.

Chu, W. (2011). Terminating hypergeometric ${}_2F_1(2)$ series, *Integral Transforms Spec. Funct.*, Vol. 22, No. 2, pp. 91–96.

Clausen, T. (1828). Ueber die Fälle wenn die Reihe $y = 1 + \frac{\alpha.\beta}{1.\gamma}x + \dots$ Ein quadrat von der form $z = 1 + \frac{\alpha'\beta'\gamma'}{1.\delta'\epsilon'}x + \dots$ hat, *J. Reine Angew Math.*, Vol. 3, pp. 89–91.

Ebisu, A. (20 Mar 2017). On a strange evaluation of the hypergeometric series by Gosper. II, arXiv: 1608.04315v2 [math.CA].

Gauss, C. F. (1866). *Disquisitiones generales circa seriem infinitam*,

$$1 + \frac{\alpha.\beta}{1.\gamma}x + \frac{\alpha(\alpha+1)\beta(\beta+1)}{1.2.\gamma(\gamma+1)}x^2 + \frac{\alpha(\alpha+1)(\alpha+2)\beta(\beta+1)(\beta+2)}{1.2.3.\gamma(\gamma+1)(\gamma+2)}x^3 + \dots,$$

Gottingen Thesis, 1812, Published in ges. Werke Gottingen, Vol. II, pp. 437-445; III, pp. 123-163; III, pp. 207-229; III, pp. 446-460.

Hài, N. T., Marichev, O. I. and Srivastava, H. M. (1992). A note on the convergence of certain families of multiple hypergeometric series, *J. Math. Anal. Appl.*, Vol. 164, pp. 104–115.

Humbert, P. (1920-21). The confluent hypergeometric functions of two variables, *Proc. Royal Soc. Edinburgh. Sec. A.*, Vol. 41, pp. 73–96.

Kampé de Fériet, J. (1921). Les fonctions hypergéométriques d'ordre supérieur à deux variables, *C. R. Acad. Sci. Paris*, Vol. 173, pp. 401–404.

Kim, Y. S. (2009). On certain reducibility of Kampé de Fériet functions, *Honam Math. J.*, Vol. 31, No. 2, pp. 167–176.

Kim, Y. S., Choi, J. and Rathie A. K. (2012). Two results for the terminating ${}_3F_2(2)$ with applications, *Bull. Korean Math. Soc.*, Vol. 49, pp. 621–633.

Kim, Y. S., Gaboury, S. and Rathie A. K. (2018). Applications of extended Watson's summation theorem, *Turk. J. Math.*, Vol. 42, pp. 418–443.

Kim, Y. S., Rakha, M. A. and Rathie A. K. (2010). Generalization of Kummer second theorem with applications, *Comput. Math. and Math. Phys.*, Vol. 50, No. 3, pp. 387–402.

Kim, Y. S., Rathie, A. K. and Paris, R. B. (2018). Evaluations of some terminating hypergeometric ${}_2F_1(2)$ series with applications, *Turk. J. Math.*, Vol. 42, pp. 2563–2575.

Kummer, E. E. (1836). *Über die hypergeometrische Reihe*

$$1 + \frac{\alpha.\beta}{1.\gamma}x + \frac{\alpha(\alpha+1)\beta(\beta+1)}{1.2.\gamma(\gamma+1)}x^2 + \frac{\alpha(\alpha+1)(\alpha+2)\beta(\beta+1)(\beta+2)}{1.2.3.\gamma(\gamma+1)(\gamma+2)}x^3 + \dots,$$

J. für die Reine und Angewandte Math., Vol. 15, pp. 39–83 and pp. 127–172.

Lauricella, G. (1893). Sulle funzioni ipergeometriche a più variabili, *Rend. Circ. Mat. Palermo*, Vol. 7, pp. 111–158.

Lavoie, J. L., Grondin, F. and Rathie, A. K. (1992). Generalizations of Watson's theorem on the sum of a ${}_3F_2$, *Indian J. Math.*, Vol. 34, No. 1, pp. 23–32.

- Lavoie, J. L., Grondin, F. and Rathie, A. K. (1996). Generalizations of Whipple's theorem on the sum of a ${}_3F_2$, *J. Comput. Appl. Math.*, Vol. 72, No. 2, pp. 293–300.
- Lavoie, J. L., Grondin, F., Rathie, A. K. and Arora, K. (1994). Generalizations of Dixon's theorem on the sum of a ${}_3F_2$, *Math. Comp.*, Vol. 62, No. 205, pp. 267–276.
- Meethal, S., Rathie, A. K. and Paris, R. B. (2015). A derivation of two quadratic transformations contiguous to that of Gauss via a differential equation approach, *Applied Mathematical Sciences*, Vol. 9, No. 17, pp. 845–851.
- Miller, A. R. (2009). Certain summation and transformation formulas for generalized hypergeometric series, *J. Comput. Appl. Math.*, Vol. 231, pp. 964–972.
- Miller, A. R. and Paris, R. B. (2011a). Certain transformations and summations for the generalized hypergeometric series with integral parameter differences, *Integral Transforms and Special Functions*, Vol. 22, pp. 67–77.
- Miller, A. R. and Paris, R. B. (2011b). Euler-type transformations for the generalized hypergeometric function ${}_{r+2}F_{r+1}(x)$, *Zeit. Angew. Math. Phys.*, Vol. 62, pp. 31–45.
- Miller, A. R. and Paris, R. B. (2013). Transformation formulas for the generalized hypergeometric function with integral parameter differences, *Rocky Mountain J. Math.*, Vol. 43, pp. 291–327.
- Miller, A. R. and Srivastava, H. M. (2010). Karlsson Minton summation theorems for the generalized hypergeometric series of unit argument, *Integral Transforms and Special Functions*, Vol. 21, pp. 603–612.
- Prudnikov, A. P., Brychkov, Yu. A. and Marichev, O. I. (1990). *Integrals and Series, Vol. 3: More Special Functions*, Nauka, Moscow, 1986 (In Russian); (Translated from the Russian by G. G. Gould), Gordon and Breach Science Publishers, New York, Philadelphia, London, Paris, Montreux, Tokyo, Melbourne.
- Qureshi, M. I., Quraishi, K. A., Khan, B. and Singh, R. (2016). Transformation formulae for ex-ton's quadruple hypergeometric functions, *South Asian Journal of Mathematics*, Vol. 6, No. 1, pp. 31–37.
- Rainville, E. D. (1960). *Special Functions*, The Macmillan Co. Inc., New York.
- Rakha, M. A and Rathie, A. K. (2011). Generalizations of classical summation theorems for the series ${}_2F_1$ and ${}_3F_2$ with Applications, *Integral Transforms and Special Functions*, Vol. 22, No. 11, pp. 823–840.
- Rathie, A. K. and Nagar, V. (1995). On Kummer's second theorem involving product of generalized hypergeometric series, *Matematiche (Catania)*, Vol. 50, No. 1, pp. 35–38.
- Srivastava, H. M. (2016). Some families of generating functions and associated hypergeometric transformation and reduction formulas, *Russian Journal of Mathematical Physics*, Vol. 23, No. 3, pp. 382–391.
- Srivastava, H. M. and Manocha, H. L. (1984). *A Treatise on Generating Functions*, Halsted Press (Ellis Horwood Limited, Chichester, U.K.) John Wiley and Sons, New York, Chichester, Brisbane and Toronto.
- Srivastava, H. M. and Panda, R. (1976). An integral representation for the product of two Jacobi polynomials, *J. London Math. Soc.*, Vol. 12, No. 2, pp. 419–425.
- Srivastava, H. M. and Pathan, M. A. (1979). Some bilateral generating functions for the extended Jacobi polynomial-I, *Comment. Math. Uni. St. Paul.*, Vol. 28, No. 1, pp. 23–30.