



4-2020

Class of Integrals Involving Generalized Hypergeometric Function

D. L. Suthar
Wollo University

Teklay Hailay
Wollo University

Hafta Amsalu
Wollo University

Jagdev Singh
JECRC University

Follow this and additional works at: <https://digitalcommons.pvamu.edu/aam>



Part of the [Analysis Commons](#), and the [Special Functions Commons](#)

Recommended Citation

Suthar, D. L.; Hailay, Teklay; Amsalu, Hafta; and Singh, Jagdev (2020). Class of Integrals Involving Generalized Hypergeometric Function, *Applications and Applied Mathematics: An International Journal (AAM)*, Vol. 15, Iss. 3, Article 7.

Available at: <https://digitalcommons.pvamu.edu/aam/vol15/iss3/7>

This Article is brought to you for free and open access by Digital Commons @PVAMU. It has been accepted for inclusion in *Applications and Applied Mathematics: An International Journal (AAM)* by an authorized editor of Digital Commons @PVAMU. For more information, please contact hvkoshy@pvamu.edu.



Class of Integrals Involving Generalized Hypergeometric Function

¹D. L. Suthar, ²Teklay Hailay, ³Hafte Amsalu and ^{4*}Jagdev Singh

^{1,2,3}Department of Mathematics, Wollo University
P.O. Box 1145, Dessie, Ethiopia

¹dlsuthar@gmail.com; ²teklayhailay@gmail.com;
³yohanahafe@gmail.com

⁴Department of Mathematics, JECRC University
Jaipur-303905, Rajasthan, India

⁴jagdevsinghrathore@gmail.com

*Corresponding Author

Received: February 29, 2020; Accepted: March 15, 2020

Abstract

In this paper, we establish some definite integrals involving generalized hypergeometric function, product of algebraic functions, Jacobi function, Legendre function and general class of polynomials. Certain special cases of the main results are also pointed out.

Keywords: Generalized hypergeometric functions; Srivastava's polynomials; Definite integrals

MSC 2010 No.: 26A33, 33C45, 33C60

1. Introduction

In past few decades, many researchers are attracted to diverse techniques of special functions and their uses in many other areas of mathematics. These functions as a part of the theory of hypergeometric functions, are important special functions and they are closely related to physics and engineering applications. The great applications in a wide variety of fields (Srivastava et al. (2014)) have played a pivotal role in the advancements of further research in special functions.

Several further properties of each of generalized incomplete hypergeometric functions and some classes of incomplete H-functions associated with them can be found in the subsequent developments presented in recent articles (Bansal et al. (2019); Bansal et al. (2020)). In view of their importance and popularity in recent years, the theory of operators of fractional calculus have been developed widely and extensively by many researchers (Chaurasia and Singh (2015); Suthar, Parmar et al. (2018)) and various generalizations associated with integrals have been studied (Chaurasia and Singh (2014); Kumar et al. (2018); Khan (2017); Nisar et al. (2017); Sarbia and Kalla (2002); Suthar et al. (2017); Suthar, Agarwal et al. (2018)).

2. Preliminaries

For our investigation, we need extended generalized hypergeometric function (Srivastava et al. (2014)) defined as:

$${}_rF_s [x] = {}_rF_s \left[\begin{matrix} (\varsigma_1, p), \varsigma_2, \dots, \varsigma_r; \\ \vartheta_1, \dots, \vartheta_s; \end{matrix} x \right] = \sum_{n=0}^{\infty} \frac{(\varsigma_1; p)_n (\varsigma_2)_n \dots (\varsigma_r)_n (x)^n}{(\vartheta_1)_n \dots (\vartheta_s)_n n!}, \quad (1)$$

where $(\lambda; p)_n$ denotes the Pochhammer symbol (see (Srivastava et al. (2014)), p. 485 Eq. (1.66)) and is defined as follow:

$$(\lambda; p)_n = \begin{cases} \frac{\Gamma_p(\lambda+n)}{\Gamma_p(\lambda)}, & (\Re(p) > 0; \lambda, n \in \mathbb{C}), \\ (\lambda)_p, & (p = 0; \lambda, n \in \mathbb{C}). \end{cases} \quad (2)$$

The generalized gamma function (Chaudhary and Zubair (2001), p. 9, Eq (1.66)) is defined as:

$$\Gamma_p(z) = \begin{cases} \int_0^{\infty} x^{z-1} \exp(-x - p/x) dx, & (\Re(p) > 0; z \in \mathbb{C}), \\ \Gamma(z), & (p = 0; \Re(z) > 0). \end{cases} \quad (3)$$

The analogous extensions of the Gauss hypergeometric and the confluent hypergeometric functions are represented as follows:

$${}_2F_1 \left[\begin{matrix} (\varsigma, p), \vartheta; \\ \gamma; \end{matrix} x \right] = \sum_{n=0}^{\infty} \frac{(\varsigma, p)_n (\vartheta)_n (x)^n}{(\gamma)_n n!}, \quad (4)$$

and

$${}_1F_1 \left[\begin{matrix} (\varsigma, p); \\ \gamma; \end{matrix} x \right] = \sum_{n=0}^{\infty} \frac{(\varsigma, p)_n (x)^n}{(\gamma)_n n!}. \quad (5)$$

For the current investigation, we need the following definitions.

Definition 2.1.

The Gamma function (Rainville (1960)) can be defined as:

$$\Gamma(\lambda) = \int_0^{\infty} e^{-t} t^{\lambda-1} dt, \quad \Re(\lambda > 0). \quad (6)$$

Definition 2.2.

The Beta function is defined (Olver et al. (2010)) as:

$$B(u, v) = \int_0^1 t^{u-1} (1-t)^{v-1} dt = \frac{\Gamma(u) \Gamma(v)}{\Gamma(u+v)} = B(v, u), \quad (u, v \in \Re). \quad (7)$$

Definition 2.3.

The Jacobi polynomial $P_n^{(\tau, v)}(x)$ (Rainville (1960), p. 254) is defined as:

$$P_n^{(\tau, v)}(x) = \frac{(\tau+1)_n}{n!} {}_2F_1 \left[\begin{matrix} -n, 1+\tau+v+n; \\ 1+\tau; \end{matrix} \frac{1-x}{2} \right], \quad (8)$$

where ${}_2F_1$ is the classical hypergeometric functions defined as:

$${}_2F_1 \left[\begin{matrix} -n, 1+\tau+v+n; \\ 1+\tau; \end{matrix} \frac{1-x}{2} \right] = \sum_{l=0}^{\infty} \frac{(-n)_l (1+\tau+v+n)_l (1-x)^l}{(1+\tau)_l 2^l l!}.$$

When $\tau = v = 0$, then the polynomial (8) becomes the Legendre polynomial (Rainville (1960), p. 157) and also, we have

$$P_n^{(\tau, v)}(1) = \frac{(\tau+1)_n}{n!}. \quad (9)$$

Definition 2.4.

The solution of Legendre equation (Rainville (1960), Sec. 3.1)

$$(1-x^2) \frac{d^2 f}{dx^2} - 2x \frac{df}{dx} + [\varepsilon(\varepsilon+1) - \tau^2(1-x^2)^{-1}] f = 0, \quad (10)$$

is presented in the form of the Legendre function. If we put $f = (x^2 - 1)^{\frac{1}{2r}} \varepsilon$, then Equation (10) reduced to

$$(1-x^2) \frac{d^2 \varepsilon}{dx^2} - 2(\tau+1)x \frac{d\varepsilon}{dx} + [\varepsilon(\tau-\varepsilon)(\tau+\varepsilon+1)^{-1}] = 0, \quad (11)$$

and if we set $\delta = \frac{1}{2} - \frac{1}{2}x$ as the independent variable, the above differential equation (10) becomes

as follows:

$$\delta(1-\delta)\frac{d^2\varepsilon}{d\delta^2} + (\tau+1)(1-2\delta)\frac{d\varepsilon}{d\delta} + [\varepsilon(\varepsilon-\tau)(\tau+\varepsilon+1)] = 0. \quad (12)$$

The solution of Equation (10) in the representation of Gauss hypergeometric equation with $a = \tau - \varepsilon$, $b = \tau + \varepsilon + 1$ and $c = \tau + 1$ is as follows:

$$f = P_\varepsilon^\tau(x) = \frac{1}{\Gamma(1-\tau)} \left(\frac{z+1}{z-1}\right)^{\frac{1}{2\tau}} F\left[-\varepsilon, \varepsilon+1; 1-\tau; \frac{1}{2} - \frac{1}{2}x\right], \quad |1-x| < 2, \quad (13)$$

where $P_\varepsilon^\tau(x)$ is well-known Legendre function of the first kind (see Erdélyi et al. (1953)).

Definition 2.5.

The general class of polynomials $S_{n_1, \dots, n_r}^{m_1, \dots, m_r}[x]$ (Srivastava (1985), p. 185, Eq. (7)) is presented in the following manner:

$$S_{n_1, \dots, n_r}^{m_1, \dots, m_r}[x] = \sum_{l_1=0}^{\lfloor \frac{n_1}{m_1} \rfloor} \cdots \sum_{l_r=0}^{\lfloor \frac{n_r}{m_r} \rfloor} \prod_{i=1}^r \frac{(-n_i)_{m_i l_i}}{l_i!} A_{n_i, l_i} x^{l_i}, \quad (14)$$

where $n_1, \dots, n_r = 0, 1, 2, \dots$; m_1, \dots, m_r are arbitrary positive integers and the coefficient A_{n_i, l_i} ($n_i, l_i \geq 0$) are arbitrary constants, (real or complex). On suitably specializing the coefficients A_{n_i, l_i} , $S_{n_1, \dots, n_r}^{m_1, \dots, m_r}[x]$ yields a number of known polynomials as its special cases. These includes Laguerre polynomials, Bessel polynomials, Jacobi polynomials, Hermite polynomials, Brafman polynomials, Gould-Hopper polynomials and several others (Srivastava and Singh (1983), pp. 158-161).

The objective of this paper is to establish some definite integrals involving generalized hypergeometric function, product of algebraic functions, Jacobi function, Legendre function and general class of polynomials.

During the course of this paper, let \mathbb{C} be sets of complex numbers, \mathbb{R} - real, \mathbb{R}^+ -positive numbers, \mathbb{Z}_0^- - non-positive and \mathbb{N} -positive integers.

3. Integrals involving generalized hypergeometric functions with algebraic function

In this section, we evaluate the following integral formulas involving generalized hypergeometric function with some algebraic functions:

$$\begin{aligned}
I_1 &= \int_0^1 x^{-\xi}(1-x)^{\xi-\omega-1} {}_rF_s [zx] dx \\
&= \int_0^1 x^{-\xi}(1-x)^{\xi-\omega-1} \sum_{n=0}^{\infty} \frac{(\varsigma_1, p)_n (\varsigma_2)_n \dots (\varsigma_r)_n (zx)^n}{(\vartheta_1)_n \dots (\vartheta_s)_n n!} dx \\
&= \sum_{n=0}^{\infty} \frac{(\varsigma_1, p)_n (\varsigma_2)_n \dots (\varsigma_r)_n z^n}{(\vartheta_1)_n \dots (\vartheta_s)_n n!} \int_0^1 x^{n-\xi}(1-x)^{\xi-\omega-1} dx \\
&= \sum_{n=0}^{\infty} \frac{(\varsigma_1, p)_n (\varsigma_2)_n \dots (\varsigma_r)_n z^n}{(\vartheta_1)_n \dots (\vartheta_s)_n n!} \frac{\Gamma(n-\xi+1) \Gamma(\xi-\omega)}{\Gamma(n-\omega+1)} \\
&= B(1-\xi, \xi-\omega) {}_{r+1}F_{s+1} \left[\begin{matrix} (\varsigma_1, p), \varsigma_2, \dots, \varsigma_r, (1-\xi) \\ \vartheta_1, \dots, \vartheta_s, (1-\omega) \end{matrix}; z \right]. \tag{15}
\end{aligned}$$

$$\begin{aligned}
I_2 &= \int_0^1 x^{\xi-1}(1-x)^{\omega-1} {}_rF_s [zx] dx \\
&= \int_0^1 x^{\xi-1}(1-x)^{\omega-1} \sum_{n=0}^{\infty} \frac{(\varsigma_1, p)_n (\varsigma_2)_n \dots (\varsigma_r)_n (zx)^n}{(\vartheta_1)_n \dots (\vartheta_s)_n n!} dx \\
&= \sum_{n=0}^{\infty} \frac{(\varsigma_1, p)_n (\varsigma_2)_n \dots (\varsigma_r)_n z^n}{(\vartheta_1)_n \dots (\vartheta_s)_n n!} \int_0^1 x^{n+\xi-1}(1-x)^{\omega-1} dx \\
&= \sum_{n=0}^{\infty} \frac{(\varsigma_1, p)_n (\varsigma_2)_n \dots (\varsigma_r)_n z^n}{(\vartheta_1)_n \dots (\vartheta_s)_n n!} \frac{\Gamma(\omega) \Gamma(n+\xi)}{\Gamma(\omega+\xi+n)}, \\
&= B(\xi, \omega) {}_{r+1}F_{s+1} \left[\begin{matrix} (\varsigma_1, p), \varsigma_2, \dots, \varsigma_r, \xi \\ \vartheta_1, \dots, \vartheta_s, \omega+\xi \end{matrix}; z \right]. \tag{16}
\end{aligned}$$

$$\begin{aligned}
I_3 &= \int_1^{\infty} x^{-\xi}(x-1)^{\omega-1} {}_rF_s \left[\frac{z}{x} \right] dx \\
&= \sum_{n=0}^{\infty} \frac{(\varsigma_1, p)_n (\varsigma_2)_n \dots (\varsigma_r)_n (\xi)_n z^n}{(\vartheta_1)_n \dots (\vartheta_s)_n (\omega+\xi)_n n!} \int_1^{\infty} x^{-\xi-n}(x-1)^{\omega-1} dx.
\end{aligned}$$

By letting $x - 1 = t$ and taking as $x \rightarrow \infty, t \rightarrow \infty$ and using the formula

$$\Gamma(\varsigma)\Gamma(\vartheta) = \Gamma(\varsigma + \vartheta) \int_0^\infty x^{\varsigma-1}(1+x)^{-(\varsigma+\vartheta)} dx, \quad (17)$$

we get the following result

$$\begin{aligned} I_3 &= \sum_{n=0}^{\infty} \frac{(\varsigma_1, p)_n (\varsigma_2)_n \dots (\varsigma_r)_n z^n}{(\vartheta_1)_n \dots (\vartheta_s)_n} \frac{z^n}{n!} \int_0^\infty t^{\omega-1} (1+t)^{-\xi-n} dt \\ &= \sum_{n=0}^{\infty} \frac{(\varsigma_1, p)_n (\varsigma_2)_n \dots (\varsigma_r)_n z^n}{(\vartheta_1)_n \dots (\vartheta_s)_n} \frac{z^n}{n!} \frac{\Gamma(\omega) \Gamma(\xi - \omega + n)}{\Gamma(\xi + n)} \\ &= \frac{\Gamma(\omega) \Gamma(\xi - \omega)}{\Gamma(\xi)} \sum_{n=0}^{\infty} \frac{(\varsigma_1, p)_n (\varsigma_2)_n \dots (\varsigma_r)_n (\xi - \omega)_n z^n}{(\vartheta_1)_n \dots (\vartheta_s)_n (\xi)_n} \frac{z^n}{n!} \\ &= B(\omega, \xi - \omega) {}_{r+1}F_{s+1} \left[\begin{matrix} (\varsigma_1, p), \varsigma_2, \dots, \varsigma_r, \xi - \omega; \\ \vartheta_1, \dots, \vartheta_s, \xi; \end{matrix} z \right]. \end{aligned} \quad (18)$$

$$\begin{aligned} I_4 &= \int_{-1}^1 (1-x)^\xi (1+x)^\omega {}_rF_s [z(1-x)^\tau] dx \\ &= \sum_{n=0}^{\infty} \frac{(\varsigma_1, p)_n (\varsigma_2)_n \dots (\varsigma_r)_n z^n}{(\vartheta_1)_n \dots (\vartheta_s)_n} \frac{z^n}{n!} \int_{-1}^1 (1-x)^{\xi+\tau n+1-1} (1+x)^{\omega+1-1} dx. \end{aligned}$$

By using the formula (Rainville (1960), p. 26)

$$\int_{-1}^1 (1-x)^{\xi+n} (1+x)^{\omega+n} dx = 2^{2n+\xi+\omega+1} B(\xi+n+1, \omega+n+1), \quad (19)$$

we evaluate the following integral:

$$\begin{aligned} I_4 &= \sum_{n=0}^{\infty} \frac{(\varsigma_1, p)_n (\varsigma_2)_n \dots (\varsigma_r)_n z^n}{(\vartheta_1)_n \dots (\vartheta_s)_n} \frac{z^n}{n!} 2^{\tau n + \omega + \xi + 1} B(1 + \xi + \tau n, 1 + \omega) \\ &= 2^{\omega + \xi + 1} \Gamma(1 + \omega) \sum_{n=0}^{\infty} \frac{(\varsigma_1, p)_n (\varsigma_2)_n \dots (\varsigma_r)_n}{(\vartheta_1)_n \dots (\vartheta_s)_n} \frac{\Gamma(1 + \xi + \tau n)}{\Gamma(\xi + \omega + 2 + \tau n)} \frac{(2^\tau z)^n}{n!}. \end{aligned} \quad (20)$$

In particular, if $\tau = 1$, then Equation (20) becomes

$$\begin{aligned} I_4 &= 2^{\omega+\xi+1} \frac{\Gamma(1+\omega)\Gamma(1+\xi)}{\Gamma(\xi+\omega+2)} \sum_{n=0}^{\infty} \frac{(\varsigma_1, p)_n (\varsigma_2)_n \dots (\varsigma_r)_n (1+\xi)_n (2z)^n}{(\vartheta_1)_n \dots (\vartheta_s)_n (\xi+\omega+2)_n n!} \\ &= 2^{\omega+\xi+1} \frac{\Gamma(1+\omega)\Gamma(1+\xi)}{\Gamma(\xi+\omega+2)} {}_{r+1}F_{s+1} \left[\begin{matrix} (\varsigma_1, p), \varsigma_2, \dots, \varsigma_r, 1+\xi; \\ \vartheta_1, \dots, \vartheta_s, \xi+\omega+2; \end{matrix} 2z \right]. \end{aligned} \quad (21)$$

4. Integrals involving generalized hypergeometric functions with Jacobi polynomials

In this section, we derive the following integral formulas involving generalized hypergeometric functions multiplied with Jacobi Polynomials:

$$\begin{aligned} I_5 &= \int_{-1}^1 x^\lambda (1-x)^\tau (1+x)^\mu P_n^{(\tau, \nu)}(x) {}_rF_s \left[z(x+1)^l \right] dx \\ &= \sum_{n=0}^{\infty} \frac{(\varsigma_1, p)_n (\varsigma_2)_n \dots (\varsigma_r)_n z^n}{(\vartheta_1)_n \dots (\vartheta_s)_n n!} \int_{-1}^1 x^\lambda (1-x)^\tau (1+x)^{\mu+nl} P_n^{(\tau, \nu)}(x) dx. \end{aligned} \quad (22)$$

Next using the following formula

$$\begin{aligned} &\int_{-1}^1 x^\lambda (1-x)^\varsigma (1+x)^\mu P_n^{(\varsigma, \vartheta)}(x) dx \\ &= \frac{(-1)^n 2^{\varsigma+\mu+1} \Gamma(\mu+1) \Gamma(n+\varsigma+1) \Gamma(\mu+\vartheta+1)}{n! \Gamma(\mu+\vartheta+n+1) \Gamma(\mu+\varsigma+n+2)} \\ &\quad \times {}_3F_2 \left[\begin{matrix} -\lambda, \tau+\vartheta+1, \tau+1; \\ \tau+\vartheta+n+1, \tau+\varsigma+n+2; \end{matrix} 1 \right], \end{aligned} \quad (23)$$

where $\varsigma > -1$ and $\vartheta > -1$. Also ${}_3F_2$ is the special case of generalized hypergeometric series. Then using Equation (22) in (23), we have

$$\begin{aligned} I_5 &= \sum_{n=0}^{\infty} \frac{(\varsigma_1, p)_n (\varsigma_2)_n \dots (\varsigma_r)_n z^n}{(\vartheta_1)_n \dots (\vartheta_s)_n n!} \\ &\quad \times \frac{(-1)^n 2^{\tau+\mu+nl+1} \Gamma(\mu+nl+1) \Gamma(n+\tau+1) \Gamma(\mu+nl+\nu+1)}{n! \Gamma(\mu+nl+\nu+n+1) \Gamma(\mu+nl+\tau+n+2)} \\ &\quad \times {}_3F_2 \left[\begin{matrix} -\lambda, \mu+nl+\nu+1, \mu+nl+1; \\ \mu+nl+\nu+n+1, \mu+nl+\tau+n+2; \end{matrix} 1 \right]. \end{aligned}$$

After some rearrangements, we get

$$\begin{aligned}
 I_5 &= 2^{\tau+\mu+1} \Gamma(\tau+1) {}_{r+1}F_s \left[\begin{matrix} (\varsigma_1, p), \varsigma_2, \dots, \varsigma_r, \tau+1; \\ \vartheta_1, \dots, \vartheta_s; \end{matrix} -2^l z \right] \\
 &\times \frac{\Gamma(\mu+nl+1) \Gamma(\mu+nl+v+1)}{n! \Gamma(\mu+n(l+1)+v+1) \Gamma(\mu+n(l+1)+\tau+2)} \\
 &\times {}_3F_3 \left[\begin{matrix} -\lambda, \mu+nl+v+1, \mu+nl+1; \\ \mu+nl+v+n+1, \mu+nl+\tau+n+2, 1; \end{matrix} 1 \right]. \tag{24}
 \end{aligned}$$

From Equation (24), we can justify the special case.

If $l = 1$, then using the basic property $(\varsigma+1)_{2n} = \left(\frac{\varsigma+1}{2}\right)_n \left(\frac{\varsigma+2}{2}\right)_n$, we get

$$\begin{aligned}
 I_5 &= 2^{\tau+\mu+1} \Gamma(\tau+1) {}_{r+1}F_s \left[\begin{matrix} (\varsigma_1, p), \varsigma_2, \dots, \varsigma_r, \tau+1; \\ \vartheta_1, \dots, \vartheta_s; \end{matrix} -2z \right] \\
 &\times {}_3F_3 \left[\begin{matrix} -\lambda, \mu+n+v+1, \mu+n+1; \\ \mu+2n+v+1, \mu+n+\tau+n+2, 1; \end{matrix} 1 \right] \\
 &\times \frac{(\mu+1)_n \Gamma(\mu+1) (\mu+v+1)_n \Gamma(\mu+v+1)}{\left(\frac{\mu+v+1}{2}\right)_n \left(\frac{\mu+v+2}{2}\right)_n \Gamma(\mu+v+1) \left(\frac{\mu+\tau+2}{2}\right)_n \left(\frac{\mu+\tau+3}{2}\right)_n \Gamma(\mu+\tau+2)} \\
 &= 2^{\tau+\mu+1} \Gamma(\tau+1) \frac{\Gamma(\mu+1) \Gamma(\mu+v+1)}{\Gamma(\mu+v+1) \Gamma(\mu+\tau+2)} \\
 &\times {}_{r+3}F_{s+4} \left[\begin{matrix} (\varsigma_1, p), \varsigma_2, \dots, \varsigma_r, \tau+1, \mu+1, \mu+v+1; \\ \vartheta_1, \dots, \vartheta_s, \frac{\mu+v+1}{2}, \frac{\mu+v+2}{2}, \frac{\mu+\tau+2}{2}, \frac{\mu+\tau+3}{2}; \end{matrix} -2z \right] \\
 &\times {}_3F_3 \left[\begin{matrix} -\lambda, \mu+n+v+1, \mu+n+1; \\ \mu+2n+v+1, \mu+n+\tau+n+2, 1; \end{matrix} 1 \right]. \tag{25}
 \end{aligned}$$

$$\begin{aligned}
 I_6 &= \int_{-1}^1 (1-x)^\delta (1+x)^\nu P_n^{(\tau, \nu)}(x) {}_rF_s \left[z(1-x)^l \right] dx \\
 &= {}_rF_s \left[\begin{matrix} (\varsigma_1, p), \varsigma_2, \dots, \varsigma_r; \\ \vartheta_1, \dots, \vartheta_s; \end{matrix} z \right] \int_{-1}^1 (1-x)^{\delta+nl} (1+x)^\nu dx \\
 &\times \frac{(1+\tau)_n}{n!} {}_2F_1 \left[\begin{matrix} -n, 1+\tau+\nu+n; \\ 1+\tau; \end{matrix} \frac{1-x}{2} \right]
 \end{aligned}$$

$$\begin{aligned}
&= \sum_{n=0}^{\infty} \frac{(\varsigma_1, p)_n (\varsigma_2)_n \cdots (\varsigma_r)_n (1+\tau)_n}{(\vartheta_1)_n \cdots (\vartheta_s)_n} \frac{z^n}{n!} \sum_{k=0}^{\infty} \frac{(-n)_k (1+\tau+\nu+n)_k}{(1+\tau)_k 2^k k!} \\
&\quad \times \int_{-1}^1 (1-x)^{\delta+nl+k} (1+x)^\nu dx.
\end{aligned} \tag{26}$$

Using (19), we obtain the following result

$$\begin{aligned}
I_6 &= \sum_{n=0}^{\infty} \frac{(\varsigma_1, p)_n (\varsigma_2)_n \cdots (\varsigma_r)_n (1+\tau)_n}{(\vartheta_1)_n \cdots (\vartheta_s)_n} \frac{z^n}{n!} \sum_{k=0}^{\infty} \frac{(-n)_k (1+\tau+\nu+n)_k (n+1)_k}{(1+\tau)_k 2^k k! (n+k)!} \\
&\quad \times 2^{\delta+nl+k+\nu+1} \frac{\Gamma(1+\delta+nl+k) \Gamma(1+\nu)}{\Gamma(\delta+nl+k+\nu+2)} \\
&= 2^{\delta+\nu+1} \Gamma(1+\nu) {}_{r+1}F_s \left[\begin{matrix} (\varsigma_1, p), \varsigma_2, \dots, \varsigma_r, 1+\tau; \\ \vartheta_1, \dots, \vartheta_s; \end{matrix} 2^l z \right] \\
&\quad \times \sum_{k=0}^{\infty} \frac{(-n)_k (1+\tau+\nu+n)_k (n+1)_k (1+\delta+nl)_k}{(1+\tau)_k k! (n+k)! (\delta+nl+\nu+2)_k} \frac{\Gamma(1+\delta+nl)}{\Gamma(\delta+nl+\nu+2)}.
\end{aligned} \tag{27}$$

By substituting $(n+k)! = (n+1)_k n! = (n+1)_k (1)_n$ into Equation (27), we get

$$\begin{aligned}
I_6 &= 2^{\delta+\nu+1} \Gamma(1+\nu) {}_{r+1}F_s \left[\begin{matrix} (\varsigma_1, p), \varsigma_2, \dots, \varsigma_r, 1+\tau; \\ \vartheta_1, \dots, \vartheta_s; \end{matrix} 2^l z \right] \\
&\quad \times \sum_{k=0}^{\infty} \frac{(-n)_k (1+\tau+\nu+n)_k (n+1)_k (1+\delta+nl)_k 1^k}{(1+\tau)_k (n+1)_k (1)_n (\delta+nl+\nu+2)_k k!} \frac{\Gamma(1+\delta+nl)}{\Gamma(\delta+nl+\nu+2)} \\
&= 2^{\delta+\nu+1} \Gamma(1+\nu) {}_{r+1}F_{s+1} \left[\begin{matrix} (\varsigma_1, p), \varsigma_2, \dots, \varsigma_r, 1+\tau; \\ \vartheta_1, \dots, \vartheta_s, 1; \end{matrix} z 2^l \right] \\
&\quad \times {}_3F_2 \left[\begin{matrix} -n, 1+\tau+\nu+n, 1+\delta+nl; \\ 1+\tau, \delta+nl+\nu+2; \end{matrix} 1 \right] \frac{\Gamma(1+\delta+nl)}{\Gamma(\delta+nl+\nu+2)}.
\end{aligned} \tag{28}$$

In particular, if $l = 1$, then

$$\begin{aligned}
I_6 &= 2^{\delta+\nu+1} B(1+\nu, 1+\delta) {}_{r+2}F_{s+2} \left[\begin{matrix} (\varsigma_1, p), \varsigma_2, \dots, \varsigma_r, 1+\tau, 1+\delta; \\ \vartheta_1, \dots, \vartheta_s, 1, \delta+\nu+2; \end{matrix} 2z \right] \\
&\quad \times {}_3F_2 \left[\begin{matrix} -n, 1+\tau+\nu+n, 1+\delta+n; \\ 1+\tau, \delta+n+\nu+2; \end{matrix} 1 \right].
\end{aligned} \tag{29}$$

$$\begin{aligned}
I_7 &= \int_{-1}^1 (1-x)^\xi (1+x)^\omega P_n^{(\tau, \nu)}(x) {}_rF_s \left[z(1-x)^l (1+x)^h \right] dx \\
&= \sum_{n=0}^{\infty} \frac{(\varsigma_1, p)_n (\varsigma_2)_n \dots (\varsigma_r)_n z^n}{(\vartheta_1)_n \dots (\vartheta_s)_n n!} \int_{-1}^1 (1-x)^{\xi+nl} (1+x)^{\omega+nh} P_n^{(\tau, \nu)} dx \\
&= {}_{r+1}F_s \left[\begin{matrix} (\varsigma_1, p), \varsigma_2, \dots, \varsigma_r, 1+\tau; \\ \vartheta_1, \dots, \vartheta_s; \end{matrix} z \right] \sum_{k=0}^{\infty} \frac{(-n)_k (1+\tau+\nu+n)_k}{(1+\tau)_k k! 2^k n!} \\
&\quad \times \int_{-1}^1 (1-x)^{\xi+nl+k} (1+x)^{\omega+nh} dx.
\end{aligned}$$

On using the result (19), we obtain

$$\begin{aligned}
I_7 &= {}_{r+1}F_s \left[\begin{matrix} (\varsigma_1, p), \varsigma_2, \dots, \varsigma_r, 1+\tau; \\ \vartheta_1, \dots, \vartheta_s; \end{matrix} z \right] \sum_{k=0}^{\infty} \frac{(-n)_k (1+\tau+\nu+n)_k}{(1+\tau)_k k! 2^k n!} \\
&\quad \times 2^{\xi+nl+k+\omega+nh+1} B(1+\xi+nl+k, 1+\omega+nh) \\
&= 2^{\xi+\omega+1} {}_{r+1}F_s \left[\begin{matrix} (\varsigma_1, p), \varsigma_2, \dots, \varsigma_r, 1+\tau; \\ \vartheta_1, \dots, \vartheta_s; \end{matrix} 2^{l+h} z \right] \sum_{k=0}^{\infty} \frac{(-n)_k (1+\tau+\nu+n)_k}{(1+\tau)_k n!} \frac{1}{k!} \\
&\quad \times \frac{\Gamma(1+\xi+nl+k) \Gamma(1+\omega+nh)}{\Gamma(\xi+n(l+h)+k+\omega+2)} \\
&= 2^{\xi+\omega+1} {}_{r+1}F_{s+1} \left[\begin{matrix} (\varsigma_1, p), \varsigma_2, \dots, \varsigma_r, 1+\tau; \\ \vartheta_1, \dots, \vartheta_s, 1; \end{matrix} 2^{l+h} z \right] \\
&\quad \times {}_3F_2 \left[\begin{matrix} -n, 1+\tau+\nu+n, 1+\xi+nl; \\ 1+\tau, \xi+n(l+h)+\omega+2; \end{matrix} 1 \right] \frac{\Gamma(1+\xi+nl) \Gamma(1+\omega+nh)}{\Gamma(\xi+n(l+h)+\omega+2)}. \tag{30}
\end{aligned}$$

For the particular value of $l = 1$ and $h = 1$, we arrive at the following result

$$\begin{aligned}
I_7 &= 2^{\xi+\omega+1} B(1+\xi, 1+\omega) {}_{r+3}F_{s+3} \left[\begin{matrix} (\varsigma_1, p), \varsigma_2, \dots, \varsigma_r, 1+\tau, 1+\xi, 1+\omega; \\ \vartheta_1, \dots, \vartheta_s, 1, \frac{\xi+\omega+2}{2}, \frac{\xi+\omega+3}{2}; \end{matrix} 4z \right] \\
&\quad \times {}_3F_2 \left[\begin{matrix} -n, 1+\tau+\nu+n, 1+\xi+n; \\ 1+\tau, \xi+2n+\omega+2; \end{matrix} 1 \right]. \tag{31}
\end{aligned}$$

$$I_8 = \int_{-1}^1 (1-x)^\xi (1+x)^\omega P_n^{(\tau, \nu)}(x) {}_rF_s \left[z(1+x)^{-l} \right] dx$$

$$\begin{aligned}
&= \sum_{n=0}^{\infty} \frac{(\varsigma_1, p)_n (\varsigma_2)_n \cdots (\varsigma_r)_n z^n}{(\vartheta_1)_n \cdots (\vartheta_s)_n} \frac{z^n}{n!} \int_{-1}^1 (1-x)^\xi (1+x)^{\omega-nl} P_n^{(\tau, \nu)}(x) dx, \\
&= {}_{r+1}F_s \left[\begin{matrix} (\varsigma_1, p), \varsigma_2, \dots, \varsigma_r, 1 + \tau; \\ \vartheta_1, \dots, \vartheta_s; \end{matrix} z \right] \sum_{k=0}^{\infty} \frac{(-n)_k (1 + \tau + \nu + n)_k}{(1 + \tau)_k k! n! 2^k} \\
&\quad \times \int_{-1}^1 (1-x)^{\xi+k} (1+x)^{\omega-nl} dx, \\
&= 2^{\xi+\omega+1} {}_{r+1}F_s \left[\begin{matrix} (\varsigma_1, p), \varsigma_2, \dots, \varsigma_r, 1 + \tau; \\ \vartheta_1, \dots, \vartheta_s; \end{matrix} 2^{-l} z \right] \\
&\quad \times \sum_{k=0}^{\infty} \frac{(-n)_k (1 + \tau + \nu + n)_k}{(1 + \tau)_k k! n!} \frac{\Gamma(1 + \xi + k) \Gamma(1 + \omega - nl)}{\Gamma(\xi + \omega + k - nl + 2)} \\
&= 2^{\xi+\omega+1} \Gamma(1 + \xi) {}_{r+1}F_{s+1} \left[\begin{matrix} (\varsigma_1, p), \varsigma_2, \dots, \varsigma_r, 1 + \tau; \\ \vartheta_1, \dots, \vartheta_s, 1; \end{matrix} 2^{-l} z \right] \\
&\quad \times \sum_{k=0}^{\infty} \frac{(-n)_k (1 + \tau + \nu + n)_k (1 + \xi)_k}{(1 + \tau)_k k!} \frac{\Gamma(1 + \omega - nl)}{\Gamma(\xi + \omega + k - nl + 2)} \\
&= 2^{\xi+\omega+1} \Gamma(1 + \xi) {}_{r+1}F_{s+1} \left[\begin{matrix} (\varsigma_1, p), \varsigma_2, \dots, \varsigma_r, 1 + \tau; \\ \vartheta_1, \dots, \vartheta_s, 1; \end{matrix} 2^{-l} z \right] \\
&\quad \times {}_3F_2 \left[\begin{matrix} -n, 1 + \tau + \nu + n, 1 + \xi; \\ 1 + \tau, \xi + \omega - nl + 2; \end{matrix} 1 \right] \frac{\Gamma(1 + \omega - nl)}{\Gamma(\xi + \omega - nl + 2)}. \tag{32}
\end{aligned}$$

However, if $l = -1$, then (32) becomes

$$\begin{aligned}
I_8 &= 2^{\xi+\omega+1} B(1 + \xi, 1 + \omega) {}_{r+2}F_{s+2} \left[\begin{matrix} (\varsigma_1, p), \varsigma_2, \dots, \varsigma_r, 1 + \tau, 1 + \omega; \\ \vartheta_1, \dots, \vartheta_s, 1, \xi + \omega + 2; \end{matrix} 2z \right] \\
&\quad \times {}_3F_2 \left[\begin{matrix} -n, 1 + \tau + \nu + n, 1 + \xi; \\ 1 + \tau, \xi + \omega + n + 2; \end{matrix} 1 \right]. \tag{33}
\end{aligned}$$

$$\begin{aligned}
I_9 &= \int_{-1}^1 (1-x)^\xi (1+x)^\omega P_n^{(\tau, \nu)}(x) {}_rF_s \left[z(1-x)^l (1+x)^{-h} \right] dx \\
&= \sum_{n=0}^{\infty} \frac{(\varsigma_1, p)_n (\varsigma_2)_n \cdots (\varsigma_r)_n z^n}{(\vartheta_1)_n \cdots (\vartheta_s)_n} \frac{z^n}{n!} \int_{-1}^1 (1-x)^{\xi+nl} (1+x)^{\omega-nh} P_n^{(\tau, \nu)}(x) dx \\
&= \sum_{n=0}^{\infty} \frac{(\varsigma_1, p)_n (\varsigma_2)_n \cdots (\varsigma_r)_n z^n}{(\vartheta_1)_n \cdots (\vartheta_s)_n} \frac{z^n}{n!} \frac{(1 + \tau)_n}{n!} \sum_{k=0}^{\infty} \frac{(-n)_k (1 + \tau + \nu + n)_k}{(1 + \tau)_k k! 2^k}
\end{aligned}$$

$$\begin{aligned} & \times \int_{-1}^1 (1-x)^{\xi+nl+k} (1+x)^{\omega-nh} dx \\ &= \sum_{n=0}^{\infty} \frac{(\varsigma_1, p)_n (\varsigma_2)_n \dots (\varsigma_r)_n (1+\tau)_n z^n}{(\vartheta_1)_n \dots (\vartheta_s)_n (1)_n n!} \sum_{k=0}^{\infty} \frac{(-n)_k (1+\tau+\nu+n)_k}{(1+\tau)_k k! 2^k} \\ & \times 2^{\xi+\omega+1+n(l-h)+k} \frac{\Gamma(\xi+nl+k+1)\Gamma(\omega-nh+1)}{\Gamma(\xi+\omega+nl-nh+k+2)} \\ &= 2^{\xi+\omega+1} {}_{r+1}F_{s+1} \left[\begin{matrix} (\varsigma_1, p), \varsigma_2, \dots, \varsigma_r, 1+\tau; \\ \vartheta_1, \dots, \vartheta_s, 1; \end{matrix} \middle| 2^{l-h} z \right] \\ & \times {}_3F_2 \left[\begin{matrix} -n, 1+\tau+\nu+n, \xi+nl+1; \\ 1+\tau, \xi+\omega+nl-nh+2; \end{matrix} \middle| 1 \right] \frac{\Gamma(\xi+nl+1)\Gamma(\omega-nh+1)}{\Gamma(\xi+\omega+nl-nh+2)}. \end{aligned}$$

If $l = 1$ and $h = -1$, then

$$\begin{aligned} I_9 &= 2^{\xi+\omega+1} B(\omega+1, \xi+1) {}_{r+3}F_{s+3} \left[\begin{matrix} (\varsigma_1, p), \varsigma_2, \dots, \varsigma_r, 1+\tau, \xi+1, \omega+1; \\ \vartheta_1, \dots, \vartheta_s, 1, ((\xi+\omega)/2)+1, ((\xi+\omega)/2)+1; \end{matrix} \middle| 4z \right] \\ & \times {}_3F_2 \left[\begin{matrix} -n, 1+\tau+\nu+n, \xi+n+1; \\ 1+\tau, \xi+\omega+2n+2; \end{matrix} \middle| 1 \right]. \end{aligned} \tag{34}$$

5. Integrals involving generalized hypergeometric functions with Legendre function

In this part, we derive integral formulas involving generalized hypergeometric functions multiplied with Legendre function.

$$\begin{aligned} I_{10} &= \int_0^1 x^{\omega-1} (1-x^2)^{\frac{\tau}{2}} P_{\nu}^{\tau}(x) {}_rF_s [zx^{\xi}] dx \\ &= \int_0^1 x^{\omega-1} (1-x^2)^{\frac{\tau}{2}} P_{\nu}^{\tau}(x) \sum_{n=0}^{\infty} \frac{(\varsigma_1, p)_n (\varsigma_2)_n \dots (\varsigma_r)_n z^n}{(\vartheta_1)_n \dots (\vartheta_s)_n n!} x^{n\xi} dx \\ &= \sum_{n=0}^{\infty} \frac{(\varsigma_1, p)_n (\varsigma_2)_n \dots (\varsigma_r)_n z^n}{(\vartheta_1)_n \dots (\vartheta_s)_n n!} \int_0^1 x^{\omega+n\xi-1} (1-x^2)^{\frac{\tau}{2}} P_{\nu}^{\tau}(x) dx. \end{aligned} \tag{35}$$

Using the formula (Erdélyi et al. (1953), Sec 3.12) for $\Re(\omega) > 0, \tau \in \mathbb{N}$

$$\int_0^1 x^{\omega-1} (1-x^2)^{\frac{\tau}{2}} P_{\nu}^{\tau}(x) dx = \frac{(-1)^{\tau} 2^{-\omega-\tau} \pi^{\frac{1}{2}} \Gamma(\omega) \Gamma(1+\tau+\nu)}{\Gamma(1-\tau+\nu) \Gamma\left(\frac{1}{2} + \frac{\omega}{2} + \frac{\tau-\nu}{2}\right) \Gamma\left(1 + \frac{\omega}{2} + \frac{\tau+\nu}{2}\right)}, \tag{36}$$

then the integral (35) becomes

$$\begin{aligned}
 I_{10} &= \sum_{n=0}^{\infty} \frac{(\varsigma_1, p)_n (\varsigma_2)_n \dots (\varsigma_r)_n z^n}{(\vartheta_1)_n \dots (\vartheta_s)_n} \frac{(-1)^\tau 2^{-\omega-\tau-n\xi} \pi^{\frac{1}{2}} \Gamma(\omega+n\xi) \Gamma(1+\tau+\nu)}{n! \Gamma(1-\tau+\nu) \Gamma\left(\frac{1+\omega+n\xi+\tau-\nu}{2}\right) \Gamma\left(1+\frac{\omega+n\xi+\tau+\nu}{2}\right)} \\
 &= \frac{(-1)^\tau 2^{-\omega-\tau} \pi^{\frac{1}{2}} \Gamma(1+\tau+\nu)}{\Gamma(1-\tau+\nu)} \sum_{n=0}^{\infty} \frac{(\varsigma_1, p)_n (\varsigma_2)_n \dots (\varsigma_r)_n (2^{-\xi} z)^n}{(\vartheta_1)_n \dots (\vartheta_s)_n} \frac{1}{n!} \\
 &\quad \times \frac{\Gamma(\omega+n\xi)}{\Gamma\left(\frac{1+\omega+n\xi+\tau-\nu}{2}\right) \Gamma\left(1+\frac{\omega+n\xi+\tau+\nu}{2}\right)}. \tag{37}
 \end{aligned}$$

In particular, if $\xi = 2$ in (37), then

$$\begin{aligned}
 I_{10} &= \frac{(-1)^\tau 2^{-\omega-\tau} \pi^{\frac{1}{2}} \Gamma(1+\tau+\nu) \Gamma(\omega)}{\Gamma(1-\tau+\nu) \Gamma\left(\frac{1+\omega+\tau-\nu}{2}\right) \Gamma\left(1+\frac{\omega+\tau+\nu}{2}\right)} \\
 &\quad \times {}_{r+2}F_{s+2} \left[\begin{matrix} (\varsigma_1, p), \varsigma_2, \dots, \varsigma_r, \frac{\omega}{2}, \frac{\omega+1}{2}; \\ \vartheta_1, \dots, \vartheta_s, \frac{1+\omega+\tau-\nu}{2}, 1+\frac{\omega+\tau+\nu}{2}; \end{matrix} \frac{z}{4} \right]. \tag{38}
 \end{aligned}$$

$$\begin{aligned}
 I_{11} &= \int_0^1 x^{\omega-1} (1-x^2)^{\frac{-\tau}{2}} P_\nu^\tau(x) {}_rF_s [zx^\xi] dx \\
 &= \sum_{n=0}^{\infty} \frac{(\varsigma_1, p)_n (\varsigma_2)_n \dots (\varsigma_r)_n z^n}{(\vartheta_1)_n \dots (\vartheta_s)_n} \frac{1}{n!} \int_0^1 x^{\omega+n\xi-1} (1-x^2)^{\frac{-\tau}{2}} P_\nu^\tau(x) dx.
 \end{aligned}$$

Using the formula (Erdélyi et al. (1953), Sec 3.12) for $\Re(\omega) > 0$, $\tau \in \mathbb{N}$,

$$\int_0^1 x^{\omega-1} (1-x^2)^{\frac{-\tau}{2}} P_\nu^\tau(x) dx = \frac{2^{\tau-\omega} \pi^{\frac{1}{2}} \Gamma(\omega)}{\Gamma\left(\frac{1+\omega-\tau-\nu}{2}\right) \Gamma\left(1+\frac{\omega-\tau-\nu}{2}\right)}, \tag{39}$$

we get

$$\begin{aligned}
 I_{11} &= \sum_{n=0}^{\infty} \frac{(\varsigma_1, p)_n (\varsigma_2)_n \dots (\varsigma_r)_n z^n}{(\vartheta_1)_n \dots (\vartheta_s)_n} \frac{1}{n!} \frac{2^{\tau-\omega-n\xi} \pi^{\frac{1}{2}} \Gamma(\omega+n\xi)}{\Gamma\left(\frac{1+\omega+n\xi-\tau-\nu}{2}\right) \Gamma\left(1+\frac{\omega+n\xi-\tau-\nu}{2}\right)} \\
 &= 2^{\tau-\omega} \pi^{\frac{1}{2}} {}_rF_s \left[\begin{matrix} (\varsigma_1, p), \varsigma_2, \dots, \varsigma_r; \\ \vartheta_1, \dots, \vartheta_s; \end{matrix} 2^{-\xi} z \right] \frac{\Gamma(\omega+n\xi)}{\Gamma\left(\frac{1+\omega+n\xi-\tau-\nu}{2}\right) \Gamma\left(1+\frac{\omega+n\xi-\tau-\nu}{2}\right)}.
 \end{aligned}$$

Particularly, if $\xi = 2$, then

$$I_{11} = 2^{\tau-\omega} \pi^{\frac{1}{2}} \frac{\Gamma(\omega)}{\Gamma\left(\frac{1+\omega-\tau-\nu}{2}\right) \Gamma\left(1+\frac{\omega-\tau-\nu}{2}\right)}$$

$$\times_{r+2}F_{s+2} \left[\begin{matrix} (\varsigma_1, p), \varsigma_2, \dots, \varsigma_r, \frac{\omega}{2}, \frac{\omega+1}{2}; \\ \vartheta_1, \dots, \vartheta_s, \frac{1+\omega-\tau-\nu}{2}, 1 + \frac{\omega-\tau-\nu}{2}; \end{matrix} \frac{z}{4} \right]. \quad (40)$$

6. Integrals involving generalized hypergeometric functions with general class of polynomials

From the basic definitions, we obtained the following results.

$$\begin{aligned} I_{12} &= \int_{-1}^1 (1-x)^{\xi-1} (1+x)^{\omega-1} S_{n_1, \dots, n_r}^{m_1, \dots, m_r} [z(1-x)^\tau (1+x)^\nu] \\ &\quad \times {}_rF_s [z(1-x)^h (1+x)^k] dx \\ &= \int_{-1}^1 (1-x)^{\xi-1} (1+x)^{\omega-1} \sum_{l_1=0}^{\lfloor \frac{n_1}{m_1} \rfloor} \dots \sum_{l_r=0}^{\lfloor \frac{n_r}{m_r} \rfloor} \prod_{i=1}^r \frac{(-n_i)_{m_i l_i}}{l_i!} A_{n_i, l_i} z^{l_i} (1-x)^{\tau l_i} \\ &\quad \times (1+x)^{\nu l_i} \sum_{n=0}^{\infty} \frac{(\varsigma_1, p)_n (\varsigma_2)_n \dots (\varsigma_r)_n z^n}{(\vartheta_1)_n \dots (\vartheta_s)_n n!} (1-x)^{nh} (1+x)^{nk} dx \\ &= \sum_{l_1=0}^{\lfloor \frac{n_1}{m_1} \rfloor} \dots \sum_{l_r=0}^{\lfloor \frac{n_r}{m_r} \rfloor} \prod_{i=1}^r \frac{(-n_i)_{m_i l_i}}{l_i!} A_{n_i, l_i} z^{l_i} \sum_{n=0}^{\infty} \frac{(\varsigma_1, p)_n (\varsigma_2)_n \dots (\varsigma_r)_n z^n}{(\vartheta_1)_n \dots (\vartheta_s)_n n!} \\ &\quad \times \int_{-1}^1 (1-x)^{\xi+\tau l_i+nh-1} (1+x)^{\omega+\nu l_i+nk-1} dx \end{aligned} \quad (41)$$

$$\begin{aligned} &= \sum_{l_1=0}^{\lfloor \frac{n_1}{m_1} \rfloor} \dots \sum_{l_r=0}^{\lfloor \frac{n_r}{m_r} \rfloor} \prod_{i=1}^r \frac{(-n_i)_{m_i l_i}}{l_i!} A_{n_i, l_i} z^{l_i} \sum_{n=0}^{\infty} \frac{(\varsigma_1, p)_n (\varsigma_2)_n \dots (\varsigma_r)_n z^n}{(\vartheta_1)_n \dots (\vartheta_s)_n n!} \\ &\quad \times 2^{\omega+\xi+(\tau+\nu)l_i+n(h+k)-1} \frac{\Gamma(\xi+\tau l_i+nh) \Gamma(\omega+\nu l_i+nk)}{\Gamma(\omega+\xi+(\tau+\nu)l_i+n(h+k))} \\ &= S_{n_1, \dots, n_r}^{m_1, \dots, m_r} [z] {}_rF_s \left[\begin{matrix} (\varsigma_1, p), \varsigma_2, \dots, \varsigma_r; \\ \vartheta_1, \dots, \vartheta_s; \end{matrix} z \right] 2^{\omega+\xi+(\tau+\nu)l_i+n(h+k)-1} \\ &\quad \times \frac{\Gamma(\xi+\tau l_i+nh) \Gamma(\omega+\nu l_i+nk)}{\Gamma(\omega+\xi+(\tau+\nu)l_i+n(h+k))}. \end{aligned} \quad (42)$$

In particular, if $h = k = 1$, then

$$\begin{aligned} I_{12} &= S_{n_1, \dots, n_r}^{m_1, \dots, m_r} [z] {}_{r+2}F_{s+2} \left[\begin{matrix} (\varsigma_1, p), \varsigma_2, \dots, \varsigma_r, \xi+\tau l_i, \omega+\nu l_i; \\ \vartheta_1, \dots, \vartheta_s, \frac{\omega+\xi+(\tau+\nu)l_i}{2}, \frac{\omega+\xi+(\tau+\nu)l_i+1}{2}; \end{matrix} 4z \right] \\ &\quad \times 2^{\omega+\xi+(\tau+\nu)l_i-1} B(\xi+\tau l_i, \omega+\nu l_i). \end{aligned} \quad (43)$$

7. Conclusion

We conclude here by remarking that the numerous further consequences of our results can easily be derived by using some known and new relationships between generalized hypergeometric functions, which is an elegant unification of various special functions such as: Gauss's hypergeometric and confluent hypergeometric functions, after some suitable parametric replacements. The results obtained here are basic in nature and are likely to find useful applications in the study of simple and multiple variable hypergeometric series which in turn are useful in statistical mechanics, electrical networks, and probability theory.

Acknowledgment:

The authors are thankful to the referees for their valuable remarks and comments for the improvement of the paper.

REFERENCES

- Bansal, M. K., Kumar, D., Khan, I., Singh J. and Nisar, K. S. (2019). Certain unified integrals associated with product of M-series and incomplete H-functions, *Mathematics*, Vol. 7, Art. No. 1191. doi:10.3390/math7121191
- Bansal, M. K., Kumar, D., Singh, J., Tassaddiq, A. and Nisar, K. S. (2020). Some new results for the Srivastava-Luo-Raina M-transform pertaining to the incomplete H-functions, *AIMS Mathematics*, Vol. 5, No. 1, pp. 717-722.
- Chaurasia, V. B. L. and Singh Y. (2014). Generalized elliptic-type integrals and generating functions, *Demonstration Mathematica*, Vol. 47, No. 2, pp. 310-323.
- Chaurasia, V. B. L. and Singh Y. (2015). Marichev-Saigo-Maeda fraction integration operators of certain special functions, *Gen. Math. Notes*, Vol. 26, No. 1, pp. 134-144.
- Chaudhary, M. A. and Zubair, S. M. (2001). *On a Class of Incomplete Gamma Functions with Applications*, Chapman and Hall, London.
- Erdélyi, A., Magnus, W., Oberhettinger, F. and Tricomi, F. G. (1953). *Higher Transcendental Functions*, Vol. I, McGraw-Hill, New York-Toronto-London.
- Khan, W. A. (2017). Some integrals involving Wright generalized Bessel-Maitland function with Jacobi polynomial, *Palestine J. Math.*, Vol. 6, No. 1, pp. 1-7.
- Kumar, D., Saxena R. K. and Ram J. (2018). Finite integral formulas involving aleph function, *Bol. Soc. Paran. Mat.*, Vol. 36, No. 1. pp. 177-193.
- Nisar, K. S., Suthar, D. L., Purohit, S. D. and Aldhaifallah, M. (2017). Some unified integral associated with the generalized Struve function, *Proc. Jangjeon Math. Soc.*, Vol. 20, No. 2, pp. 261-267.
- Olver, F. W. J., Lozier, D. W., Boisvert, R. F. and Clark, C. W. (2010). *NIST Handbook of Mathematical Functions*, Cambridge University Press, New York.
- Rainville, E. D. (1960). *Special Functions*, Macmillan Co., New York.

- Sarbia, J. and Kalla, S. L. (2002). On a generalized Jacobi transform, *J. Appl. Math. Stoch. Anal.*, Vol. 15, No. 4, pp. 371-384.
- Srivastava, H. M. (1985). A multilinear generating function for the Konhauser sets of biorthogonal polynomials suggested by the Laguerre polynomials, *Pacific J. Math.*, Vol. 117, No. 1, pp. 183-191.
- Srivastava, H. M., Cetinkaya, A. and Kiyamaz, I. O. (2014). A certain generalized Pochhammer symbol and its applications to hypergeometric functions, *Appl. Math. Comput.*, Vol. 226, pp. 484-491.
- Srivastava, H. M. and Singh N. P. (1983). The integration of certain of product multivariable H-function with a general class of polynomials, *Rend. Circ. Mat. Palermo Ser.*, Vol. 32, No. 2, pp. 157-187.
- Suthar, D. L., Agarwal, S. and Kumar D. (2019). Certain integrals involving the product of Gaussian hypergeometric function and Aleph function, *Honam Mathematical J.*, Vol. 41, No. 1, pp. 1-17.
- Suthar, D. L., Habenom, H. and Tadesse, H. (2017). Certain integrals involving Aleph function and Wright's generalized hypergeometric function, *Acta Univ. Apulensis Math. Inform.*, Vol. 52, pp. 01-10.
- Suthar, D. L., Parmar, R. K. and Purohit S. D. (2017). Fractional calculus with complex order and generalized hypergeometric functions, *Nonlinear Sci. Lett. A*, Vol. 8, No. 2, pp. 156-161.
- Suthar, D. L., Purohit, S. D. and Agarwal, S. (2017). Class of integrals involving generalized hypergeometric function and Srivastava's polynomials, *Int. J. Appl. Comput. Math.*, Vol. 3, No. 1, pp. 1197-1203.
- Suthar, D. L., Purohit, S. D. and Nisar, K. S. (2018). Certain integrals associated with generalized Hypergeometric functions, *Acta Univ. Apulensis Math. Inform.*, Vol. 55, pp. 105-112.
- Suthar, D. L., Tadesse, H. and Tilahun K. (2018). Integrals involving Jacobi polynomials and M-series, *J. Fractional Calc. Appl.*, Vol. 9, No. 2, pp. 287-294.