



4-2020

## Finite and Infinite Integral Formulas Involving the Family of Incomplete H-Functions

Manish K. Bansal  
*Government Engineering College*

Devendra Kumar  
*University of Rajasthan*

Jagdev Singh  
*JECRC University*

Kottakkaran Soopy Nisar  
*Prince Sattam bin Abdulaziz University*

Follow this and additional works at: <https://digitalcommons.pvamu.edu/aam>



Part of the [Special Functions Commons](#)

### Recommended Citation

Bansal, Manish K.; Kumar, Devendra; Singh, Jagdev; and Nisar, Kottakkaran Soopy (2020). Finite and Infinite Integral Formulas Involving the Family of Incomplete H-Functions, *Applications and Applied Mathematics: An International Journal (AAM)*, Vol. 15, Iss. 3, Article 2.

Available at: <https://digitalcommons.pvamu.edu/aam/vol15/iss3/2>

This Article is brought to you for free and open access by Digital Commons @PVAMU. It has been accepted for inclusion in *Applications and Applied Mathematics: An International Journal (AAM)* by an authorized editor of Digital Commons @PVAMU. For more information, please contact [hvkoshy@pvamu.edu](mailto:hvkoshy@pvamu.edu).



## Finite and Infinite Integral Formulas Involving the Family of Incomplete H-Functions

<sup>1</sup>Manish Kumar Bansal, <sup>2</sup>Devendra Kumar, <sup>3</sup>Jagdev Singh and <sup>4,\*</sup>Kottakkaran Soopy Nisar

<sup>1</sup>Department of Applied Sciences  
Government Engineering College  
Banswara-327001, Rajasthan, India  
[bansalmanish443@gmail.com](mailto:bansalmanish443@gmail.com)

<sup>2</sup>Department of Mathematics  
University of Rajasthan  
Jaipur 302004, Rajasthan, India  
[devendra.maths@gmail.com](mailto:devendra.maths@gmail.com)

<sup>3</sup>Department of Mathematics  
JECRC University  
Jaipur 303905, Rajasthan, India  
[jagdevsinghrathore@gmail.com](mailto:jagdevsinghrathore@gmail.com)

<sup>4,\*</sup>Department of Mathematics  
College of Arts and Sciences  
Prince Sattam bin Abdulaziz University  
Wadi Aldawaser 11991, Saudi Arabia  
[ksnisar1@gmail.com](mailto:ksnisar1@gmail.com)

\*Corresponding Author

Received: February 29, 2020; Accepted: March 15, 2020

### Abstract

Recent research focuses on the integral representations of the various type of special functions due to their potential applicability in different disciplines. In this line, we deal with several finite and infinite integrals involving the family of incomplete H-functions. Further, we point out some known and new special cases of these integrals. Finally, we establish the integral representation of incomplete H-functions.

**Keywords:** Incomplete  $H$ -functions; Gamma Function; Incomplete generalized hypergeometric function; Mellin-Barnes type integrals

**MSC 2010 No.:** 33C60, 33B15, 33D15, 33D70

## 1. Introduction

A large number of integral formulas involving the various kind of special functions have been established by several authors (Saxena et al. (2018); Suthar et al. (2019); Kumar et al. (2018); Bansal et al. (2019b); Bansal et al. (2019a); Bansal, Kumar, Khan et al. (2019)). In recent years, these integral formulas have many applications in the potential field of physics, applied sciences, engineering and chemical science. In this article, we deal with those finite and infinite integral formulas which are most general in nature.

## 2. Preliminaries

Very often used *incomplete Gamma functions*  $\Gamma(p, v)$  and  $\gamma(p, v)$  are represented in the following way:

$$\gamma(p, v) := \int_0^v e^{-t} t^{p-1} dt, \quad (\Re(p) > 0; v \geq 0), \quad (1)$$

and

$$\Gamma(p, v) := \int_v^\infty e^{-t} t^{p-1} dt, \quad (v \geq 0; \Re(p) > 0 \text{ when } v = 0). \quad (2)$$

The *incomplete Gamma functions*  $\Gamma(p, v)$  and  $\gamma(p, v)$  are holding the subsequent decomposition relation

$$\Gamma(p, v) + \gamma(p, v) = \Gamma(p), \quad (\Re(p) > 0). \quad (3)$$

The condition that we have employed on the parameter  $v$  and any place of the present study is unrestrained  $\Re(w)$  ( $w \in \mathbb{C}$ ).

We recall here incomplete  $H$ -functions (IHF's)  $\gamma_{p,q}^{m,n}(w)$  and  $\Gamma_{p,q}^{m,n}(w)$  which was introduced by Srivastava et al. (2018), Equations (2.1)-(2.4) in the following manner:

$$\begin{aligned} \Gamma_{p,q}^{m,n}(w) &= \Gamma_{p,q}^{m,n} \left[ w \left| \begin{array}{l} (g_1, \mathcal{G}_1, u), (g_j, \mathcal{G}_j)_{2,p} \\ (h_j, H_j)_{1,q} \end{array} \right. \right] \\ &= \Gamma_{p,q}^{m,n} \left[ w \left| \begin{array}{l} (g_1, \mathcal{G}_1, u), (g_2, \mathcal{G}_2), \dots, (g_p, \mathcal{G}_p) \\ (h_1, H_1), (h_2, H_2), \dots, (h_q, H_q) \end{array} \right. \right] \\ &= \frac{1}{2\pi i} \int_{\mathfrak{L}} f(\xi, u) w^{-\xi} d\xi, \end{aligned} \quad (4)$$

where

$$\begin{aligned}
 f(\xi, u) &= \frac{\Gamma(1 - g_1 - \mathcal{G}_1\xi, u) \prod_{j=1}^m \Gamma(h_j + H_j\xi) \prod_{j=2}^n \Gamma(1 - g_j - \mathcal{G}_j\xi)}{\prod_{j=m+1}^q \Gamma(1 - h_j - H_j\xi) \prod_{j=n+1}^p \Gamma(g_j + \mathcal{G}_j\xi)}, \\
 \gamma_{p,q}^{m,n}(w) &= \gamma_{p,q}^{m,n} \left[ w \left| \begin{array}{l} (g_1, \mathcal{G}_1, u), (g_j, \mathcal{G}_j)_{2,p} \\ (h_j, H_j)_{1,q} \end{array} \right. \right] \\
 &= \gamma_{p,q}^{m,n} \left[ w \left| \begin{array}{l} (g_1, \mathcal{G}_1, u), (g_2, \mathcal{G}_2), \dots, (g_p, \mathcal{G}_p) \\ (h_1, H_1), (h_2, H_2), \dots, (h_q, H_q) \end{array} \right. \right] \\
 &= \frac{1}{2\pi i} \int_{\mathcal{E}} F(\xi, u) w^{-\xi} d\xi, \tag{5}
 \end{aligned}$$

and

$$F(\xi, u) = \frac{\Gamma(1 - g_1 - \mathcal{G}_1\xi, u) \prod_{j=1}^m \Gamma(h_j + H_j\xi) \prod_{j=2}^n \Gamma(1 - g_j - \mathcal{G}_j\xi)}{\prod_{j=m+1}^q \Gamma(1 - h_j - H_j\xi) \prod_{j=n+1}^p \Gamma(g_j + \mathcal{G}_j\xi)}.$$

If  $u \geq 0$ , then the IHF's  $\Gamma_{p,q}^{m,n}(w)$  and  $\gamma_{p,q}^{m,n}(w)$  are exist and conditions of IHF's are stated in various articles given by Kilbas et al. (2006), Mathai and Saxena (1978), and Mathai et al. (2009). A large number of special cases of IHF's are presented in the articles Srivastava et al. (2018), Bansal, Kumar, Khan et al. (2019), Bansal and Choi (2019), and Bansal et al. (2020).

Next, we call here Euler's Beta and Mellin Transform of IHF which was given by Srivastava et al. (2018) due to evaluate some integrals in Section 2.

**Euler's Beta Transform**

The *Euler's Beta* transform of incomplete H-function is given by Srivastava et al. (2018), p. 124, Equation (3.12) in the following way:

$$\begin{aligned}
 &\int_0^t w^{-\alpha} (t - w)^{\alpha - \beta - 1} \Gamma_{p,q}^{m,n} \left[ w \left| \begin{array}{l} (g_1, \mathcal{G}_1, u), (g_j, \mathcal{G}_j)_{2,p} \\ (h_j, H_j)_{1,q} \end{array} \right. \right] dw \\
 &= \frac{\Gamma(\alpha - \beta)}{t^\beta} \Gamma_{p+1,q+1}^{m,n+1} \left[ t \left| \begin{array}{l} (g_1, \mathcal{G}_1, u), (\alpha, 1), (g_j, \mathcal{G}_j)_{2,p} \\ (h_j, H_j)_{1,q}, (\beta, 1) \end{array} \right. \right], \tag{6}
 \end{aligned}$$

provided that conditions are given in Srivastava et al. (2018).

## Mellin Transform

The Mellin transform of incomplete H-function is given by Srivastava et al. (2018), p. 122, Equation (3.4) in the following way:

$$\begin{aligned} & \mathfrak{M} \left\{ \Gamma_{p,q}^{m,n} \left[ \begin{matrix} (\mathfrak{g}_1, \mathcal{G}_1, \mathfrak{u}), (\mathfrak{g}_j, \mathcal{G}_j)_{2,p} \\ (\mathfrak{h}_j, \mathfrak{H}_j)_{1,q} \end{matrix} \right] ; \mathfrak{s} \right\} \\ &= \mathfrak{w}^{-\mathfrak{s}} \frac{\Gamma(1 - \mathfrak{g}_1 - \mathcal{G}_1 \mathfrak{s}, \mathfrak{u}) \prod_{j=1}^m \Gamma(\mathfrak{h}_j + \mathfrak{H}_j \mathfrak{s}) \prod_{j=2}^n \Gamma(1 - \mathfrak{g}_j - \mathcal{G}_j \mathfrak{s})}{\prod_{j=m+1}^q \Gamma(1 - \mathfrak{h}_j - \mathfrak{H}_j \mathfrak{s}) \prod_{j=n+1}^p \Gamma(\mathfrak{g}_j + \mathcal{G}_j \mathfrak{s})}, \end{aligned} \quad (7)$$

provided that conditions are given in Srivastava et al. (2018).

## 3. Main Results

In this section, we evaluate several finite and infinite integrals associated with family of incomplete H-functions (4) and (5).

### Theorem 3.1.

If

$$\begin{aligned} & \lambda > 0, \quad \mathfrak{u} \geq 0, \quad \mathfrak{v} \geq 0, \\ & \lambda \max_{1 \leq j \leq n} \Re \left\{ \frac{\mathfrak{g}_j - 1}{\mathcal{G}_j} \right\} - \min_{1 \leq j \leq M} \Re \left\{ \frac{b_j}{B_j} \right\} < \Re(\alpha) < \lambda \min_{1 \leq j \leq m} \Re \left\{ \frac{\mathfrak{h}_j}{\mathfrak{H}_j} \right\} + \max_{1 \leq j \leq N} \Re \left\{ \frac{1 - a_j}{A_j} \right\}, \end{aligned}$$

then the following improper integral holds:

$$\begin{aligned} & \int_0^\infty t^{\alpha-1} \Gamma_{p,q}^{m,n} \left[ \begin{matrix} (\mathfrak{g}_1, \mathcal{G}_1, \mathfrak{u}), (\mathfrak{g}_j, \mathcal{G}_j)_{2,p} \\ (\mathfrak{h}_j, \mathfrak{H}_j)_{1,q} \end{matrix} \right] \Gamma_{P,Q}^{M,N} \left[ \begin{matrix} (a_1, A_1, \mathfrak{v}), (a_j, A_j)_{2,P} \\ (b_j, B_j)_{1,Q} \end{matrix} \right] dt \\ &= \mathfrak{s}^{-\alpha} \Gamma_{p+P, q+Q}^{m+M, n+N} \left[ \begin{matrix} A^* \\ \mathfrak{w} \mathfrak{s}^\lambda \\ B^* \end{matrix} \right], \end{aligned} \quad (8)$$

and

$$\int_0^\infty t^{\alpha-1} \gamma_{p,q}^{m,n} \left[ \begin{matrix} (g_1, G_1, u), (g_j, G_j)_{2,p} \\ (h_j, H_j)_{1,q} \end{matrix} \right] \gamma_{P,Q}^{M,N} \left[ \begin{matrix} (a_1, A_1, v), (a_j, A_j)_{2,P} \\ (b_j, B_j)_{1,Q} \end{matrix} \right] dt$$

$$= s^{-\alpha} \gamma_{p+P, q+Q}^{m+M, n+N} \left[ \begin{matrix} A^* \\ ws^\lambda \\ B^* \end{matrix} \right], \tag{9}$$

where

$$A^* = (g_1, G_1, u), (a_1 + \alpha A_1, \lambda A_1, v), (g_j, G_j)_{2,n}, (a_j + \alpha A_j, \lambda A_j)_{2,P}, (g_j, G_j)_{n+1,p},$$

$$B^* = (h_j, H_j)_{1,m}, (b_j + \alpha B_j, \lambda B_j)_{1,Q}, (h_j, H_j)_{m+1,q},$$

provided that conditions of IHF's  $\Gamma_{p,q}^{m,n}(w)$  and  $\gamma_{p,q}^{m,n}(w)$  in (4) and (5) are satisfied.

**Proof:**

To prove the assertion (8), firstly, we write the Mellin-Barnes contour integral form of IHF with the help of (4), we get (say  $\Delta$ )

$$\Delta = \int_0^\infty t^{\alpha-1} \Gamma_{P,Q}^{M,N} \left[ \begin{matrix} (a_1, A_1, v), (a_j, A_j)_{2,P} \\ (b_j, B_j)_{1,Q} \end{matrix} \right] \left[ \frac{1}{2\pi i} \int_{\mathcal{L}} f(\xi, u) (wt^{-\lambda})^{-\xi} d\xi \right] dt.$$

Further, changing the order of integration and with the help of (7), we obtain

$$\Delta = \frac{s^{-\alpha}}{2\pi i} \int_{\mathcal{L}} f(\xi, u) (ws^\lambda)^{-\xi}$$

$$\times \frac{\Gamma(1 - a_1 - A_1(\alpha + \lambda\xi)) \prod_{j=1}^M \Gamma(b_j + B_j(\alpha + \lambda\xi)) \prod_{j=2}^N \Gamma(1 - a_j - A_j(\alpha + \lambda\xi))}{\prod_{j=M+1}^Q \Gamma(1 - b_j - B_j(\alpha + \lambda\xi)) \prod_{j=N+1}^P \Gamma(a_j + A_j(\alpha + \lambda\xi))} d\xi.$$

With the help of Equation (4), we get the required result. ■

**Theorem 3.2.**

If

$$\lambda > 0, \qquad \beta > 0, \qquad u \geq 0,$$

$$-\lambda \min_{1 \leq j \leq m} \Re \left\{ \frac{h_j}{H_j} \right\} < \Re(\alpha) < \lambda \max_{1 \leq j \leq n} \Re \left\{ \frac{1 - g_j}{G_j} \right\},$$

then the following improper integral holds:

$$\int_0^\infty t^{\alpha-1} (t + \mathbf{a})^{-\beta} \Gamma_{p,q}^{m,n} \left[ \begin{matrix} \mathbf{wt}^\lambda \\ (h_j, H_j)_{1,q} \end{matrix} \middle| \begin{matrix} (g_1, \mathcal{G}_1, \mathbf{u}), (g_j, \mathcal{G}_j)_{2,p} \end{matrix} \right] dt$$

$$= \frac{\mathbf{a}^{\alpha-\beta}}{\Gamma(\beta)} \Gamma_{p+1,q+1}^{m+1,n+1} \left[ \begin{matrix} \mathbf{wa}^\lambda \\ (\beta - \alpha, \lambda), (h_j, H_j)_{1,q} \end{matrix} \middle| \begin{matrix} (g_1, \mathcal{G}_1, \mathbf{u}), (1 - \alpha, \lambda), (g_j, \mathcal{G}_j)_{2,p} \end{matrix} \right], \quad (10)$$

and

$$\int_0^\infty t^{\alpha-1} (t + \mathbf{a})^{-\beta} \gamma_{p,q}^{m,n} \left[ \begin{matrix} \mathbf{wt}^\lambda \\ (h_j, H_j)_{1,q} \end{matrix} \middle| \begin{matrix} (g_1, \mathcal{G}_1, \mathbf{u}), (g_j, \mathcal{G}_j)_{2,p} \end{matrix} \right] dt$$

$$= \frac{\mathbf{a}^{\alpha-\beta}}{\Gamma(\beta)} \gamma_{p+1,q+1}^{m+1,n+1} \left[ \begin{matrix} \mathbf{wa}^\lambda \\ (\beta - \alpha, \lambda), (h_j, H_j)_{1,q} \end{matrix} \middle| \begin{matrix} (g_1, \mathcal{G}_1, \mathbf{u}), (1 - \alpha, \lambda), (g_j, \mathcal{G}_j)_{2,p} \end{matrix} \right], \quad (11)$$

provided that condition of incomplete  $H$ -functions  $\Gamma_{p,q}^{m,n}(z)$  and  $\gamma_{p,q}^{m,n}(z)$  in (4) and (5) are satisfied.

**Proof:**

To prove the result (10), firstly, we write the Mellin-Barnes contour integral form of IHF with the help of (4), we get (say  $\Omega$ )

$$\Omega = \int_0^\infty t^{\alpha-1} (t + \mathbf{a})^{-\beta} \left[ \frac{1}{2\pi i} \int_{\mathcal{L}} f(\xi, \mathbf{u}) (\mathbf{wt}^\lambda)^{-\xi} d\xi \right] dt.$$

Further, changing the order of integration and with the help of Beta function definition, we obtained

$$\Omega = \frac{\mathbf{a}^{\alpha-\beta}}{\Gamma(\beta)} \frac{1}{2\pi i} \int_{\mathcal{L}} f(\xi, \mathbf{u}) \Gamma(\alpha - \lambda\xi) \Gamma(\beta - \alpha + \lambda\xi) (\mathbf{wa}^\lambda)^{-\xi} d\xi.$$

With the help of equation (4), we get the required result. ■

**Theorem 3.3.**

If

$$\lambda > 0, \quad \mu > 0, \quad \mathbf{u} \geq 0,$$

$$\Re(\alpha) + \lambda \min_{1 \leq j \leq m} \Re\left(\frac{h_j}{H_j}\right) > 0, \quad \Re(\beta) + \mu \min_{1 \leq j \leq m} \Re\left(\frac{h_j}{H_j}\right) > 0,$$

then the following improper integral is hold:

$$\int_0^t x^{\alpha-1}(t-x)^{\beta-1} \Gamma_{p,q}^{m,n} \left[ \begin{matrix} wx^\lambda(t-x)^\mu \\ (h_j, H_j)_{1,q} \end{matrix} \middle| \begin{matrix} (g_1, \mathcal{G}_1, u), (g_j, \mathcal{G}_j)_{2,p} \end{matrix} \right] dx$$

$$= t^{\alpha+\beta-1} \Gamma_{p+2,q+1}^{m,n+2} \left[ \begin{matrix} wt^{\lambda+\mu} \\ (\beta-\alpha, \lambda), (h_j, H_j)_{1,q}, (1-\alpha-\beta, \lambda+\mu) \end{matrix} \middle| \begin{matrix} (g_1, \mathcal{G}_1, u), (1-\alpha, \lambda), (1-\beta, \mu), (g_j, \mathcal{G}_j)_{2,p} \end{matrix} \right], \tag{12}$$

and

$$\int_0^t x^{\alpha-1}(t-x)^{\beta-1} \gamma_{p,q}^{m,n} \left[ \begin{matrix} wx^\lambda(t-x)^\mu \\ (h_j, H_j)_{1,q} \end{matrix} \middle| \begin{matrix} (g_1, \mathcal{G}_1, u), (g_j, \mathcal{G}_j)_{2,p} \end{matrix} \right] dx$$

$$= t^{\alpha+\beta-1} \gamma_{p+2,q+1}^{m,n+2} \left[ \begin{matrix} wt^{\lambda+\mu} \\ (\beta-\alpha, \lambda), (h_j, H_j)_{1,q}, (1-\alpha-\beta, \lambda+\mu) \end{matrix} \middle| \begin{matrix} (g_1, \mathcal{G}_1, u), (1-\alpha, \lambda), (1-\beta, \mu), (g_j, \mathcal{G}_j)_{2,p} \end{matrix} \right], \tag{13}$$

provided that the condition of incomplete  $H$ -functions  $\Gamma_{p,q}^{m,n}(w)$  and  $\gamma_{p,q}^{m,n}(w)$  in (4) and (5) are satisfied.

**Proof:**

To prove the assertion (12), firstly, we write the Mellin-Barnes contour integral form of IHF with the help of (4), we get (say  $\Xi$ )

$$\Xi = \int_0^t x^{\alpha-1}(t-x)^{-\beta} \left[ \frac{1}{2\pi i} \int_{\mathfrak{L}} f(\xi, u) (wx^\lambda(t-x)^\mu)^{-\xi} d\xi \right] dt.$$

Further, changing the order of integration and with the help of Beta function definition, we obtained

$$\Xi = \frac{t^{\alpha+\beta-1}}{2\pi i} \int_{\mathfrak{L}} f(\xi, u) \frac{\Gamma(\alpha - \lambda\xi)\Gamma(\beta - \mu\xi)}{\Gamma(\alpha + \beta - (\lambda + \mu)\xi)} (wt^{\lambda+\mu})^{-\xi} d\xi.$$

With the help of (4), we get the required result. ■

**Theorem 3.4.**

If

$$\rho > 0, \quad \sigma > 0, \quad u \geq 0, \quad v \geq 0, \quad \lambda + \mu \geq 1$$

$$\Re(\alpha) + \lambda \min_{1 \leq j \leq m} \Re\left(\frac{h_j}{H_j}\right) > 0, \quad \Re(\beta) + \mu \min_{1 \leq j \leq m} \Re\left(\frac{h_j}{H_j}\right) > 0,$$

then the following improper integral is hold:



$$\begin{aligned}
& \int_0^t x^{\alpha-1} (t-x)^{\beta-1} {}_P\Gamma_Q \left[ \begin{matrix} (a_1, \mathbf{u}), a_2, \dots, a_P; \\ b_1, \dots, b_Q; \end{matrix} \alpha x^\lambda (t-x)^\mu \right] \\
& \Gamma_{p,q}^{m,n} \left[ \mathbf{w}x^\rho (t-x)^\sigma \left| \begin{matrix} (\mathbf{g}_1, \mathcal{G}_1, \mathbf{v}), (\mathbf{g}_j, \mathcal{G}_j)_{2,p} \\ (\mathbf{h}_j, \mathbf{H}_j)_{1,q} \end{matrix} \right. \right] dx = t^{\alpha+\beta-1} \frac{\prod_{j=1}^Q \Gamma(b_j)}{\prod_{j=1}^P \Gamma(a_j)} \\
& \times \sum_{r=0}^{\infty} f(r) \Gamma_{p+2,q+1}^{m,n+2} \left[ \mathbf{w}t^{\rho+\sigma} \left| \begin{matrix} (\mathbf{g}_1, \mathcal{G}_1, \mathbf{u}), (1-\alpha-\lambda r, \rho), (1-\beta-\mu r, \sigma), (\mathbf{g}_j, \mathcal{G}_j)_{2,p} \\ (\mathbf{h}_j, \mathbf{H}_j)_{1,q}, (1-\alpha-\beta-(\mu+\lambda)r, \rho+\sigma) \end{matrix} \right. \right] t^{(\lambda+\mu)r},
\end{aligned} \tag{14}$$

where

$$\begin{aligned}
f(r) &= \frac{\Gamma(a_1+r, \mathbf{u}) \prod_{j=2}^P \Gamma(a_j+r)}{\prod_{j=1}^Q \Gamma(b_j+r)} \frac{\mathbf{a}^r}{r!}, \\
& \int_0^t x^{\alpha-1} (t-x)^{\beta-1} {}_P\gamma_Q \left[ \begin{matrix} (a_1, \mathbf{u}), a_2, \dots, a_P; \\ b_1, \dots, b_Q; \end{matrix} \alpha x^\lambda (t-x)^\mu \right] \\
& \gamma_{p,q}^{m,n} \left[ \mathbf{w}x^\rho (t-x)^\sigma \left| \begin{matrix} (\mathbf{g}_1, \mathcal{G}_1, \mathbf{v}), (\mathbf{g}_j, \mathcal{G}_j)_{2,p} \\ (\mathbf{h}_j, \mathbf{H}_j)_{1,q} \end{matrix} \right. \right] dx = t^{\alpha+\beta-1} \frac{\prod_{j=1}^Q \Gamma(b_j)}{\prod_{j=1}^P \Gamma(a_j)} \\
& \times \sum_{r=0}^{\infty} g(r) \gamma_{p+2,q+1}^{m,n+2} \left[ \mathbf{w}t^{\rho+\sigma} \left| \begin{matrix} (\mathbf{g}_1, \mathcal{G}_1, \mathbf{u}), (1-\alpha-\lambda r, \rho), (1-\beta-\mu r, \sigma), (\mathbf{g}_j, \mathcal{G}_j)_{2,p} \\ (\mathbf{h}_j, \mathbf{H}_j)_{1,q}, (1-\alpha-\beta-(\mu+\lambda)r, \rho+\sigma) \end{matrix} \right. \right] t^{(\lambda+\mu)r},
\end{aligned} \tag{15}$$

and

$$g(r) = \frac{\gamma(a_1+r, \mathbf{u}) \prod_{j=2}^P \Gamma(a_j+r)}{\prod_{j=1}^Q \Gamma(b_j+r)} \frac{\mathbf{a}^r}{r!},$$

provided that the condition of incomplete  $H$ -functions  $\Gamma_{p,q}^{m,n}(\mathbf{w})$  and  $\gamma_{p,q}^{m,n}(\mathbf{w})$  in (4) and (5) are satisfied.

**Proof:**

To prove the assertion (14), firstly, we express the incomplete generalized hypergeometric function in series form, we get (say  $\Upsilon$ )

$$\Upsilon = \int_0^t x^{\alpha-1} (t-x)^{\beta-1} \left[ \frac{\prod_{j=1}^Q \Gamma(b_j)}{\prod_{j=1}^P \Gamma(a_j)} \sum_{r=0}^{\infty} \frac{\Gamma(a_1+r, u) \prod_{j=2}^P \Gamma(a_j+r)}{\prod_{j=1}^Q \Gamma(b_j+r)} \frac{(ax^\lambda(t-x)^\mu)^r}{r!} \right] \\ \times \Gamma_{p,q}^{m,n} \left[ wx^\rho(t-x)^\sigma \left| \begin{array}{c} (\mathfrak{g}_1, \mathcal{G}_1, \nu), (\mathfrak{g}_j, \mathcal{G}_j)_{2,p} \\ (\mathfrak{h}_j, \mathbb{H}_j)_{1,q} \end{array} \right. \right] dx.$$

Further, changing the order of integration and summation (under the permissible conditions) and with the help of (12), we obtain

$$\Upsilon = \frac{\prod_{j=1}^Q \Gamma(b_j)}{\prod_{j=1}^P \Gamma(a_j)} \sum_{r=0}^{\infty} \frac{\Gamma(a_1+r, u) \prod_{j=2}^P \Gamma(a_j+r)}{\prod_{j=1}^Q \Gamma(b_j+r)} \frac{a^r}{r!} \\ \times \int_0^t x^{\alpha+\lambda r-1} (t-x)^{\beta+\mu r-1} \Gamma_{p,q}^{m,n} \left[ wx^\rho(t-x)^\sigma \left| \begin{array}{c} (\mathfrak{g}_1, \mathcal{G}_1, \nu), (\mathfrak{g}_j, \mathcal{G}_j)_{2,p} \\ (\mathfrak{h}_j, \mathbb{H}_j)_{1,q} \end{array} \right. \right] dx.$$

Next, with the help of (12), we get the required result after a little simplification. ■

**Theorem 3.5.**

$$\lambda > 0, \quad \mu > 0, \quad u \geq 0, \\ \Re(\alpha) - \lambda \max_{1 \leq j \leq n} \Re \left\{ \frac{\mathfrak{g}_j - 1}{\mathcal{G}_j} \right\} > 0, \quad \Re(\beta) - \mu \max_{1 \leq j \leq n} \Re \left\{ \frac{\mathfrak{g}_j - 1}{\mathcal{G}_j} \right\} > 0,$$

then the following improper integral is hold:

$$\begin{aligned} & \int_0^t x^{\alpha-1}(t-x)^{\beta-1} \Gamma_{p,q}^{m,n} \left[ \left. \begin{matrix} wx^{-\lambda}(t-x)^{-\mu} \\ (\mathbf{g}_1, \mathcal{G}_1, \mathbf{u}), (\mathbf{g}_j, \mathcal{G}_j)_{2,p} \\ (\mathbf{h}_j, \mathbf{H}_j)_{1,q} \end{matrix} \right| \right] dx \\ &= t^{\alpha+\beta-1} \Gamma_{p+1,q+2}^{m+2,n} \left[ \left. \begin{matrix} wt^{-\lambda-\mu} \\ (\mathbf{g}_1, \mathcal{G}_1, \mathbf{u}), (\mathbf{g}_j, \mathcal{G}_j)_{2,p}, (\alpha + \beta, \mu + \lambda) \\ (\alpha, \lambda), (\beta, \mu), (\mathbf{h}_j, \mathbf{H}_j)_{1,q} \end{matrix} \right| \right], \end{aligned} \quad (16)$$

and

$$\begin{aligned} & \int_0^t x^{\alpha-1}(t-x)^{\beta-1} \gamma_{p,q}^{m,n} \left[ \left. \begin{matrix} wx^{-\lambda}(t-x)^{-\mu} \\ (\mathbf{g}_1, \mathcal{G}_1, \mathbf{u}), (\mathbf{g}_j, \mathcal{G}_j)_{2,p} \\ (\mathbf{h}_j, \mathbf{H}_j)_{1,q} \end{matrix} \right| \right] dx \\ &= t^{\alpha+\beta-1} \gamma_{p+1,q+2}^{m+2,n} \left[ \left. \begin{matrix} wt^{-\lambda-\mu} \\ (\mathbf{g}_1, \mathcal{G}_1, \mathbf{u}), (\mathbf{g}_j, \mathcal{G}_j)_{2,p}, (\alpha + \beta, \mu + \lambda) \\ (\alpha, \lambda), (\beta, \mu), (\mathbf{h}_j, \mathbf{H}_j)_{1,q} \end{matrix} \right| \right], \end{aligned} \quad (17)$$

provided that the condition of incomplete  $H$ -functions  $\Gamma_{p,q}^{m,n}(w)$  and  $\gamma_{p,q}^{m,n}(w)$  in (4) and (5) are satisfied.

**Proof:**

To prove the assertion (16), firstly, we write the Mellin-Barnes contour integral form of IHF with the help of (4), we get (say  $\Xi$ )

$$\Xi = \int_0^t x^{\alpha-1}(t-x)^{-\beta} \left[ \frac{1}{2\pi i} \int_{\mathfrak{L}} f(\xi, \mathbf{u}) (wx^{-\lambda}(t-x)^{-\mu})^{-\xi} d\xi \right] dt.$$

Further, changing the order of integration and with the help of Beta function definition, we obtained

$$\Xi = \frac{t^{\alpha+\beta-1}}{2\pi i} \int_{\mathfrak{L}} f(\xi, \mathbf{u}) \frac{\Gamma(\alpha + \lambda\xi)\Gamma(\beta + \mu\xi)}{\Gamma(\alpha + \beta + (\lambda + \mu)\xi)} (wt^{-\lambda-\mu})^{-\xi} d\xi.$$

With the help of equation (4), we get the required result. ■

**Theorem 3.6.**

$$\begin{aligned} \rho > 0, \quad \sigma > 0, \quad \mathbf{u} \geq 0, \quad \mathbf{v} \geq 0, \quad \lambda + \mu \geq 1, \\ \Re(\alpha) - \rho \max_{1 \leq j \leq n} \Re \left\{ \frac{\mathbf{g}_j - 1}{\mathcal{G}_j} \right\} > 0, \quad \Re(\beta) - \sigma \max_{1 \leq j \leq n} \Re \left\{ \frac{\mathbf{g}_j - 1}{\mathcal{G}_j} \right\} > 0, \end{aligned}$$

then the following improper integral is hold:

$$\int_0^t x^{\alpha-1}(t-x)^{\beta-1} {}_P\Gamma_Q \left[ \begin{matrix} (a_1, \mathbf{u}), a_2, \dots, a_P; \\ b_1, \dots, b_Q; \end{matrix} \mathbf{a}x^\lambda(t-x)^\mu \right] \\ \Gamma_{p,q}^{m,n} \left[ \mathbf{w}x^{-\rho}(t-x)^{-\sigma} \left| \begin{matrix} (\mathbf{g}_1, \mathcal{G}_1, \mathbf{v}), (\mathbf{g}_j, \mathcal{G}_j)_{2,p} \\ (\mathbf{h}_j, \mathbf{H}_j)_{1,q} \end{matrix} \right. \right] dx = t^{\alpha+\beta-1} \frac{\prod_{j=1}^Q \Gamma(b_j)}{\prod_{j=1}^P \Gamma(a_j)} \\ \times \sum_{r=0}^{\infty} f(r) \Gamma_{p+1,q+2}^{m+2,n} \left[ \mathbf{w}t^{-\rho-\sigma} \left| \begin{matrix} (\mathbf{g}_1, \mathcal{G}_1, \mathbf{v}), (\mathbf{g}_j, \mathcal{G}_j)_{2,p}, (\alpha + \beta + (\mu + \lambda)r, \rho + \sigma) \\ (\alpha + \lambda r, \rho), (\beta + \mu r, \sigma), (\mathbf{h}_j, \mathbf{H}_j)_{1,q}, \end{matrix} \right. \right] t^{(\lambda+\mu)r}, \quad (18)$$

where

$$f(r) = \frac{\Gamma(a_1 + r, \mathbf{u}) \prod_{j=2}^P \Gamma(a_j + r)}{\prod_{j=1}^Q \Gamma(b_j + r)} \frac{\mathbf{a}^r}{r!},$$

and

$$\int_0^t x^{\alpha-1}(t-x)^{\beta-1} {}_P\gamma_Q \left[ \begin{matrix} (a_1, \mathbf{u}), a_2, \dots, a_P; \\ b_1, \dots, b_Q; \end{matrix} \mathbf{a}x^\lambda(t-x)^\mu \right] \\ \gamma_{p,q}^{m,n} \left[ \mathbf{w}x^{-\rho}(t-x)^{-\sigma} \left| \begin{matrix} (\mathbf{g}_1, \mathcal{G}_1, \mathbf{v}), (\mathbf{g}_j, \mathcal{G}_j)_{2,p} \\ (\mathbf{h}_j, \mathbf{H}_j)_{1,q} \end{matrix} \right. \right] dx = t^{\alpha+\beta-1} \frac{\prod_{j=1}^Q \Gamma(b_j)}{\prod_{j=1}^P \Gamma(a_j)} \\ \times \sum_{r=0}^{\infty} g(r) \cdot \gamma_{p+1,q+2}^{m+2,n} \left[ \mathbf{w}t^{-\rho-\sigma} \left| \begin{matrix} (\mathbf{g}_1, \mathcal{G}_1, \mathbf{v}), (\mathbf{g}_j, \mathcal{G}_j)_{2,p}, (\alpha + \beta + (\mu + \lambda)r, \rho + \sigma) \\ (\alpha + \lambda r, \rho), (\beta + \mu r, \sigma), (\mathbf{h}_j, \mathbf{H}_j)_{1,q}, \end{matrix} \right. \right] t^{(\lambda+\mu)r}, \quad (19)$$

where

$$g(r) = \frac{\Gamma(a_1 + r, \mathbf{u}) \prod_{j=2}^P \Gamma(a_j + r)}{\prod_{j=1}^Q \Gamma(b_j + r)} \frac{\mathbf{a}^r}{r!},$$

provided that the condition of incomplete  $H$ -functions  $\Gamma_{p,q}^{m,n}(\mathbf{w})$  and  $\gamma_{p,q}^{m,n}(\mathbf{w})$  in (4) and (5) are satisfied.

**Proof:**

To prove the assertion (18), firstly, we express the incomplete generalized hypergeometric function in series form, we get (say  $\Upsilon$ )

$$\Upsilon = \int_0^t x^{\alpha-1} (t-x)^{\beta-1} \left[ \frac{\prod_{j=1}^Q \Gamma(b_j)}{\prod_{j=1}^P \Gamma(a_j)} \sum_{r=0}^{\infty} \frac{\Gamma(a_1+r, u) \prod_{j=2}^P \Gamma(a_j+r)}{\prod_{j=1}^Q \Gamma(b_j+r)} \frac{(ax^\lambda(t-x)^\mu)^r}{r!} \right] \\ \times \Gamma_{p,q}^{m,n} \left[ wx^{-\rho} (t-x)^{-\sigma} \left| \begin{array}{c} (\mathfrak{g}_1, \mathcal{G}_1, \nu), (\mathfrak{g}_j, \mathcal{G}_j)_{2,p} \\ (\mathfrak{h}_j, \mathbf{H}_j)_{1,q} \end{array} \right. \right] dx.$$

Further, changing the order of integration and summation (under the permissible conditions) and with the help of (12), we obtained

$$\Upsilon = \frac{\prod_{j=1}^Q \Gamma(b_j)}{\prod_{j=1}^P \Gamma(a_j)} \sum_{r=0}^{\infty} \frac{\Gamma(a_1+r, u) \prod_{j=2}^P \Gamma(a_j+r)}{\prod_{j=1}^Q \Gamma(b_j+r)} \frac{a^r}{r!} \\ \times \int_0^t x^{\alpha+\lambda r-1} (t-x)^{\beta+\mu r-1} \Gamma_{p,q}^{m,n} \left[ wx^{-\rho} (t-x)^{-\sigma} \left| \begin{array}{c} (\mathfrak{g}_1, \mathcal{G}_1, \nu), (\mathfrak{g}_j, \mathcal{G}_j)_{2,p} \\ (\mathfrak{h}_j, \mathbf{H}_j)_{1,q} \end{array} \right. \right] dx.$$

Next, with the help of (16), we get the required result after a little simplification. ■

#### 4. Integral representation of incomplete H-functions

Recently, Srivastava et al. (2012) established the integral representation of the incomplete Gauss hypergeometric functions. So, motivated by the work of Srivastava et al. (2012), we give the integral representation of incomplete  $H$ -functions (4) and (5).

**Theorem 4.1.**

If  $u \geq 0$  and  $\Re(\mathfrak{g}_1) > 0$ , then the following integral representation formula holds:

$$\Gamma_{p,q}^{m,n} \left[ w \left| \begin{array}{c} (1 - \mathfrak{g}_1, \mathcal{G}_1, u), (\mathfrak{g}_j, \mathcal{G}_j)_{2,p} \\ (\mathfrak{h}_j, \mathbf{H}_j)_{1,q} \end{array} \right. \right] = \int_u^\infty t^{\mathfrak{g}_1-1} e^{-t} \mathcal{H}_{p-1,q}^{m,n-1} \left[ wt^{\mathfrak{G}_1} \left| \begin{array}{c} (\mathfrak{g}_j, \mathcal{G}_j)_{2,p} \\ (\mathfrak{h}_j, \mathbf{H}_j)_{1,q} \end{array} \right. \right] dt. \quad (20)$$

**Proof:**

To prove the assertion (20), we write the Mellin-Barnes contour integral form of well known  $\mathcal{H}$ -function and then with the help of (2), we get the desired result. ■

**Theorem 4.2.**

If  $u \geq 0$  and  $\Re(g_1) > 0$ , then the following integral representation formula holds:

$$\gamma_{p,q}^{m,n} \left[ w \left| \begin{array}{c} (1 - g_1, \mathcal{G}_1, u), (g_j, \mathcal{G}_j)_{2,p} \\ (h_j, H_j)_{1,q} \end{array} \right. \right] = \int_0^u t^{g_1-1} e^{-t} \mathcal{H}_{p-1,q}^{m,n-1} \left[ wt^{g_1} \left| \begin{array}{c} (g_j, \mathcal{G}_j)_{2,p} \\ (h_j, H_j)_{1,q} \end{array} \right. \right] dt. \quad (21)$$

**Proof:**

To prove the assertion (21), we write the Mellin-Barnes contour integral form of well known  $\mathcal{H}$ -function and then with the help of (1), we get the desired result. ■

**Remark 4.3.**

If incomplete  $H$ -function reduces to the familiar of Fox's  $\mathcal{H}$ -function and incomplete generalised hypergeometric function reduce into generalised hypergeometric function, then results are recorded in the text book of Srivastava et al. (1982).

**5. Conclusions**

Many authors produced a large number of research articles on the integral formulas involving the various kind of special functions. In this regard, we deal with several finite and infinite integrals involving the family of incomplete  $H$ -functions(IHF's). Further, we point out some known and new special cases of these integrals. Finally, we established the integral representation of IHF's.

**Acknowledgment:**

*The author K.S. Nisar expresses his thanks to the Deanship of Scientific Research (DSR), Prince Sattam bin Abdulaziz University, Saudi Arabia for providing facilities and support.*

**REFERENCES**

Bansal, M.K. and Choi, J. (2019). A note on pathway fractional integral formulas associated with the incomplete  $H$ -functions, Int. J. Appl. Comput. Math, Vol. 5, No. 5, pp. 1–10.

- Bansal, M.K., Kumar, D. and Jain, R. (2019a). Interrelationships between Marichev-Saigo-Maeda fractional integral operators, the Laplace Transform and the  $\bar{H}$ -function, *Int. J. Appl. Comput. Math*, Vol. 5, No. 4, pp. 1–13.
- Bansal, M.K., Kumar, D. and Jain, R. (2019b). A study of Marichev-Saigo-Maeda fractional integral operators associated with S-generalized Gauss hypergeometric function, *Kyungpook Math. J.*, Vol. 59, No. 3, pp. 433–443.
- Bansal, M.K., Kumar, D., Khan, I., Singh, J. and Nisar, K.S. (2019). Certain unified integrals associated with product of  $M$ -series and incomplete  $H$ -functions, *Mathematics*, Vol. 7, No. 12, pp. 1–11.
- Bansal, M.K., Kumar, D., Singh, J., Tassaddiq, A. and Nisar, K.S. (2020). Some new results for the Srivastava-Luo-Raina M-transform pertaining to the incomplete H-functions, *AIMS-Mathematics*, Vol. 5, No. 1, pp. 717-722.
- Kilbas, A.A., Srivastava, H.M. and Trujillo, J.J. (2006). *Theory and Applications of Fractional Differential Equations* (North-Holland Mathematical Studies), Elsevier (North Holland) Science Publishers, Amsterdam, London and New York.
- Kumar, D., Gupta, R.K., Rawat, D.S. and Daiya, J. (2018). Hypergeometric fractional integrals of multiparameter K-Mittag-Leffler function, *Nonlinear Sci. Lett, A Math. Phys. Mech.*, Vol. 9, No. 1, pp. 17–26.
- Mathai, A.M. and Saxena, R.K. (1978). *The H-function with Applications in Statistics Other Disciplines*, Wiley Eastern, New Delhi & Wiley Halsted, New York.
- Mathai, A.M., Saxena, R.K. and Haubold, H.J. (2009). *The H-function: Theory and Applications*, Springer, New York.
- Saxena, R.K., Purohit, S.D. and Kumar, D. (2018). Integral inequalities associated with Gauss hypergeometric function fractional integral operator, *Proc. Natl. Acad. Sci., India, Sect. A Phys. Sci.*, Vol. 88, No. 1, pp. 27–31.
- Srivastava, H.M., Chaudhary, M.A. and Agarwal, R.P.(2012). The incomplete Pochhammer symbols and their applications to hypergeometric and related functions, *Integral Transforms Spec. Funct.*, Vol. 23, pp. 659–683.
- Srivastava, H.M., Gupta, K.C. and Goyal, S.P. (1982). *The H-Functions of One and Two Variables with Applications*, South Asian Publishers, New Delhi.
- Srivastava, H.M., Saxena, R.K. and Parmar, R.K. (2018). Some Families of the Incomplete  $H$ -functions and the Incomplete  $\bar{H}$ -functions and Associated Integral Transforms and Operators of Fractional Calculus with Applications, *Russ. J. Math. Phys.*, Vol. 25, No. 1, pp. 116–138.
- Suthar, D.L., Agarwal, S. and Kumar, D. (2019). Certain integrals involving the product of Gaussian hypergeometric function and Aleph function, *Honam Math. J.*, Vol. 41, No. 1, pp. 1–17.